

MET in Lung Cancer: Biomarker Selection Based on Scientific Rationale

Ravi Salgia

Abstract

MET or hepatocyte growth factor (HGF) receptor pathway signaling mediates wound healing and hepatic regeneration, with pivotal roles in embryonic, neuronal, and muscle development. However, dysregulation of MET signaling mediates proliferation, apoptosis, and migration and is implicated in a number of malignancies. In non-small cell lung cancer (NSCLC), aberrant MET signaling can occur through a number of mechanisms that collectively represent a significant proportion of patients. These include MET or HGF protein overexpression, *MET* gene amplification, *MET* gene mutation or fusion/rearrangement, or aberrations in downstream signaling or regulatory components. Responses to MET tyrosine kinase inhibitors have been documen-

ted in clinical trials in patients with *MET*-amplified or *MET*-overexpressing NSCLC, and case studies or case series have shown that *MET* mutation/deletion is a biomarker that is also predictive of response to these agents. However, other recent clinical data have highlighted an urgent need to elucidate optimal biomarkers based on genetic and/or protein diagnostics to correctly identify patients most likely to benefit in ongoing clinical trials of an array of *MET*-targeted therapies of differing class. The latest advances in the development of *MET* biomarkers in NSCLC have been reviewed, toward establishing appropriate *MET* biomarker selection based on a scientific rationale. *Mol Cancer Ther*; 16(4); 555–65. ©2017 AACR.

Introduction

The *MET* proto-oncogene was originally identified as a fusion partner with the translocated promoter region of the *TPR* gene in a chemically transformed osteosarcoma-derived cell line (1). The *MET* protein encoded by this proto-oncogene was later found to be a transmembrane receptor tyrosine kinase (RTK) activated by an endogenous ligand, scatter factor, or hepatocyte growth factor (HGF; refs. 2–5). Binding of HGF to *MET* [or HGF receptor (HGFR)] results in receptor dimerization and phosphorylation of tyrosine residues, ultimately leading to the phosphorylation of intracellular docking sites where adaptor proteins bind to activate downstream signaling (4, 6, 7). Activated signaling pathways include mitogen-activated protein kinase (MAPK), phosphoinositide 3-kinase (PI3K)/AKT (protein kinase B), signal transducer and activator of transcription proteins, and nuclear factor- κ B (8–10). In normal physiology, these signaling pathways promote activation of cytoplasmic and nuclear processes, which lead to a variety of cellular functions, including proliferation and protection from apoptosis (8–10). The *MET* pathway also mediates functions such as wound healing and hepatic regeneration, and has pivotal roles in normal liver development (11), embryonic

placental development, and the formation of muscle and neurons (12–15).

HFG/MET in Lung Cancer

Dysregulation of *MET* signaling-mediated proliferation, apoptosis, and migration through overexpression of *MET* and amplification or mutation of the *MET* gene has been widely demonstrated in oncogenic processes across multiple tumor types and has been reviewed elsewhere (10, 16–18). Moreover, it is notable that all three of these mechanisms of *MET*/*MET* dysregulation have been documented in non-small cell lung cancer (NSCLC; refs. 19–22). Early studies established that *MET* can be overexpressed or activated [as measured by phosphorylation of the catalytic domain as well as juxtamembrane (JM) domain], or the gene mutated (in the semaphorin or JM domains) and/or amplified in lung cancer. For instance, studies on small cell lung cancer (SCLC) cell lines established the multipurpose nature of *MET*/HGF pathway activation during tumor progression and invasion, which occurs via dysregulation of diverse biological functions such as proliferation and differentiation, transcriptional control, cell-cycle G_1/S checkpoint, cytoskeletal functions, survival, motility, and apoptosis (23). Both epidermal growth factor receptor (EGFR) and *MET* are widely expressed on cancer cells, and both RTKs are implicated in these diverse signaling processes. Indeed, synergistic effects of epidermal growth factor (EGF) and HGF on proliferation, and additive effects on motility, were noted in preclinical studies in NSCLC cells. For example, increased membrane ruffling to form a motile cell surface was observed when cells were stimulated with HGF and EGF independently, and when these growth factors were combined, an additive effect was observed (24). These preclinical studies suggested that the combination of inhibitors for *MET* and EGFR RTKs could potentially produce synergistic antitumor effects (24). Indeed, a synergistic effect on inhibition of cell proliferation and apoptosis was

City of Hope, Department of Medical Oncology and Therapeutics Research, Duarte, California.

Note: Supplementary data for this article are available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

Corresponding Author: Ravi Salgia, City of Hope, Department of Medical Oncology and Therapeutics Research, 1500 E. Duarte Road, Duarte, CA 91010-3000. Phone: 626-471-9200; Fax: 626-471-7322; E-mail: rsalgia@coh.org

doi: 10.1158/1535-7163.MCT-16-0472

©2017 American Association for Cancer Research.

seen when a novel first-generation MET inhibitor, SU11274, was combined with EGFR inhibitors such as AG1478 or gefitinib (24). The synergism and cross-talk of EGFR and MET pathways in NSCLC, and the potential of simultaneous inhibition, were thus recognized in these studies.

Therapy combining various targeted agents or other standard therapies with MET inhibitors has also been explored in other preclinical studies, including synergistic effects of when MET and EGFR inhibitors were combined in NSCLC cell lines, and head and neck squamous cell carcinoma (SCCHN) cells (24, 25). Combined inhibition of MET and mammalian target of rapamycin resulted in cooperative inhibition of cell growth in *TPR-MET* transformed cells expressing a constitutively active variant of MET, and in MET-expressing NSCLC cells (26). MET inhibition is also synergistic with cisplatin in SCCHN cancer cell lines (25) and also appears to be synergistic with radiation in cell lines in some studies, although crizotinib (Pfizer), an inhibitor of MET, anaplastic lymphoma kinase (ALK), and ROS1 kinases, failed to enhance the effects of radiation in SCCHN xenograft models (27). Furthermore, MET inhibition was synergistic with topoisomerase-I inhibition in SCLC cell lines, with a significant positive correlation observed between *MET* gene copy number (GCN) and topoisomerase-I nuclear expression (28).

Role of MET and HGF in Induction of Drug Resistance

Several RTKs often expressed on cancer cells activate signaling pathways that converge on common downstream effectors. This "overlap" may, in part, be implicated in resistance to RTK-based treatment, which is commonly observed in cancer patients. Resistance to targeted agents may be mediated by the stroma, and preclinical investigations of growth-factor-driven resistance have shown that increasing levels of ligands with a similar signaling output, such as PI3K or MAPK, may confer innate or acquired resistance to inhibitors of an oncogenic kinase (29, 30).

Loss of drug sensitivity in tumor cells through exposure to RTK ligands in the tumor microenvironment was demonstrated in *BRAF*-mutant melanoma cells, with HGF conferring resistance to the *BRAF* inhibitor PLX4032 (vemurafenib; ref. 30). Stromal secretion of HGF resulted in activation of MET, thereby reactivating the MAPK and PI3K/AKT pathways (30). Consequently, in cell models at least, it is feasible that dual inhibition of RAF and MET can reverse drug resistance (30). These data highlight the redundancy of RTK-transduced signaling in cancer cells and the wide-ranging effects of RTK ligands that lead to innate and acquired resistance, which may potentially be overcome through combinations of targeted agents (29, 30).

MET amplification in NSCLC is implicated in acquired resistance to EGFR inhibitors and has been reported in approximately one-fifth of cases with EGFR inhibitor resistance (31–34). This provides further therapeutic rationale for combinations of MET inhibitors with EGFR inhibitors to treat selected patients with NSCLC.

MET Biomarkers in NSCLC

The varied mechanisms of MET activation in lung cancer, including overexpression of MET and/or its ligand, HGF, and genetic alterations to *MET* (e.g., mutations, amplification, translocation, or dysregulated transcription), and impaired degrada-

tion of MET, provide an array of potential biomarkers (Table 1). The challenge now faced is to identify which of these biomarkers holds the most promise to select appropriate patients for MET-targeted treatment with the array of agents currently in development.

Expression of MET/p-MET and HGF proteins

Human tissue microarray studies reveal that while HGF is widely expressed in both tumor and nonmalignant tissue, MET is differentially expressed in solid tumors (35). Positive staining for MET and HGF, which is thought to have a progenitor role, was observed in the bronchioalveolar junctions of lung tumors (35). Overexpression of MET occurs with a high frequency (35%–72%) in NSCLC tumors (Table 1). For example, in a recent study of more than 200 NSCLC samples, 37% were scored as immunohistochemistry (IHC) $\geq 2+$ for MET expression (36). In another study, MET was detected in eight of nine NSCLC cell lines and in all of 23 NSCLC tumor samples examined (37). Furthermore, 61% of tumor tissues strongly expressed MET, with high MET expression being confirmed as particularly common in adenocarcinoma (67%). It is noteworthy that increased levels of circulating MET mRNA, which were 1.4–8 times above normal concentrations in 68% of patients with overexpression of MET in their tumors, have been found to correlate with early disease recurrence in NSCLC patients (38).

In addition to total levels of the protein, MET activated by ligand to induce phosphorylation of the JM domain can be assayed via phospho-MET (p-MET). Using IHC, specific expression of p-MET has been observed in approximately two-thirds of lung cancer samples and has also been reported to be preferentially expressed at the invasive fronts of NSCLC tumors (35, 37). In a study of the expression and prognostic role of MET, p-MET, and HGF in patients with NSCLC and SCLC ($N = 129$), high expression of two specific forms of p-MET—cytoplasmic expression of Y1003 and nuclear expression of Y1365—was associated with significantly worse overall survival [OS; $P = 0.016$; hazard ratio (HR), 1.86; 95% confidence interval: 1.12–3.07; and $P = 0.034$; HR: 1.70; 95% confidence interval: 1.04–2.78, respectively]. Consequently, specific forms of p-MET may also serve as potential biomarkers in lung cancer (39).

Serum HGF (s-HGF) is also feasible as a biomarker in MET-addicted cancer. Levels of s-HGF were significantly elevated in patients with SCLC compared with healthy individuals (0.40 ± 0.17 vs. 0.26 ± 0.093 , $P = 0.0083$; ref. 40). A high s-HGF level has also been shown to be associated with epithelial-to-mesenchymal transition in patients with SCLC ($N = 112$; ref. 41). Of these patients with stage IV disease, increased s-HGF levels at response evaluation ($P = 0.042$) and at progression ($P = 0.003$) were associated with poor outcome (41).

MET mutations

Mutations in the splice site of *MET* that result in skipping of exon 14 are important molecular drivers in NSCLC (37, 42). Such mutations have recently been shown to occur in 3% to 4% of NSCLC adenocarcinomas, 2% of squamous cell carcinomas, and 1% to 8% of other subtypes of lung cancer (Table 1; refs. 43–46). Novel JM domain (exon 14/15) mutations in *MET* were first identified in SCLC, in three of 10 cell lines and in four of 32 SCLC tumor tissue samples examined (42). *MET* alterations included two different missense mutations in the JM domain (R988C found in NCI-H69 and H249 cell lines; and T1010I in SCLC

Table 1. Reported incidence and functional consequences of MET biomarkers in lung cancer

MET dysregulation	Functional consequences	Biomarker	Reported incidence in samples from lung cancer patients	Reference
Gene overexpression	Reduces or removes the requirement for ligand activation, leading to sustained or altered signaling properties of the receptor	MET/p-MET expression by IHC	NSCLC	
			37% IHC $\geq 2+$	(36)
			61% IHC $\geq 2+$	(37)
			ADC	
			35%	(76)
			67% IHC $\geq 2+$	(37)
			72%	(77)
HGF expression	Ligand-induced activation could cause sustained or altered signaling	Circulating plasma HGF	SCC	
			38%	(77)
Gene mutation	<i>MET</i> mutation can lead to reduced degradation, with consequent overexpression and sustained or altered signaling	<i>MET</i> exon 14 skipping mutation	p-MET in NSCLC	
			67%	(35, 37)
			Elevated in SCLC	(40)
			ADC	
			3%	(43–46)
Gene amplification	Can lead to overexpression and reduce or remove the requirement for ligand activation, leading to sustained or altered signaling properties of the MET receptor	<i>MET</i> GCN MET/CEP7 ratio	4%	(43, 44, 46, 78)
			SCC	
			2%	(45)
			Other lung cancer subtypes	
			2%	(43, 44, 46)
Gene rearrangement	May reduce or remove the requirement for ligand activation, leading to sustained or altered signaling properties of the MET receptor	<i>MET</i> rearrangement	1%–8%	(45)
			Newly diagnosed ADC	
			2%	(46)
			4%	(21, 56)
			5%	(55)
			EGFR TKI-resistant ADC	
			5%	(57, 58)
Downstream MET signaling alteration	Decreases RTK turnover to perpetuate oncogenic activation of MET	<i>CBL</i> mutation or LOH	17%	(33)
			21%	(31)
			22%	(34)
			Identified in an ADC patient	(67)
			Detected in NSCLC patients	(69)

Abbreviations: ADC, adenocarcinoma; CEP, chromosome enumeration probe; SCC, squamous cell carcinoma.

tumor sample). Also, a semaphorin domain missense mutation (E168D in SCLC tumor sample), two-base-pair insertional mutations [IVS13-(52–53)insCT in SCLC tumor samples] within the pre-JM intron 13, as well as an alternative transcript involving exon 10 (H128 cell line), were identified (42). The two reported JM mutations affected cell proliferation, resulting in small but significant growth factor independence in the IL3-dependent BaF3 cell line, and were found to regulate cell morphology and adhesion, and enhanced tumorigenicity when introduced into an SCLC cell line (42). The JM mutations also altered MET RTK signaling, resulting in preferentially increased constitutive tyrosine phosphorylation of various cellular proteins, with significant implications in cytoskeletal functions and metastatic potential. These novel *MET* JM gain-of-function somatic mutations associated with a more aggressive phenotype were among those mutations subsequently identified in NSCLC adenocarcinoma tissues (R988C, R988C + T1010I, S1058P, and an alternative exon 14 splice variant product skipping the entire JM domain; ref. 37).

Using NSCLC tissues and cell lines, we (37) and Kong-Beltran and colleagues (22) functionally characterized tumor-specific somatic intronic *MET* mutations, which immediately flank exon 14 (22). Exon 14 was found to encode the JM domain and Y1003 residue that serves as the binding site for casitas B-lineage lym-

phoma (CBL), the E3 ubiquitin ligase that controls MET turnover (22). In each case of *MET* exon 14 skipping, confirmed by reverse transcriptase polymerase chain reaction, the result was a decrease in the ubiquitination of MET and consequent delayed receptor downregulation after stimulation with HGF. Downstream ligand-dependent signaling through MAPK was also prolonged in cells transfected with a *MET* exon 14 splice variant (22). Furthermore, a xenograft model of Rat1A fibroblasts stably transfected with a *MET* exon 14 splice variant developed particularly aggressive tumors compared with wild-type *MET* (22). Overall, the biological effects of *MET* JM mutations are increased tumorigenicity, reduced adhesion, and disorganized cytoarchitecture compared with wild-type, increased cell survival, motility and migration, increased phosphorylation of focal adhesion proteins, such as paxillin, and HGF independence (23, 37, 47).

Elegant studies have validated the nematode *C. elegans* as a model for rapid screening of the functional aspects of mutant forms of cancer genes, with METR988C and METT1010I harboring wild-type or frequently seen mutant forms of MET in lung cancer (48). Expression of these mutations in this model led to low fecundity and abnormal vulval development characterized by hyperplasia. In addition, exposure of *MET*-mutant transgenic worms to nicotine resulted in enhanced abnormal vulval

Table 2. Case reports and series of patients with lung adenocarcinomas and *MET* exon 14 alterations responding to *MET* inhibitors

Patient no.	Age and gender	Smoking status	<i>MET</i> exon 14 alterations	<i>MET</i> IHC	<i>MET</i> amp	Best response	Reference
1	86 M	NS	Splice acceptor deletion	2+	NA	PR to crizotinib	Jenkins <i>et al.</i> , 2015 (50)
2	71 M	ES	D1028H Splice donor mutation	NA	No	PR to crizotinib	Waqar <i>et al.</i> , 2015 (51)
3	76 F	ES	D1010H	NA	NA	PR to crizotinib	Mendenhall <i>et al.</i> , 2015 (52)
4	80 F	NS	Splice donor mutation	3+	Yes	CR (PERCIST) to cabozantinib	Paik <i>et al.</i> , 2015 (44)
5	78 M	ES	Splice donor deletion	3+	NA	PR to crizotinib (lung) PD to crizotinib (liver)	Paik <i>et al.</i> , 2015 (44)
6	65 M	ES	Splice donor mutation	NA	NA	PR to crizotinib	Paik <i>et al.</i> , 2015 (44)
7	90 F	NS	Splice donor mutation	NA	NA	PR to crizotinib	Paik <i>et al.</i> , 2015 (44)
8	67 F	NS	D1028N Splice donor mutation	NA	NA	PR to crizotinib	Mahjoubi <i>et al.</i> , 2016 (79)

Abbreviations: CR, complete response; ES, ever-smoker; F, female; M, male; NA, not available; NS, never-smoker; PERCIST, PET Response Criteria in Solid Tumors; PR, partial response.

development, fecundity, and locomotion (48). This model also demonstrated colocalization of *MET* and *EGL5* (*PAX8* ortholog) proteins in embryos of the organism (49). *PAX8* provides signals for growth and motility of NSCLC cells and is required for *MET* and *RON* expression; also, it may have therapeutic potential (49).

Responses to the *MET* inhibitors crizotinib and cabozantinib have been documented in case reports of patients with lung adenocarcinoma and *MET* exon 14 alterations (Table 2; refs. 44, 50–52). In phase I clinical studies of the investigational *MET* inhibitor capmatinib (INC280, Novartis), two patients with *MET*-dependent NSCLC and *MET* exon 14 alterations were identified by comprehensive genomic profiling. In one patient with large-cell carcinoma who was treated for over 5 months, there was a partial response comprising a 53% reduction in tumor, and in the other patient, who had squamous NSCLC that had failed prior therapies and was treated for 13 months, there was a partial response comprising a 61% reduction in tumor (43). *MET* mutations in the semaphorin domain have been shown to affect ligand binding: *MET*-N375S, the most frequent mutation of *MET*, most common among male smokers and squamous cell carcinoma, confers resistance to *MET* inhibition based on HGF binding, molecular modeling, and apoptotic susceptibility to *MET* inhibitor studies (53).

Larger clinical studies focusing on patients with *MET* mutations, particularly exon 14 alterations, are now required to prospectively obtain response rates associated with *MET* inhibitors in this patient population. These studies will also need to evaluate any association of *MET* inhibitor efficacy with known disease driving mutations such as *KRAS*, *EGFR*, *BRAF*, or *ALK* (54). Nonetheless, because the potentially actionable genetic alterations within exon 14 are diverse, in-depth molecular profiling of all lung cancer patients, irrespective of additional disease driving mutations, is recommended (54).

MET amplification

In NSCLC, amplification of *MET* typically occurs in about 2% to 5% of newly diagnosed adenocarcinomas (Table 1; refs. 21, 46, 55, 56). Interestingly, a much greater incidence of *MET* amplification, ranging from 5% to 22%, has been reported in patients with NSCLC following treatment with erlotinib/gefitinib (Table 1; refs. 31, 33, 34, 57, 58). Amplification of *MET* (and overexpression of the protein) is also a common event in brain metastases of NSCLC (59). Furthermore, fluorescence *in situ* hybridization (FISH)-positive *MET* status predicts worse survival in patients with advanced NSCLC (56, 60). An analysis of OS based on *MET* FISH status-derived GCN revealed that increased GCN is an independent negative prognostic factor in surgically resected

NSCLC, with OS of 25.8 months for patients with *MET* ≥ 5 copies/cell compared with 47.5 months for patients with *MET* < 5 copies/cell ($P = 0.0045$; ref. 21). These data support further investigation of anti-*MET* therapeutic strategies in appropriately selected patients (21). The question remains as to how biomarkers should be best utilized for patient selection.

While preclinical studies indicated that agents targeting *MET* are effective in the presence of high levels of *MET* gene amplification (61, 62), there is currently no clear consensus on how best to determine *MET* gene amplification in the clinical setting. In a phase I study of capmatinib, preliminary antitumor activity was seen in patients with *EGFR*-wild-type NSCLC and a high level of *MET* amplification (*MET* GCN ≥ 6 ; ref. 63), while a study of capmatinib plus gefitinib in patients with *EGFR*-mutant, *MET*-positive NSCLC reported an overall response rate of 50% in patients with *MET* GCN ≥ 6 (64). Although, based on preliminary data, *MET* GCN appears to be a good predictive biomarker, the FISH *MET*/chromosome enumeration probe 7 (CEP7) ratio is also a relatively simple primary measure of amplification. In a study of crizotinib in *MET*-amplified NSCLC, as determined by *MET*/CEP7 ratio [≥ 1.8 to ≤ 2.2 (low), > 2.2 to < 5 (intermediate) and ≥ 5 (high)], antitumor activity was seen in patients with intermediate and high ratios, with a high response to therapy only observed in individuals in the gene ratio ≥ 5 category (65). One possible drawback of using the *MET*/CEP7 gene ratio is that this technique may not identify all amplified patients due to the unique pathophysiology of NSCLC. In some cases, amplicons occur that include the centromere control protein and the *MET* gene or the centromere protein but not the *MET* gene; in the latter case, the ratio may be falsely lowered (66).











MET rearrangement

Compared with mutations and amplification of *MET*, gene rearrangements are less well documented. However, the kinase fusion *KIF5B-MET* has been identified in a case of lung adenocarcinoma, and it is feasible that this translocation event could potentially account for a significant portion of *MET*-driven oncogenesis (67). This fusion-driven activation of *MET* is most likely due to constitutive dimerization and is likely to be an actionable target for drug-induced inhibition, similar to other fusions in lung cancer such as those involving *ALK*, *ROS1*, and *RET* (67).

MET processing: degradation/transcription


CBL is a mammalian gene encoding an E3 ubiquitin ligase and adaptor protein involved in cell signaling and protein ubiquitination (68). *CBL* thus has an important role in RTK down-regulation and degradation (68). Somatic mutations [or loss of

Table 3. MET-targeted therapies in development for NSCLC and/or solid tumors

Agent and structure	Target(s)	Company	Status (MET ⁺ indications)
Small molecule tyrosine kinase inhibitors			
Crizotinib (PF-02341066) 	ALK/ROS/MET	Pfizer	Phase II (NSCLC, GC, UC, pRCC)
Capmatinib (INC280, INCB28060) 	MET	Novartis	Phase II (NSCLC, HCC, pRCC, CRC, HNSCC)
SARI25844 	MET	Sanofi	Phase II (NSCLC)
Cabozantinib (XL184) 	MET/RET/others	Exelixis	Phase II (NSCLC)
Glesatinib (MGCD265) 	MET/AXL/others	Mirati Therapeutics	Phase II (NSCLC)
Tepotinib (EMD-1214063, MSC2156119J) 	MET	Merck KGaA	Phase II (NSCLC, HCC)
Merestinib (LY2801653) 	MET/ROS1/AXL/FLT3/others	Eli Lilly	Phase II (NSCLC)
S49076 	MET/AXL/FGFR1-3	Servier	Phase I/II (NSCLC)
AMG337 	MET	Amgen	Phase I (GC, ST)
Savolitinib (AZD6094, HMPL-504, volitinib) 	MET	AstraZeneca/Hutchison Medi Pharma	Phase I (pRCC, GC, NSCLC)

(Continued on the following page)

Table 3. MET-targeted therapies in development for NSCLC and/or solid tumors. (Cont'd)

Agent and structure	Target(s)	Company	Status (MET ⁺ indications)
Sitravatinib (MGCD516)	MET/VEGFR/others	Mirati Therapeutics	Phase I (NSCLC, ST)
			
Monoclonal antibodies			
Emibetuzumab (LY2875358)	MET	Eli Lilly	Phase II (NSCLC, GC)
Ficlatuzumab (AV-299)	HGF	AVEO	Phase II (NSCLC); Phase I (HNSCC)

Abbreviations: CRC, colorectal cancer; FGFR, fibroblast growth factor receptor; FLT3, Fms-related tyrosine kinase 3; GC, gastric or esophageal carcinoma; HCC, hepatocellular carcinoma; HNSCC, squamous cell carcinoma of the head and neck; pRCC, papillary renal cell carcinoma; ST, solid tumors; UC, urothelial cancer; VEGFR, vascular endothelial growth factor receptor.

heterozygosity (LOH)] in *CBL* have been detected in NSCLC patients, and expression of these mutations in cell lines was found to result in increased proliferation and cell motility (69). *CBL* LOH is associated with either *MET* or *EGFR* mutations and may contribute to their oncogenic potential (69). As already described, it is noteworthy that the JM domain of MET is involved in the binding and E3 activity of CBL, and *MET* JM mutations (e.g., Y1003) therefore affect CBL binding and decrease RTK turnover to perpetuate oncogenic activation of MET (22, 70).






Discussion: Potential of MET as a Biomarker in Lung Cancer

The growing prominence of MET inhibition in lung cancer is reflected in the number of molecular aberrations with oncogenic potential that occur in this disease, and in the number and diversity of MET-targeted agents in clinical development in this indication. These include the monoclonal antibodies emibetuzumab, ficlatuzumab, and rilotuzumab, and tyrosine kinase inhibitors (TKI) such as crizotinib, tepotinib, cabozantinib, and capmatinib (Table 3). Recent negative or disappointing clinical trials results pose the question as to whether the biomarkers and their related cutoff values have been chosen appropriately to select patients for enrollment in all studies to date. For instance, despite positive phase II data (71), the phase III METLung trial ($N = 499$) of onartuzumab plus erlotinib failed to show clinical benefit compared with placebo plus erlotinib in patients with MET⁺ NSCLC (Table 4; ref. 72). In this study, patient biomarker-based selection of patients with MET-overexpressing tumors as measured by IHC (MET SP44) was used to determine eligibility. These negative data suggest that IHC may not be sufficiently sensitive as a diagnostic tool for MET positivity; its use as a standard biomarker for overexpression is further compromised by the lack of standardized interpretation or consensus on optimized cutoff values. Moreover, MET protein expression may have a low predictive value as a tool to detect MET activation and may not always reflect tumor dependency on MET signaling (73). Heterogeneity in the expression of MET within a tumor or across metastatic sites may also lead to unreliable results.

Circulating HGF (cHGF) or MET are attractive potential alternative biomarkers for ligand or receptor overexpression, respectively. For example, elevated cHGF, as measured by ELISA, has been used as a pharmacodynamic biomarker of activity with onartuzumab (74). However, in cases of ligand-independent activation of MET, it is feasible that monoclonal antibody therapy, without drug internalization, may be a less effective therapeutic


strategy than TKIs that target the receptor protein kinase directly. Recent data suggest that HGF/MET protein expression alone may be an oversimplification of the oncogenic driver status of "MET-positive" NSCLC, where mutations or translocations and amplification reduce the requirement for ligand activation and lead to sustained or altered signaling properties of the receptor. Although IHC data have been shown to correlate with MET amplification (66), clinical study biomarker data (summarized in Table 4) have not confirmed any clear-cut relationships between MET mutation, amplification, and overexpression, when collectively applied as predictive biomarkers for MET-targeted therapy. IHC-based MET expression has not been a successful biomarker approach in clinical studies of monoclonal antibodies, and current clinical and biomarker data suggest that genetic changes in *MET*, in particular gene amplification, may be the preferred biomarkers for MET TKI therapy (21, 63–65). The data summarized in Table 4 also indicate that biomarkers for MET TKI therapy need to be optimized based on not only *MET* amplification but also *MET* mutation or translocation status, which constitutes an additional and numerically significant (>4%) molecular subgroup of NSCLC (46). Mutations or altered expression of signaling proteins downstream of MET signaling, such as *CBL* mutation, are also emerging biomarkers in NSCLC and extend further the range and diversity of potential MET-related biomarkers in this disease. There is therefore an urgent need to elucidate both optimal biomarkers for MET dysregulation, and their application, based on our growing understanding of this oncogenic driver in NSCLC. To facilitate this goal, the medical oncologist and pathologist now have at their disposal a panel of genetic and protein biomarkers for MET dysregulation that together constitute a significant proportion of lung cancer molecular subgroups. Indeed, current data indicate that panels of MET biomarkers are likely to be necessary in the future, and measurements of potential biomarkers should therefore be included in new clinical trial designs for MET inhibitors to facilitate the robust definition of appropriate therapies for specific MET-dysregulated NSCLC subsets. Furthermore, the recent report of a response to crizotinib in a patient with lung adenocarcinoma with MET copy-number gain but without a high-level MET/CEP7 ratio, MET overexpression, or exon 14 splicing mutation (75) indicates that the list of independent predictive biomarkers for response to MET inhibitors may well be extended further. Importantly, since alterations in *MET* gene status have been found to occur in both untreated patients and those who have developed resistance to other targeted therapies, new clinical study designs should consider both patient groups. This

Table 4. Predictive biomarkers evaluated in clinical studies of MET-targeted therapies in patients with NSCLC

Agent	Study	Biomarker (assay)	Comment	Reference
Tyrosine kinase inhibitors				
 Cabozantinib (XL184)	Phase II cabozantinib + erlotinib in pretreated EGFR-mut NSCLC (N = 37)	MET amplification/GCN (FISH)	Clinical activity (ORR/DCR 8%/65%) was unrelated to MET amplification (not detected in any patients)	Reckamp et al., 2015 (80)
 Capmatinib (INC280, INCB28060)	Phase I MET+ (H-score ≥ 150 or MET/centromere ratio ≥ 2.0 or MET GCN ≥ 5 , or $\geq 50\%$ of tumor cells IHC 2/3+ or IHC 3+ [expansion]) NSCLC (N = 55)	MET expression (IHC) MET amplification/GCN (FISH) MET Δ ex14 mutation (NGS)	10/18 MET IHC 3+ patients showing tumor shrinkage also had MET GCN ≥ 6 ORR/DCR 20%/51% in all patients; ORR/DCR 47%/80% in patients with MET GCN ≥ 6 and 24%/60% in patients with IHC 3+ Preliminary median PFS 3.6 months, increased to 7.4 months for patients with MET GCN ≥ 6 Preliminary activity in patients with either a high level of MET amplification and/or cMET Δ ex14 mutation	Schuler et al., 2016 (63)
 Crizotinib (PF-02341066)	Phase II capmatinib + gefitinib in MET+ (MET GCN ≥ 4 , or $\geq 50\%$ of tumor cells IHC 3+)/EGFR-mut NSCLC (N = 14)	MET expression (IHC) MET amplification/GCN (FISH)	ORR/DCR 3%/81% GCN <4: ORR/DCR 14%/77% 4 \leq GCN <6: ORR/DCR 24%/88% GCN ≥ 6 : ORR/DCR 50%/84%	Wu et al., 2016 (64)
 Glesatinib (MGCD265)	Phase I MET+ (MET-amplified) NSCLC (PROFILE 1001; N = 21) Phase I advanced solid tumors (N = 25) including NSCLC (n = 11)	MET amplification as MET/CEP7 ratio (FISH) MET mutation or amplification	Tumor shrinkage seen in intermediate MET (MET/CEP7 ratio > 2.2 to < 5 ; ORR 17%) and high MET (MET/CEP7 ratio ≥ 5 ; ORR 67%) cohorts Clinically meaningful antitumor activity in this subgroup ORR/DCR 4.4%/94% The first three NSCLC patients (two with MET Δ ex14 and one with MET GCN gain) demonstrated tumor regression	Camidge et al., 2015 (65) Drilon et al., 2016 (78) Kollmannsberger et al., 2015 (81)
 Tepotinib (EMD-1214063, MSC2156119J)	Phase I advanced solid tumors/NSCLC (N = 143)	MET expression (IHC) MET amplification/GCN (FISH)	Exploratory biomarker analyses suggested MET expression and amplification were associated with response	Falchook et al., 2015 (82)

(Continued on the following page)

Table 4. Predictive biomarkers evaluated in clinical studies of MET-targeted therapies in patients with NSCLC. (Cont'd)

Agent	Study	Biomarker (assay)	Comment	Reference
 Tivantinib (ARQ197)	Phase II tivantinib + erlotinib (T + E) in advanced NSCLC (N = 167)	MET amplification/GCN (FISH)	Patients with increased MET GCN demonstrated a trend towards PFS benefit from T + E MET GCN 2, 3, 4, and 5 revealed PFS HRs of 0.92, 0.75, 0.71, and 0.42, respectively Similar but less vigorous trend for OS No evidence that T + E was worse than E + placebo in patients with low MET GCN (<2 or <3)	Sequist <i>et al.</i> , 2011 (83)
	Phase III erlotinib ± tivantinib in advanced NSCLC (N = 1,048)	MET expression (IHC) MET amplification/GCN (FISH)	Exploratory subgroup analysis suggested OS improvement in patients with high MET expression (HR 0.70) Longer OS in patients with tumors with MET GCN >4 (HR 0.83), but limited sample size Study discontinued for futility	Scagliotti <i>et al.</i> , 2015 (84)
Erimbetuzumab (LY2875358)	Phase III (Asia) erlotinib ± tivantinib in advanced NSCLC (ATTENTION; N = 307)	MET expression (IHC) MET amplification/GCN (FISH) HGF expression (IHC) Serum HGF (ELISA)	Tivantinib was associated with a weak OS benefit in patient with high MET expression (HR 0.83) Significant OS benefit in patients with normal MET GCN (<4, HR 0.51) Significant OS benefit in patients with high HGF expression (H-score ≥200; HR 0.54) Favorable effect on OS in patients with high serum HGF concentration Enrollment stopped for safety reasons (LD incidence)	Yoshioka <i>et al.</i> , 2015 (85)
	Phase I (MET ⁺ (IHC ≥2+) solid tumors (N = 62), including NSCLC (n = 19)	MET expression (IHC)	Preliminary, but limited, single-agent clinical activity was observed (DCR 26% in patients with MET IHC ≥2+ NSCLC) IHC was not considered to be a sufficient biomarker	Bankk <i>et al.</i> , 2015 (86)
Ficlatuzumab	Phase II (Asia) gefitinib ± ficlatuzumab (G + F) in lung adenocarcinoma (N = 188)	MET expression (IHC) HGF expression (IHC)	No statistically significant improvement in ORR/PFS Notable difference seen in low MET group (ORR 41% vs. 22%) and median PFS (7.3 vs. 2.8 months) favoring G + F Preliminary OS favored G + F in patients with high stromal HGF (P = 0.03), and high MET (P = 0.18) biomarkers	Mok <i>et al.</i> , 2012 (87)
Onartuzumab	Phase I (Japan) solid tumors and MET ⁺ lung cancer	MET expression (IHC)	PR observed in a patient with an IHC 3+ tumor	Nishio <i>et al.</i> , 2015 (88)
	Phase II onartuzumab + erlotinib in advanced NSCLC (N = 137)	MET expression (IHC) MET expression (IHC) MET amplification/GCN (FISH) MET mRNA expression (RT-PCR) HGF mRNA (RT-PCR) Circulating plasma HGF (ELISA) MET exon 14 (RT-PCR)	Benefit maintained in patients with IHC 2/3+ Detriment in patients with IHC 0/1+ Benefit was proportional to the intensity of expression MET IHC was the most robust predictor of OS and PFS benefit Non-significant OS improvement in patients with high MET GCN (mean ≥5 copies/cell) Benefit maintained in MET IHC ⁺ /MET FISH ⁻ patients (HR 0.37; P = 0.01) MET and HGF mRNA levels did not predict significant benefit; nonsignificant OS improvement in patients with high tumor MET mRNA levels (HR 0.59) OS favored onartuzumab in patients with low baseline plasma HGF (HR 0.52; P = 0.09)	Spigel <i>et al.</i> , 2013 (71) Koeppen <i>et al.</i> , 2014 (89)
	Phase III onartuzumab + erlotinib in advanced MET ⁺ (IHC 2/3+) NSCLC (N = 499)	MET expression (IHC)	No improvement in OS, PFS, or ORR Trial stopped for futility	Spigel <i>et al.</i> , 2014 (72)

Abbreviations: DCR, disease control rate; ELISA, enzyme-linked immunosorbent assay; IHC, interstitial lung disease; METΔex14, MET exon 14 deletion; NGS, next-generation sequencing; ORR, overall response rate; PFS, progression-free survival; PR, partial response; RT-PCR, reverse transcription polymerase chain reaction.

highlights the future importance of both upfront and resistance-based genetic testing in lung cancer patients, which should include MET as the probable next major biomarker in lung cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

This manuscript was written by the author with medical editorial assistance provided by Matthew Naylor PhD, funded by Novartis Pharmaceuticals Corporation.

Grant Support

Medical editorial assistance was funded by Novartis Pharmaceuticals Corporation.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received July 20, 2016; revised November 21, 2016; accepted December 19, 2016; published online April 3, 2017.

References

- Cooper CS, Park M, Blair DC, Tainsky MA, Huebner K, Croce CM, et al. Molecular cloning of a new transforming gene from a chemically transformed human cell line. *Nature* 1984;311:29–33.
- Giordano S, Ponzetto C, Di Renzo MF, Cooper CS, Comoglio PM. Tyrosine kinase receptor indistinguishable from the c-met protein. *Nature* 1989; 339:155–6.
- Naldini L, Vigna E, Narsimhan RP, Gaudino G, Zarnegar R, Michalopoulos GK, et al. Hepatocyte growth factor (HGF) stimulates the tyrosine kinase activity of the receptor encoded by the proto-oncogene c-MET. *Oncogene* 1991;6:501–4.
- Fixman ED, Fournier TM, Kamikura DM, Naujokas MA, Park M. Pathways downstream of Shc and Grb2 are required for cell transformation by the tpr-Met oncoprotein. *J Biol Chem* 1996;271:13116–22.
- Bottaro DP, Rubin JS, Faletto DL, Chan AM, Kmiecik TE, Vande Woude GF, et al. Identification of the hepatocyte growth factor receptor as the c-met proto-oncogene product. *Science* 1991;251:802–4.
- Ponzetto C, Bardelli A, Zhen Z, Maina F, dalla Zonca P, Giordano S, et al. A multifunctional docking site mediates signaling and transformation by the hepatocyte growth factor/scatter factor receptor family. *Cell* 1994;77:261–71.
- Weidner KM, Di Cesare S, Sachs M, Brinkmann V, Behrens J, Birchmeier W. Interaction between Gab1 and the c-Met receptor tyrosine kinase is responsible for epithelial morphogenesis. *Nature* 1996;384:173–6.
- Sipeki S, Bander E, Buday L, Farkas G, Bacsy E, Ways DK, et al. Phosphatidylinositol 3-kinase contributes to Erk1/Erk2 MAP kinase activation associated with hepatocyte growth factor-induced cell scattering. *Cell Signal* 1999;11:885–90.
- Zhang YW, Wang LM, Jove R, Vande Woude GF. Requirement of Stat3 signaling for HGF/SF-Met mediated tumorigenesis. *Oncogene* 2002;21: 217–26.
- Van Der Steen N, Pauwels P, Gil-Bazo I, Castanon E, Ruez L, Cappuzzo F, et al. cMET in NSCLC: Can we cut off the head of the hydra? From the pathway to the resistance. *Cancers (Basel)* 2015;7:556–73.
- Schmidt C, Bladt F, Goedecke S, Brinkmann V, Zschiesche W, Sharpe M, et al. Scatter factor/hepatocyte growth factor is essential for liver development. *Nature* 1995;373:699–702.
- Uehara Y, Minowa O, Mori C, Shiota K, Kuno J, Noda T, et al. Placental defect and embryonic lethality in mice lacking hepatocyte growth factor/scatter factor. *Nature* 1995;373:702–5.
- Maina F, Hilton MC, Ponzetto C, Davies AM, Klein R. Met receptor signaling is required for sensory nerve development and HGF promotes axonal growth and survival of sensory neurons. *Genes Dev* 1997;11: 3341–50.
- Chmielowiec J, Borowiak M, Morkel M, Stradal T, Munz B, Werner S, et al. c-Met is essential for wound healing in the skin. *J Cell Biol* 2007;177:151–62.
- Huh CG, Factor VM, Sanchez A, Uchida K, Conner EA, Thorgeirsson SS. Hepatocyte growth factor/c-met signaling pathway is required for efficient liver regeneration and repair. *Proc Natl Acad Sci USA* 2004;101:4477–82.
- Liu X, Newton RC, Scherle PA. Developing c-MET pathway inhibitors for cancer therapy: progress and challenges. *Trends Mol Med* 2010;16:37–45.
- Sierra JR, Tsao MS. c-MET as a potential therapeutic target and biomarker in cancer. *Ther Adv Med Oncol* 2011;3:S21–35.
- Smyth EC, Sclafani F, Cunningham D. Emerging molecular targets in oncology: clinical potential of MET/hepatocyte growth-factor inhibitors. *Onco Targets Ther* 2014;7:1001–14.
- Park S, Choi YL, Sung CO, An J, Seo J, Ahn MJ, et al. High MET copy number and MET overexpression: poor outcome in non-small cell lung cancer patients. *Histol Histopathol* 2012;27:197–207.
- Tsuta K, Kozu Y, Mimae T, Yoshida A, Kohno T, Sekine I, et al. c-MET/phospho-MET protein expression and MET gene copy number in non-small cell lung carcinomas. *J Thorac Oncol* 2012;7:331–9.
- Cappuzzo F, Marchetti A, Skokan M, Rossi E, Gajopathy S, Felicioni L, et al. Increased MET gene copy number negatively affects survival of surgically resected non-small-cell lung cancer patients. *J Clin Oncol* 2009;27:1667–74.
- Kong-Beltran M, Seshagiri S, Zha J, Zhu W, Bhawe K, Mendoza N, et al. Somatic mutations lead to an oncogenic deletion of met in lung cancer. *Cancer Res* 2006;66:283–9.
- Ma PC, Tretiakova MS, Nallasura V, Jagadeeswaran R, Husain AN, Salgia R. Downstream signalling and specific inhibition of c-MET/HGF pathway in small cell lung cancer: implications for tumour invasion. *Br J Cancer* 2007;97:368–77.
- Puri N, Salgia R. Synergism of EGFR and c-Met pathways, cross-talk and inhibition, in non-small cell lung cancer. *J Carcinog* 2008;7:9.
- Seiwert TY, Jagadeeswaran R, Faoro L, Janamanchi V, Nallasura V, El Dinali M, et al. The MET receptor tyrosine kinase is a potential novel therapeutic target for head and neck squamous cell carcinoma. *Cancer Res* 2009; 69:3021–31.
- Ma PC, Schaefer E, Christensen JG, Salgia R. A selective small molecule c-MET inhibitor, PHA665752, cooperates with rapamycin. *Clin Cancer Res* 2005;11:2312–9.
- Baschnagel AM, Galoforo S, Thibodeau BJ, Ahmed S, Nirmal S, Akervall J, et al. Crizotinib fails to enhance the effect of radiation in head and neck squamous cell carcinoma xenografts. *Anticancer Res* 2015;35:5973–82.
- Rolle CE, Kanteti R, Surati M, Nandi S, Dhanasingh I, Yala S, et al. Combined MET inhibition and topoisomerase I inhibition block cell growth of small cell lung cancer. *Mol Cancer Ther* 2014;13:576–84.
- Wilson TR, Fridlyand J, Yan Y, Penuel E, Burton L, Chan E, et al. Widespread potential for growth-factor-driven resistance to anticancer kinase inhibitors. *Nature* 2012;487:505–9.
- Straussman R, Morikawa T, Shee K, Barzily-Rokni M, Qian ZR, Du J, et al. Tumour micro-environment elicits innate resistance to RAF inhibitors through HGF secretion. *Nature* 2012;487:500–4.
- Bean J, Brennan C, Shih JY, Riely G, Viale A, Wang L, et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc Natl Acad Sci USA* 2007;104:20932–7.
- Sadiq AA, Salgia R. MET as a possible target for non-small-cell lung cancer. *J Clin Oncol* 2013;31:1089–96.
- Chen HJ, Mok TS, Chen ZH, Guo AL, Zhang XC, Su J, et al. Clinicopathologic and molecular features of epidermal growth factor receptor T790M mutation and c-MET amplification in tyrosine kinase inhibitor-resistant Chinese non-small cell lung cancer. *Pathol Oncol Res* 2009; 15:651–8.
- Engelman JA, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park JO, et al. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science* 2007;316:1039–43.
- Ma PC, Tretiakova MS, MacKinnon AC, Ramnath N, Johnson C, Dietrich S, et al. Expression and mutational analysis of MET in human solid cancers. *Genes Chromosomes Cancer* 2008;47:1025–37.

36. Watermann I, Schmitt B, Stellmacher F, Muller J, Gaber R, Kugler C, et al. Improved diagnostics targeting c-MET in non-small cell lung cancer: expression, amplification and activation? *Diagn Pathol* 2015;10:130.
37. Ma PC, Jagadeeswaran R, Jagadeesh S, Tretiakova MS, Nallasura V, Fox EA, et al. Functional expression and mutations of c-Met and its therapeutic inhibition with SU11274 and small interfering RNA in non-small cell lung cancer. *Cancer Res* 2005;65:1479–88.
38. Cheng TL, Chang MY, Huang SY, Sheu CC, Kao EL, Cheng YJ, Chong IW. Overexpression of circulating c-met messenger RNA is significantly correlated with nodal stage and early recurrence in non-small cell lung cancer. *Chest* 2005;128:1453–60.
39. Tretiakova M, Salama AK, Karrison T, Ferguson MK, Husain AN, Vokes EE, et al. MET and phosphorylated MET as potential biomarkers in lung cancer. *J Environ Pathol Toxicol Oncol* 2011;30:341–54.
40. Takigawa N, Segawa Y, Maeda Y, Takata I, Fujimoto N. Serum hepatocyte growth factor/scatter factor levels in small cell lung cancer patients. *Lung Cancer* 1997;17:211–8.
41. Canadas I, Taus A, Gonzalez I, Villanueva X, Gimeno J, Pijuan L, et al. High circulating hepatocyte growth factor levels associate with epithelial to mesenchymal transition and poor outcome in small cell lung cancer patients. *Oncotarget* 2014;5:5246–56.
42. Ma PC, Kijima T, Maulik G, Fox EA, Sattler M, Griffin JD, et al. c-MET mutational analysis in small cell lung cancer: novel juxtamembrane domain mutations regulating cytoskeletal functions. *Cancer Res* 2003;63:6272–81.
43. Frampton GM, Ali SM, Rosenzweig M, Chmielecki J, Lu X, Bauer TM, et al. Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov* 2015;5:850–9.
44. Paik PK, Drilon A, Fan PD, Yu H, Rekhman N, Ginsberg MS, et al. Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov* 2015;5:842–9.
45. Schrock AB, Frampton GM, Suh J, Chalmers ZR, Rosenzweig M, Erlich RL, et al. Characterization of 298 lung cancer patients harboring MET exon 14 skipping (METex14) alterations. *J Thorac Oncol* 2016;11:1493–502.
46. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 2014;511:543–50.
47. Maulik G, Shrikhande A, Kijima T, Ma PC, Morrison PT, Salgia R. Role of the hepatocyte growth factor receptor, c-Met, in oncogenesis and potential for therapeutic inhibition. *Cytokine Growth Factor Rev* 2002;13:41–59.
48. Siddiqui SS, Loganathan S, Krishnaswamy S, Faoro L, Jagadeeswaran R, Salgia R. *C. elegans* as a model organism for in vivo screening in cancer: effects of human c-Met in lung cancer affect *C. elegans* vulva phenotypes. *Cancer Biol Ther* 2008;7:856–63.
49. Kanteti R, El-Hashani E, Dhanasingh I, Tretiakova M, Husain AN, Sharma S, et al. Role of PAX8 in the regulation of MET and RON receptor tyrosine kinases in non-small cell lung cancer. *BMC Cancer* 2014;14:185.
50. Jenkins RW, Oxnard GR, Elkin S, Sullivan EK, Carter JL, Barbie DA. Response to crizotinib in a patient with lung adenocarcinoma harboring a MET splice site mutation. *Clin Lung Cancer* 2015;16:e101–4.
51. Waqar SN, Morgensztern D, Sehn J. MET mutation associated with responsiveness to crizotinib. *J Thorac Oncol* 2015;10:e29–31.
52. Mendenhall MA, Goldman JW. MET-mutated NSCLC with major response to crizotinib. *J Thorac Oncol* 2015;10:e33–4.
53. Krishnaswamy S, Kanteti R, Duke-Cohan JS, Loganathan S, Liu W, Ma PC, et al. Ethnic differences and functional analysis of MET mutations in lung cancer. *Clin Cancer Res* 2009;15:5714–23.
54. Sattler M, Salgia R. MET in the driver's seat: exon 14 skipping mutations as actionable targets in lung cancer. *J Thorac Oncol* 2016;11:1381–3.
55. Kawakami H, Okamoto I, Okamoto W, Tanizaki J, Nakagawa K, Nishio K. Targeting MET amplification as a new oncogenic driver. *Cancers (Basel)* 2014;6:1540–52.
56. Go H, Jeon YK, Park HJ, Sung SW, Seo JW, Chung DH. High MET gene copy number leads to shorter survival in patients with non-small cell lung cancer. *J Thorac Oncol* 2010;5:305–13.
57. Yu HA, Arcila ME, Rekhman N, Sima CS, Zakowski MF, Pao W, et al. Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin Cancer Res* 2013;19:2240–7.
58. Sequist LV, Waltman BA, Dias-Santagata D, Digumarthy S, Turke AB, Fidias P, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med* 2011;3:75ra26.
59. Preusser M, Streubel B, Berghoff AS, Hainfellner JA, von Deimling A, Widhalm G, et al. Amplification and overexpression of CMET is a common event in brain metastases of non-small cell lung cancer. *Histopathology* 2014;65:684–92.
60. Dimou A, Non L, Chae YK, Tester WJ, Syrigos KN. MET gene copy number predicts worse overall survival in patients with non-small cell lung cancer (NSCLC): a systematic review and meta-analysis. *PLoS One* 2014;9:e107677.
61. Smolen GA, Sordella R, Muir B, Mohapatra G, Barmettler A, Archibald H, et al. Amplification of MET may identify a subset of cancers with extreme sensitivity to the selective tyrosine kinase inhibitor PHA-665752. *Proc Natl Acad Sci USA* 2006;103:2316–21.
62. Tanizaki J, Okamoto I, Okamoto K, Takezawa K, Kuwata K, Yamaguchi H, et al. MET tyrosine kinase inhibitor crizotinib (PF-02341066) shows differential antitumor effects in non-small cell lung cancer according to MET alterations. *J Thorac Oncol* 2011;6:1624–31.
63. Schuler MH, Berardi R, Lim W, Geel RV, De Jonge MJ, Bauer TM, et al. Phase (Ph) I study of the safety and efficacy of the cMET inhibitor capmatinib (INC280) in patients (pts) with advanced cMET+ non-small cell lung cancer (NSCLC). *J Clin Oncol* 2016;34:9067.
64. Wu Y, Kim D, Felip E, Zhang L, Liu X, Zhou CC, et al. Phase (Ph) II safety and efficacy results of a single-arm ph Ib/II study of capmatinib (INC280) + gefitinib in patients (pts) with EGFR-mutated (mut), cMET-positive (cMET+) non-small cell lung cancer (NSCLC). *J Clin Oncol* 2016;34:9020.
65. Camidge DR, Ou SI, Shapiro G, Otterson GA, Villaruz LC, Villalona-Calero MA, et al. Efficacy and safety of crizotinib in patients with advanced c-MET-amplified non-small cell lung cancer (NSCLC). *J Clin Oncol* 2014;32:8001.
66. Schildhaus HU, Schultheis AM, Ruschoff J, Binot E, Merkelbach-Bruse S, Fassunke J, et al. MET amplification status in therapy-naïve adeno- and squamous cell carcinomas of the lung. *Clin Cancer Res* 2015;21:907–15.
67. Stransky N, Cerami E, Schalm S, Kim JL, Lengauer C. The landscape of kinase fusions in cancer. *Nat Commun* 2014;5:4846.
68. Swaminathan G, Tsygankov AY. The Cbl family proteins: ring leaders in regulation of cell signaling. *J Cell Physiol* 2006;209:21–43.
69. Tan YH, Krishnaswamy S, Nandi S, Kanteti R, Vora S, Onel K, et al. CBL is frequently altered in lung cancers: its relationship to mutations in MET and EGFR tyrosine kinases. *PLoS One* 2010;5:e8972.
70. Onozato R, Kosaka T, Kuwano H, Sekido Y, Yatabe Y, Mitsudomi T. Activation of MET by gene amplification or by splice mutations deleting the juxtamembrane domain in primary resected lung cancers. *J Thorac Oncol* 2009;4:5–11.
71. Spigel DR, Ervin TJ, Ramlau RA, Daniel DB, Goldschmidt JH Jr, Blumenschein GR Jr, et al. Randomized phase II trial of onartuzumab in combination with erlotinib in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2013;31:4105–14.
72. Spigel DR, Edelman MJ, O'Byrne K, Paz-Ares L, Shames DS, Yu W, et al. Onartuzumab plus erlotinib versus erlotinib in previously treated stage IIIB or IV NSCLC: results from the pivotal phase III randomized, multicenter, placebo-controlled METLung (OAM4971g) global trial. *J Clin Oncol* 2014;32:8000.
73. Finocchiaro G, Toschi L, Gianoncelli L, Baretta M, Santoro A. Prognostic and predictive value of MET deregulation in non-small cell lung cancer. *Ann Transl Med* 2015;3:83.
74. Penuel E, Li C, Parab V, Burton L, Cowan KJ, Merchant M, et al. HGF as a circulating biomarker of onartuzumab treatment in patients with advanced solid tumors. *Mol Cancer Ther* 2013;12:1122–30.
75. Zhang Y, Wang W, Wang Y, Xu Y, Tian Y, Huang M, et al. Response to crizotinib observed in lung adenocarcinoma with MET copy number gain but without a high-level MET/CEP7 ratio, MET overexpression, or exon 14 splicing mutations. *J Thorac Oncol* 2016;11:e59–62.
76. Tsao MS, Liu N, Chen JR, Pappas J, Ho J, To C, et al. Differential expression of Met/hepatocyte growth factor receptor in subtypes of non-small cell lung cancers. *Lung Cancer* 1998;20:1–16.
77. Ichimura E, Maeshima A, Nakajima T, Nakamura T. Expression of c-met/HGF receptor in human non-small cell lung carcinomas in vitro and in vivo and its prognostic significance. *Jpn J Cancer Res* 1996;87:1063–9.

78. Drilon AE, Camidge DR, Ou SI, Clark JW, Socinski MA, Weiss J, et al. Efficacy and safety of crizotinib in patients (pts) with advanced MET exon 14-altered non-small cell lung cancer (NSCLC). *J Clin Oncol* 2016;34:108.
79. Mahjoubi L, Gazzah A, Besse B, Lacroix L, Soria JC. A never-smoker lung adenocarcinoma patient with a MET exon 14 mutation (D1028N) and a rapid partial response after crizotinib. *Invest New Drugs* 2016;34:397–8.
80. Reckamp KL, Mack PC, Ruel N, Frankel PH, Gitlitz BJ, Li T, et al. Biomarker analysis of a phase II trial of cabozantinib and erlotinib in patients (pts) with EGFR-mutant NSCLC with epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) resistance: a California Cancer Consortium Phase II trial (NCI 9303). *J Clin Oncol* 2015;33:8087.
81. Kollmannsberger CK, Sharma S, Shapiro G, Chi KN, Christensen J, Tassell VR, et al. Phase I study of receptor tyrosine kinase (RTK) inhibitor, MGCD265, in patients (pts) with advanced solid tumors. *J Clin Oncol* 2015;33:2589.
82. Falchook GS, Kurzrock R, Amin HM, Fu S, Piha-Paul SA, Janku F, et al. Efficacy, safety, biomarkers, and phase II dose modeling in a phase I trial of the oral selective c-Met inhibitor tepotinib (MSC2156119J). *J Clin Oncol* 2015;33:2591.
83. Sequist LV, von Pawel J, Garmey EG, Akerley WL, Brugger W, Ferrari D, et al. Randomized phase II study of erlotinib plus tivantinib versus erlotinib plus placebo in previously treated non-small-cell lung cancer. *J Clin Oncol* 2011;29:3307–15.
84. Scagliotti G, von Pawel J, Novello S, Ramlau R, Favaretto A, Barlesi F, et al. Phase III multinational, randomized, double-blind, placebo-controlled study of tivantinib (ARQ 197) plus erlotinib versus erlotinib alone in previously treated patients with locally advanced or metastatic nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2015;33:2667–74.
85. Yoshioka H, Azuma K, Yamamoto N, Takahashi T, Nishio M, Katakami N, et al. A randomized, double-blind, placebo-controlled, phase III trial of erlotinib with or without a c-Met inhibitor tivantinib (ARQ 197) in Asian patients with previously treated stage IIIB/IV nonsquamous non-small-cell lung cancer harboring wild-type epidermal growth factor receptor (ATTENTION study). *Ann Oncol* 2015;26:2066–72.
86. Banck MS, Chugh R, Natale RB, Algazi A, Carthon BC, Rosen LS, et al. Phase 1 results of emibetuzumab (LY2875358), a bivalent MET antibody, in patients with advanced castration-resistant prostate cancer, and MET positive renal cell carcinoma, non-small cell lung cancer, and hepatocellular carcinoma. *Mol Cancer Ther* 2015;14:A55.
87. Mok TSK, Park K, Geater SL, Agarwal S, Han M, Credi M, et al. A randomized phase (Ph) 2 study with exploratory biomarker analysis of ficlatuzumab (F) a humanized hepatocyte growth factor (HGF) inhibitory mAb in combination with gefitinib (G) versus G in Asian patients (pts) with lung adenocarcinoma (LA). *Ann Oncol* 2012;23:1198P.
88. Nishio M, Horiike A, Nokihara H, Horinouchi H, Nakamichi S, Wakui H, et al. Phase I study of the anti-MET antibody onartuzumab in patients with solid tumors and MET-positive lung cancer. *Invest New Drugs* 2015; 33:632–40.
89. Koeppen H, Yu W, Zha J, Pandita A, Penuel E, Rangell L, et al. Biomarker analyses from a placebo-controlled phase II study evaluating erlotinib ± onartuzumab in advanced non-small cell lung cancer: MET expression levels are predictive of patient benefit. *Clin Cancer Res* 2014;20:4488–98.

Molecular Cancer Therapeutics

MET in Lung Cancer: Biomarker Selection Based on Scientific Rationale

Ravi Salgia

Mol Cancer Ther 2017;16:555-565.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/16/4/555>

Cited articles This article cites 89 articles, 28 of which you can access for free at:
<http://mct.aacrjournals.org/content/16/4/555.full#ref-list-1>

Citing articles This article has been cited by 1 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/16/4/555.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.