

1,1-Bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes inhibit colon cancer cell and tumor growth through PPAR γ -dependent and PPAR γ -independent pathways

Sudhakar Chintharlapalli,¹ Sabitha Papineni,² and Stephen Safe^{1,2,3}

Departments of ¹Biochemistry and Biophysics and ²Physiology and Pharmacology, Texas A&M University, College Station, Texas and ³Institute of Biosciences and Technology, Texas A&M University Health Science Center, Houston, Texas

Abstract

1,1-Bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes containing *p*-trifluoromethyl, *t*-butyl, and phenyl [1,1-bis(3'-indolyl)-1-(*p*-phenyl)methane (DIM-C-pPhC₆H₅)] substituents induce peroxisome proliferator-activated receptor γ (PPAR γ)-mediated transactivation in SW480 colon cancer cells. These PPAR γ -active compounds also inhibit cell proliferation and modulate some cell cycle proteins. At concentrations from 2.5 to 7.5 μ mol/L, the PPAR γ agonists induce caveolin-1 and phosphorylation of Akt and cotreatment with the PPAR γ antagonist GW9662 inhibited the induction response. In contrast, higher concentrations (10 μ mol/L) of 1,1-bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes containing 1,1-bis(3'-indolyl)-1-(*p*-trifluoromethyl)methane and DIM-C-pPhC₆H₅ induce apoptosis, which is PPAR γ independent. This was accompanied by loss of caveolin-1 induction but induction of proapoptotic nonsteroidal anti-inflammatory drug activated gene-1. In athymic nude mice bearing SW480 cell xenografts, DIM-C-pPhC₆H₅ inhibits tumor growth at doses of 20 and 40 mg/kg/d and immunohistochemical staining of the tumors showed induction of apoptosis and nonsteroidal anti-inflammatory drug activated gene-1 expression. Thus, the indole-derived PPAR γ -active compounds induce both receptor-dependent and receptor-independent responses in SW480 cells, which are separable over a narrow range of concentrations. This dual mechanism of action enhances their antiproliferative and anticancer activities. [Mol Cancer Ther 2006;5(5):1362–70]

Introduction

Peroxisome proliferator-activated receptor γ (PPAR γ) is a member of the nuclear receptor superfamily of ligand-activated transcription factors (1–6). Although the endogenous ligand for PPAR γ has not been determined, several endogenous molecules, such as 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂, other prostaglandin-derived compounds, fatty acids, and flavonoids, activate PPAR γ . It has recently been reported that nitrolinoleic acid, a stress-induced fatty acid oxidation product, may be an endogenous PPAR γ agonist (7). Several different structural classes of PPAR γ agonists have been synthesized and these include thiazolidinediones, 2-cyano-3,12-dioxoolean-1,9-dien-28-oic acid (CDDO) and related triterpenoids, 3-benzyl and acyl-substituted indoles, substituted chromane carboxylic acids, and phosphonophosphates (8–13). Among the synthetic PPAR γ agonists, the thiazolidinediones are now being extensively used as insulin-sensitizing agents for treatment of type II diabetes (1, 3). PPAR γ is highly expressed in many tumor samples and cancer cell lines derived from hematopoietic and nonhematopoietic tumors (14), and several studies show that PPAR γ agonists inhibit growth and/or induce apoptosis in multiple cancer cell lines and in *in vivo* tumor models (15–33).

3,3'-Diindolylmethane is a metabolite of the phytochemical indole-3-carbinol and both compounds exhibit a broad spectrum of anticancer activities (34, 35). Methylene-substituted 3,3'-diindolylmethanes (C-DIM) are synthetic analogues of 3,3'-diindolylmethane, which also exhibit anticancer activity and bind to orphan receptors, such as PPAR γ and Nur77 (36–40). The 1,1-bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes containing *p*-trifluoromethyl [1,1-bis(3'-indolyl)-1-(*p*-trifluoromethyl)methane (DIM-C-pPhCF₃)], *t*-butyl [1,1-bis(3'-indolyl)-1-(*p*-*t*-butyl)methane (DIM-C-pPh_tBu)], and phenyl [1,1-bis(3'-indolyl)-1-(*p*-phenyl)methane (DIM-C-pPhC₆H₅)] substituents activate PPAR γ in several cancer cell lines, including colon cancer cells (36–39). The effects of these compounds depend on cell context and this is commonly observed for other PPAR γ agonists. The PPAR γ -active C-DIMs inhibit growth and/or induce apoptosis in cancer cells; in HT-29 and HCT-15 colon cancer cells, these compounds induce caveolin-1, whereas p21 is induced in Panc28 pancreatic cancer cells and both responses are inhibited by PPAR γ antagonists (37, 38). In contrast, induction of cyclin D1 down-regulation and apoptosis in breast cancer cells (36) and induction of nonsteroidal anti-inflammatory drug activated gene-1 (NAG-1) are PPAR γ independent (41, 42).

In this study, we show that PPAR γ -active C-DIMs inhibit growth of SW480 colon cancer cells and this is associated with PPAR γ -dependent induction of caveolin-1 and

Received 1/3/06; revised 2/2/06; accepted 3/17/06.

Grant support: NIH grants ES09106 and CA112337 and Texas Agricultural Experiment Station.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Stephen Safe, Department of Physiology and Pharmacology, Texas A&M University, 4466 TAMU, Veterinary Research Building 410, College Station, TX 77843-4466. Phone: 979-845-5988; Fax: 979-862-4929. E-mail: ssafe@cvm.tamu.edu

Copyright © 2006 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-06-0002

receptor-independent activation of both NAG-1 and poly (ADP-ribose) polymerase (PARP) cleavage. Moreover, in athymic nude mice bearing SW480 cell xenografts, DIM-C-pPhC₆H₅ (20 and 40 mg/kg/d) inhibits tumor growth and immunostaining of tumors shows both overexpression of caspase-3 and NAG-1 in the treated groups. These data show that PPAR γ -active C-DIMs inhibit colon cancer cell/tumor growth and confirm that PPAR γ agonists are an important class of compounds for potential applications in the treatment of colon cancer.

Materials and Methods

Cell Lines, Plasmids, Chemicals, and Reagents

The SW480 human colon cancer cell line was provided by Dr. S. Hamilton (M. D. Anderson Cancer Center, Houston, TX). SW480 cells were maintained in DMEM/Ham's F-12 (Sigma, St. Louis, MO) with phenol red supplemented with 0.22% sodium bicarbonate, 0.011% sodium pyruvate, and 5% fetal bovine serum (FBS) and 10 ml/L of 100 \times antibiotic-antimycotic solution (Sigma). The Gal4 reporter containing 5 \times Gal4DBD (Gal4Luc) was kindly provided by Dr. Marty Mayo (University of North Carolina, Chapel Hill, NC). Gal4DBD-PPAR γ construct was a gift of Dr. Jennifer L. Oberfield (Glaxo Wellcome Research and Development, Research Triangle Park, NC) and chimeric pM-PPAR γ coactivator-1 (PGC-1) was a gift of Dr. Bruce M. Spiegelman (Harvard University, Boston, MA). The PPAR γ 2-VP16 fusion plasmid (VP-PPAR γ) contained the DEF region of PPAR γ (amino acids 183–505) fused to the pVP16 expression vector and the Gal4-coactivator fusion plasmids pM-SRC-1, pM-AIBI, pM-TIFII, pM-DRIP205, pM-TRAP220, and pM-CARM-1 were kindly provided by Dr. Shigeaki Kato (University of Tokyo, Tokyo, Japan). Rosiglitazone was purchased from LKT Laboratories, Inc. (St. Paul, MN). Horseradish peroxidase substrate for Western blot analysis was purchased from NEN Life Science Products (Boston, MA). Cell lysis buffer and luciferase reagent were purchased from Promega (Madison, WI), and β -galactosidase reagent was from Tropix (Bedford, MA). Antibodies for cyclin D1 (sc-718), p27 (sc-528), phosphorylated Akt (sc-7985R), Akt (sc-8312), Bcl-2 (sc-7382), Bax (sc-20067), caveolin-1 (sc-894), and PARP (sc-8007) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). NAG-1 antibodies were obtained from Upstate Biotechnology (Lake Placid, NY); monoclonal anti- β -actin was purchased from Sigma.

Cell Proliferation Assay

Cells were plated at a density of 2 \times 10⁴ per well in 12-well plates in DMEM/Ham's F-12 and 5% FBS, and after 24 hours, this was replaced with DMEM/Ham's F-12 containing 2.5% charcoal-stripped FBS. Cells were then treated with either vehicle (DMSO) or the indicated ligand in DMSO. Fresh medium and compounds were added every 48 hours. Cells were counted at the indicated times using a Coulter Z1 cell counter (Beckman-Coulter, Fullerton, CA). Each experiment was carried out in triplicate and results are expressed as mean \pm SE for each determination.

Transfection and Luciferase Assay

SW480 cells were plated in 12-well plates at 1 \times 10⁵ per well in DMEM/Ham's F-12 supplemented with 2.5% charcoal-stripped FBS. After growth for 16 hours, various amounts of DNA [i.e., Gal4Luc (0.4 μ g), β -galactosidase (0.04 μ g), VP-PPAR γ (0.04 μ g), pM-SRC-1 (0.04 μ g), pM-TIFII (0.04 μ g), pM-AIBI (0.04 μ g), pM-DRIP205 (0.04 μ g), and pM-CARM-1 (0.04 μ g)] were transfected by LipofectAMINE (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. Briefly, 2 μ L LipofectAMINE was mixed with 50 μ L serum-free medium. Appropriate concentrations of plasmid DNA were mixed with 50 μ L serum-free medium. The LipofectAMINE solution was then mixed with the DNA solution and the mixture was allowed to incubate at room temperature for 45 minutes to form the DNA/LipofectAMINE complex. In the mean time, cells grown in the presence DMEM/Ham's F-12 supplemented with 2.5% charcoal-stripped FBS were washed with serum-free DMEM/Ham's F-12, and 400 μ L serum-free medium was added. Following 45-minute incubation, the LipofectAMINE/DNA complex was carefully dropped over the cells and incubated for 5 to 6 hours at 37 $^{\circ}$ C. Five hours after transfection, the transfection mix was replaced with complete medium containing either vehicle (DMSO) or the indicated ligand for 20 to 22 hours. Cells were then lysed with 100 μ L of 1 \times reporter lysis buffer, and 30 μ L cell extracts were used for luciferase and β -galactosidase assays. A Lumicount luminometer (Perkin-Elmer Life Sciences, Boston, MA) was used to quantitate luciferase and β -galactosidase activities, and the luciferase activities were normalized to β -galactosidase activity.

Western Blot Analysis

SW480 cells were seeded in DMEM/Ham's F-12 containing 2.5% charcoal-stripped FBS for 24 hours and then treated with either the vehicle (DMSO) or the indicated compounds. Whole-cell lysates were obtained using high-salt buffer [50 mmol/L HEPES, 500 mmol/L NaCl, 1.5 mmol/L MgCl₂, 1 mmol/L EGTA, 10% glycerol, 1% Triton X-100 (pH 7.5), 5 μ L/mL protease inhibitor cocktail (Sigma)].

Protein samples were incubated at 100 $^{\circ}$ C for 2 to 3 minutes, separated on 10% SDS-PAGE at 120 V for 3 to 4 hours in 1 \times running buffer [25 mmol/L Tris, 192 mmol/L glycine, 0.1% SDS (pH 8.3)], and transferred to polyvinylidene difluoride (Bio-Rad, Hercules, CA) at 0.1 V for 16 hours at 4 $^{\circ}$ C in 1 \times transfer buffer (48 mmol/L Tris-HCl, 39 mmol/L glycine, 0.025% SDS). The polyvinylidene difluoride membrane was blocked in 5% TBS-Tween 20-Blotto [10 mmol/L Tris-HCl, 150 mmol/L NaCl (pH 8.0), 0.05% Triton X-100, 5% nonfat dry milk] with gentle shaking for 30 minutes and incubated in fresh 5% TBS-Tween 20-Blotto with 1:1,000 (for CD1, p27, Bcl-2, Bax, and caveolin-1), 1:500 (for PARP and NAG-1), and 1:5,000 (for β -actin) primary antibody overnight with gentle shaking at 4 $^{\circ}$ C. After washing with TBS-Tween 20 for 10 minutes, the polyvinylidene difluoride membrane was incubated with secondary antibody (1:5,000) in 5% TBS-Tween 20-Blotto for 90 minutes. The membrane was washed with TBS-Tween

20 for 10 minutes and incubated with 10 mL chemiluminescence substrate (Perkin-Elmer Life Sciences) for 1 minute and exposed to Kodak X-OMAT AR autoradiography film (Eastman Kodak, Rochester, NY).

Xenograft Experiment

Male athymic BALB/c nude mice (age 4–6 weeks) were purchased from Harlan (Indianapolis, IN). Mice were housed and maintained in laminar flow cabinets under specific pathogen-free conditions. SW480 cells were concentrated to 2×10^6 per 200 μ L and injected s.c. into the left flank of each mouse using a 30-gauge needle. Six days after cell inoculation, animals were divided into three equal groups of 10 mice each. The first group received 70 μ L vehicle (corn oil) by oral gavage and the second and third groups of animals received 20 and 40 mg/kg/d doses of DIM-C-pPhC₆H₅ in vehicle every second day for 20 days (10 doses). The mice were weighed, and tumor areas were measured throughout the study. After 21 days, the animals were sacrificed, final body and tumor weights were determined, and selected tissues were further examined by routine H&E staining and immunohistochemical analysis.

Immunohistochemistry

Tissue sections (4–5 μ m thick) mounted on poly-L-lysine-coated slide were deparaffinized by standard methods. Endogenous peroxidase was blocked by the use of 3% hydrogen peroxide in PBS for 10 minutes. Antigen retrieval for NAG-1 staining was done for 7.5 minutes in 10 mmol/L sodium citrate buffer (pH 6) heated at 95°C in a steamer followed by cooling for 15 minutes. The slides were washed with PBS and incubated for 30 minutes at room temperature with a protein blocking solution (Biostain Rabbit IgG System, Biomedica, Foster City, CA). Excess blocking

solution was drained, and the samples were incubated overnight at 4°C with one of the following: a 1:100 dilution of activated caspase-3 antibody or a 1:100 dilution of NAG-1 antibody.

Sections were then incubated with biotinylated secondary antibody followed by streptavidin (Biostain Rabbit IgG System). The color was developed by exposing the peroxidase to diaminobenzidine reagent (Vector Laboratories, Burlingame, CA), which forms a brown reaction product. The sections were then counterstained with Gill's hematoxylin. Activated caspase-3 and NAG-1 expression was identified by the brown cytoplasmic staining.

Assessment of Apoptotic Cells

Four slides per group were stained and apoptotic cells were identified by the dark brown cytoplasmic staining. Ten microscopic fields in each slide were counted for cytoplasmic staining and averaged.

Results

Inhibition of Cell Proliferation and Activation of PPAR γ

Previous studies in colon and pancreatic cancer cells showed that PPAR γ -active C-DIMs inhibit cell growth (38, 39), and results in Fig. 1 also show the DIM-C-pPhCF₃, DIM-C-pPhTbu, and DIM-C-pPhC₆H₅ inhibit SW480 cancer cell proliferation. IC₅₀s varied between 1 and 10 μ mol/L and the relative order of potency was DIM-C-pPhCF₃ > DIM-C-pPhTbu > DIM-C-pPhC₆H₅. In contrast, 1 to 10 μ mol/L rosiglitazone did not significantly affect growth of SW480 cells.

Activation of PPAR γ was also investigated in SW480 cells transfected with chimeric PPAR γ -Gal4 and a Gal4 response




Figure 1. Cell proliferation assays. SW480 cells were treated with DIM-C-pPhCF₃ (A), DIM-C-pPhTbu (B), DIM-C-pPhC₆H₅ (C), or rosiglitazone (D), and cell numbers were determined after 2, 4, and 6 d as described in Materials and Methods. Significant ($P < 0.05$) inhibition of cell proliferation was observed for the C-DIMs ($\geq 5 \mu$ mol/L). Points, mean of three replicate determinations for each treatment group; bars, SE.

element reporter plasmid containing five-tandem Gal4 response elements linked to a luciferase reporter gene. The results showed that DIM-C-pPhCF₃, DIM-C-pPhtBu, and DIM-C-pPhC₆H₅ significantly induced transactivation with a maximal 17-fold increase observed for 10 μmol/L DIM-C-pPhCF₃, whereas <4-fold induction of luciferase activity was observed for rosiglitazone (Fig. 2A). We also investigated ligand-induced interactions of PPARγ with a panel of coactivators in a mammalian two-hybrid assay in SW480 cells (Fig. 2B). The results show that the PPARγ-active C-DIMs induce interactions between PPARγ and PGC-1 and similar results were obtained for the compounds in previous studies in HCT-15 cells, whereas in HT-29 cells PGC-1-PPARγ interactions were induced by DIM-C-pPhCF₃ and DIM-C-pPhtBu but not DIM-C-pPhC₆H₅ (37). In contrast to results obtained in HT-29/HCT-15 cells, the PPARγ-active C-DIMs also significantly induced interactions of PPARγ with SRC-1, SRC-2 (TIFII), SRC-3 (AIBI), and the corepressor SMRT, and this pattern of ligand-induced PPARγ-coactivator/corepressor interactions in SW480 cells was different from that observed in HCT-15 and HT-29 cells. These cell context-dependent differences suggest that these compounds are selective PPARγ modulators. In addition, rosiglitazone was inactive in the mammalian two-hybrid assay in SW480 cells and this assay distinguished between the activities of this compound and the PPARγ-active C-DIMs. The selectivity of these ligand-induced PPARγ-coactivator interactions indicates the potential for cell context-dependent differences in the activation of PPARγ-dependent responses by the C-DIMs.

Effects of PPARγ-Active C-DIMs on Expression of Proteins Associated with Cell Proliferation and Cell Death

The effects of two prototypical PPARγ-active C-DIMs on expression of various cell cycle proteins and apoptosis (PARP cleavage) was investigated in SW480 cells treated with 5 to 10 μmol/L DIM-C-pPhCF₃, DIM-C-pPhC₆H₅, or rosiglitazone for 24 hours (Fig. 3A). The PPARγ-active C-DIMs enhanced p27 expression and down-regulated cyclin D1 protein only at the highest concentration and this was also accompanied by PARP cleavage. Previous studies with different structural classes of PPARγ agonists have reported induction of p27 in colon and other cancer cell lines, and we have also observed cyclin D1 down-regulation and PARP cleavage in other cancer cell lines treated with C-DIM compounds (36–39, 41, 42). There were minimal compound-induced changes in Bax levels and decreased expression of Bcl-2 at the 10 μmol/L concentrations of DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅, whereas up to 10 μmol/L rosiglitazone did not affect expression of apoptotic or cell cycle proteins. The growth inhibitory effects of PPARγ-active C-DIMs in HT-29 and HCT-15 cells have been linked, in part, to induction of caveolin-1 (37) and the results in Fig. 3B show that at doses as low as 2.5 μmol/L there was induction of caveolin-1 in SW480 cells. This is a response that was not observed until 48 hours after treatment and only minimal apoptosis was induced in the




Figure 2. C-DIMs activate PPARγ. **A**, transactivation. SW480 cells were transfected with PPARγ-Gal4/pGal4 and treated with DMSO or different concentrations of compounds, and luciferase activity was determined as described in Materials and Methods. **B**, mammalian two-hybrid assay. SW480 cells were transfected with Gal4 chimeras, VP-PPARγ (LBD) and pGal4Luc, and treated with various compounds [rosiglitazone (R), DIM-C-pPhCF₃ (1), DIM-C-pPhtBu (4), and DIM-C-pPhC₆H₅ (9)], and luciferase activity was determined as described in Materials and Methods. Columns, mean of three replicate determinations for each treatment group; bars, SE. *, $P < 0.05$, significant induction.

7.5 μmol/L DIM-C-pPhCF₃ treatment group (data not shown), suggesting that caveolin-1 was primarily associated with growth inhibitory responses observed at lower concentrations (≤ 7.5 μmol/L). Induction of caveolin-1 by 5 μmol/L DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ was inhibited after cotreatment with the PPARγ antagonist GW9662 (Fig. 3C), showing that this response was PPARγ dependent, and this was observed previously in HT-29 and HCT-15 cells (37). We also recently reported that PPARγ-dependent induction of caveolin-1 was observed in SW480 cells treated with CDDO and related compounds (43) and this was accompanied by induction of phosphatidylinositol 3-kinase-dependent phosphorylation of Akt. The results in Fig. 3D also show that 5 μmol/L DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ also induce Akt phosphorylation, and in

SW480 cells cotreated with GW9662, there was a dose-dependent decrease in Akt phosphorylation. Thus, the PPAR γ -active C-DIMs coordinately induced caveolin-1 and phosphorylated Akt in SW480 cells and it has been reported that combined activation of caveolin-1 and phosphatidylinositol 3-kinase enhances the effects of agents that decrease cancer cell survival (44, 45).

The role of PPAR γ in mediating DIM-C-pPhCF $_3$ - and DIM-C-pPhC $_6$ H $_5$ -induced apoptosis was investigated in SW480 cells cotreated with C-DIMs and the PPAR γ antagonist GW9662 (Fig. 4A). The results show that GW9662 did not affect induced PARP cleavage, suggesting that this response was PPAR γ independent and similar results were observed for cyclin D1 down-regulation (data not shown). Interestingly, we also observed that at higher concentrations (>7.5 μ M/L) of C-DIMs that induce apoptosis in SW480 cells (Fig. 4A) there is a loss of caveolin-1 induction by these same compounds (Fig. 4B). The reason for this concentration-dependent switch between PPAR γ -

dependent and PPAR γ -independent responses is unknown; however, we have also observed similar responses for CDDO and related compounds (data not shown). Recent studies in this laboratory show that PPAR γ -active C-DIMs induced cell death through activation of endoplasmic reticulum stress in pancreatic cancer cells (42) or through induction of NAG-1 in HCT-116 colon cancer cells (41). Results in Fig. 4C show that induction of apoptosis in SW480 cells treated with DIM-C-pPhCF $_3$ or DIM-C-pPhC $_6$ H $_5$ for 24 hours is not accompanied by induction of the stress protein GRP78.

NAG-1 is a transforming growth factor- β -like peptide, which is induced by C-DIMs and many other growth inhibitory agents, including rosiglitazone, in selected cell lines, including HCT-116 colon cancer cells. Preliminary studies did not detect induction of NAG-1 by C-DIMs in SW480 cells (42); however, repetition of this experiment with a different antibody showed that within 24 hours after treatment with DIM-C-pPhCF $_3$ or DIM-C-pPhC $_6$ H $_5$ there




Figure 3. Cell cycle proteins, caveolin-1 expression, and Akt phosphorylation. **A**, cell cycle proteins. SW480 cells were treated with different concentrations of compounds for 24 h, and whole-cell lysates were analyzed by Western blot analysis as described in Materials and Methods. Induction of caveolin-1 (**B**, **C**), Akt phosphorylation (**D**), and inhibition by GW9662 (**C**, **D**). SW480 cells were treated with different concentrations of compounds alone or in combination, and whole-cell lysates were analyzed by Western blot analysis as described in Materials and Methods. β -Actin was used as a loading control for these experiments.




Figure 4. Effects of higher concentrations of C-DIMs on PARP cleavage (**A**) and caveolin-1 (**B**) and GRP78 (**C**) expression. SW480 cells were treated with different compounds (alone or combined) for 24 h (**A** and **C**) or 72 h (**B**), and whole-cell lysates were assayed by Western blot analysis as described in Materials and Methods. β -Actin served as a loading control for all experiments.

was induction of NAG-1 protein (Fig. 5A). Moreover, in SW480 cells cotreated with PPAR γ -active C-DIMs (5–10 $\mu\text{mol/L}$) plus the PPAR γ agonist GW9662 (10 $\mu\text{mol/L}$), the antagonist did not block induction of NAG-1 (Fig. 5B), suggesting that this response was PPAR γ independent as reported previously in HCT-116 cells (42). We also observed induction of NAG-1 after treatment with C-DIMs for 96 hours (Fig. 5C), and at both 24- and 96-hour time points, effective doses were in the 7.5 to 10 $\mu\text{mol/L}$ range.

The *in vivo* antitumor activity of PPAR γ -active C-DIMs was further investigated in male athymic nude mice bearing SW480 cell xenografts. After palpable tumors were first observed (<15 mm²), animals were treated with corn oil (70 $\mu\text{L}/\text{mouse}$) or DIM-C-pPhC₆H₅ (20 or 40 mg/kg) in corn oil by oral gavage every second day for 21 days and tumor areas were determined. The results (Fig. 6A) show that tumor areas were significantly decreased after treatment with DIM-C-pPhC₆H₅ compared with the corn oil (control) group, and after sacrifice, tumor weights were also decreased in both treatment groups (Fig. 6B). We also investigated the effects of DIM-C-pPhC₆H₅ (40 mg/kg/d) on expression of the apoptotic gene caspase-3 (Fig. 6C and D) and NAG-1 in tumors by immunohistochemical analysis. DIM-C-pPhC₆H₅ significantly induced caspase-3 expression compared with control (corn oil) tumors (Fig. 6C). NAG-1 was not detected in tumors from control animals, whereas massive expression (brown staining) of NAG-1 was detected in tumors from DIM-C-pPhC₆H₅-treated mice (Fig. 6D),

suggesting an important role for this protein in the antitumor activity of this C-DIM compound. In contrast, minimal NAG-1 expression was observed at the 20 mg/kg dose and the available caveolin-1 antibodies did not give consistent staining. These data complement the *in vitro* studies and show that PPAR γ -active C-DIMs inhibit colon cancer cell/tumor growth and these compounds are being developed for future clinical applications.

Discussion

Several studies report that PPAR γ agonists inhibit growth of colon cancer cells; however, their induced responses and mechanisms of action are dependent on multiple factors, including ligand structure, cell context, and wild-type or variant PPAR γ expression. For example, troglitazone (a thiazolidinedione), 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂, and PPAR γ -active C-DIMs induce the transforming growth factor- β -like peptide NAG-1 in HCT-116 cells and GW9662 inhibits the effects of 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ but had no effect on troglitazone or the PPAR γ -active C-DIMs (27, 41). Although troglitazone, DIM-C-pPhCF₃, DIM-C-pPhTbu, and DIM-C-pPhC₆H₅ induce NAG-1 through prior activation of early growth response gene (27, 41), the former response is dependent on activation of mitogen-activated protein kinase, whereas the C-DIMs activate EGR-1 and NAG-1 through induction of phosphatidylinositol 3-kinase. Dubois and coworkers (20, 21, 23) have reported that the effects of rosiglitazone on cell proliferation and differentiation in several colon cancer cell lines




Figure 5. Induction of NAG-1 by C-DIM compounds. SW480 cells were treated for 24 h with DIM-C-pPhCF₃ or DIM-C-pPhC₆H₅ alone (**A**) or in combination with GW9662 (**B**) or for 96 h (**C**). Whole-cell lysates were analyzed for NAG-1 expression by Western blot analysis as described in Materials and Methods. β -Actin was used as a loading control for these experiments.




Figure 6. *In vivo* antitumor activity of DIM-C-pPhC₆H₅. Tumor areas (A) and weights (B). When palpable tumors were first observed (15 mm²), athymic nude mice bearing SW480 colon cancer cell xenografts were given corn oil (control) or DIM-C-pPhC₆H₅ (20 or 40 mg/kg/d) in corn oil by oral gavage, and tumor areas and weights were determined as described in Materials and Methods. Points (A) and columns (B), mean for 10 animals for each treatment group; bars, SE. *, $P < 0.05$, significant decrease in tumor areas and weights in the treatment groups compared with the corn oil (control) group. C, caspase-3 expression was quantitated by immunohistochemical analysis of caspase-3 in tumor sections from control and treated (DIM-C-pPhC₆H₅, 40 mg/kg/d) mice as described in Materials and Methods. Arrows, positive caspase-3 staining. D, immunohistochemical analysis of NAG-1. Tumors from control or DIM-C-pPhC₆H₅-treated mice were analyzed for NAG-1 expression by immunohistochemistry as described in Materials and Methods. There were no significant differences in animal weight gain, organ weights, or histopathology in any of the treatment groups.

were dependent on their expression of wild-type or mutant K422Q PPAR γ where rosiglitazone responsiveness was observed only in cells (e.g., HT-29) expressing wild-type receptor (21).

PPAR γ -active C-DIMs inhibit growth of HT-29 and HCT-15 colon cancer cell lines expressing wild-type and mutant PPAR γ , respectively, and these compounds also induce

caveolin-1 in both cell lines (37). The results in Figs. 1 and 2 show that the C-DIM compounds but not rosiglitazone inhibit SW480 cancer cell growth and induced PPAR γ -dependent transactivation. These data are similar to those previously reported in rosiglitazone-nonresponsive HCT-15 cells, suggesting that SW480 cells may also express mutant PPAR γ , which is not responsive to the growth inhibitory effects of rosiglitazone.

The short-term effects of DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ on cell cycle protein expression and apoptosis in SW480 (Fig. 4) versus HT-29/HCT-15 cells (37) were different. Growth inhibitory concentrations (5.0 and 7.5 μ mol/L) did not affect p21 (data not shown) or cyclin D1 expression in SW480, HT-29, or HCT-15 cells, whereas p27 protein levels were slightly elevated in SW480 cells after treatment with 7.5 μ mol/L DIM-C-pPhCF₃ and 10 μ mol/L DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅. Another major difference observed in this study with SW480 cells was the induction of apoptosis and down-regulation of cyclin D1 after treatment with 10 μ mol/L DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ for 24 hours, whereas these responses were not observed in HT-29 or HCT-15 cells (37).

Despite the cell context-dependent differences in the effects of PPAR γ -active C-DIMs in colon cancer cells, the results in Fig. 3 show that PPAR γ -active C-DIMs induce caveolin-1 expression in SW480 cells after prolonged treatment (3 days) with concentrations as low as 2.5 μ mol/L. This response in SW480 cells was inhibited by GW9662 and similar results were observed for PPAR γ -active C-DIMs and for CDDO and related compounds in colon cancer cells (37, 43). In this study, we also observed PPAR γ -dependent up-regulation of phosphorylated Akt in SW480 cells and similar results have been observed for CDDO and related compounds in the same cell line (43). The induction of caveolin-1 at the lower doses (<7.5 μ mol/L) of DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ is consistent with the growth inhibitory effects of these compounds because caveolin-1 expression has been linked to inhibition of colon cancer cell/tumor proliferation in both *in vitro* and *in vivo* models (44, 45). The observation that DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ induce phosphatidylinositol 3-kinase-dependent phosphorylation of Akt, which is normally a cell survival pathway, was also consistent with the observed growth inhibitory effects of these compounds. Phosphatidylinositol 3-kinase and caveolin-1 coexpression sensitizes HeLa and 293 cells to the cytotoxicity of arsenite and hydrogen peroxide (46) and sensitizes L929 cells to tumor necrosis factor α -induced cell death (47).

Several studies show that PPAR γ agonists induce both receptor-dependent and receptor-independent responses (26, 27, 29, 36–39, 41–43), and in this study, these pathways are separable at different concentrations of C-DIMs. At low concentrations (\leq 7.5 μ mol/L), DIM-C-pPhCF₃ and DIM-C-pPhC₆H₅ induce receptor-dependent up-regulation of caveolin-1 (Fig. 3), whereas at higher concentrations (>7.5 μ mol/L) these same compounds induce apoptosis, which was not inhibited by GW9662, a PPAR γ antagonist (Fig. 4A). Similar results were observed for CDDO and

related compounds in SW480 cells (43). Interestingly, there seems to be a narrow concentration range for the switch between receptor-dependent and receptor-independent responses. Moreover, induction of PPAR γ -dependent caveolin-1 expression by ≥ 7.5 $\mu\text{mol/L}$ DIM-C-pPhCF $_3$ or DIM-C-pPhC $_6$ H $_5$ is lost in cells treated with 10 $\mu\text{mol/L}$ concentrations of these same compounds (Fig. 4B) where extensive apoptosis is observed. In other cell lines, we have recently shown that C-DIMs induce endoplasmic reticulum stress, which leads to activation of DR5 and the extrinsic pathway for apoptosis (41); however, this response was not observed in SW480 cells (Fig. 4C). In contrast, both DIM-C-pPhC $_6$ H $_5$ and DIM-C-pPhCF $_3$ induced NAG-1 expression at concentrations (≥ 7.5 $\mu\text{mol/L}$) similar to those required for induction of apoptosis (PARP cleavage) and both responses were PPAR γ independent. NAG-1 is up-regulated by different classes of compounds that inhibit growth and induce apoptosis in colon and other cancer cell lines (29, 41, 48–51), and a recent study showed that inhibition of NAG-1 induction by sulindac sulfide using RNA interference also reversed the growth inhibitory effects of this compound in SKOV3 cells (50). Moreover, overexpression of NAG-1 in breast cancer cells significantly decreased cell viability (52), and purified NAG-1 induced apoptosis in prostate cancer cells (53). These results suggest that PPAR γ -independent induction of apoptosis in SW480 cells by C-DIM compounds is due to up-regulation of NAG-1 expression. These *in vitro* responses are also paralleled in the athymic nude mouse xenograft studies where DIM-C-pPhC $_6$ H $_5$ not only inhibits tumor growth but also induces caspase-3 and NAG-1 expression in the tumors (Fig. 6).

In summary, our results show that PPAR γ -active C-DIMs inhibit SW480 cell proliferation and inhibit colon tumor growth in an athymic nude mouse xenograft model (Fig. 6). The antitumorigenic activity of these compounds may be due to activation of both PPAR γ -dependent (caveolin-1) or PPAR γ -independent (NAG-1) responses; however, the relative contributions of these pathways to the antitumorigenic action of the C-DIMs is unknown. Ongoing studies using RNA interference (*in vitro*) and transgenic mouse models (*in vivo*) will provide critical insights on the relative importance of the receptor-dependent and receptor-independent responses to the antitumorigenic activity of C-DIMs and guide development of more effective C-DIM analogues for applications in colon cancer chemotherapy.

References

- Rosen ED, Spiegelman BM. PPAR γ : a nuclear regulator of metabolism, differentiation, and cell growth. *J Biol Chem* 2001;276:37731–4.
- Willson RM, Lambert MH, Kliewer SA. Peroxisome proliferator-activated receptor γ and metabolic disease. *Annu Rev Biochem* 2001;70:341–67.
- Lee CH, Olson P, Evans RM. Minireview: lipid metabolism, metabolic diseases, and peroxisome proliferator-activated receptors. *Endocrinology* 2003;144:2201–7.
- Desvergne B, Wahli W. Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 1999;20:649–88.
- Escher P, Wahli W. Peroxisome proliferator-activated receptors: insight into multiple cellular functions. *Mutat Res* 2000;448:121–38.
- Fajas L, Debril MB, Auwerx J. Peroxisome proliferator-activated receptor- γ : from adipogenesis to carcinogenesis. *J Mol Endocrinol* 2001;27:1–9.
- Schopfer FJ, Lin Y, Baker PR, et al. Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor γ ligand. *Proc Natl Acad Sci U S A* 2005;102:2340–5.
- Rieusset J, Touri F, Michalik L, et al. A new selective peroxisome proliferator-activated receptor γ antagonist with antiobesity and antidiabetic activity. *Mol Endocrinol* 2002;16:2628–44.
- Berger JP, Petro AE, Macnaul KL, et al. Distinct properties and advantages of a novel peroxisome proliferator-activated protein γ selective modulator. *Mol Endocrinol* 2003;17:662–76.
- Koyama H, Miller DJ, Boueres JK, et al. (2R)-2-Ethylchromane-2-carboxylic acids: discovery of novel PPAR α/γ dual agonists as antihyperglycemic and hypolipidemic agents. *J Med Chem* 2004;47:3255–63.
- Acton JJ III, Black RM, Jones AB, et al. Benzoyl 2-methyl indoles as selective PPAR γ modulators. *Bioorg Med Chem Lett* 2005;15:357–62.
- Liu K, Black RM, Acton JJ III, et al. Selective PPAR γ modulators with improved pharmacological profiles. *Bioorg Med Chem Lett* 2005;15:2437–40.
- Suh N, Wang Y, Honda T, et al. A novel synthetic oleanane triterpenoid, 2-cyano-3,12-dioxolean-1,9-dien-28-oic acid, with potent differentiating, antiproliferative, and anti-inflammatory activity. *Cancer Res* 1999;59:336–41.
- Ikezoe T, Miller CW, Kawano S, et al. Mutational analysis of the peroxisome proliferator-activated receptor γ gene in human malignancies. *Cancer Res* 2001;61:5307–10.
- Place AE, Suh N, Williams CR, et al. The novel synthetic triterpenoid, CDDO imidazolide, inhibits inflammatory response and tumor growth *in vivo*. *Clin Cancer Res* 2003;9:2798–806.
- Qin C, Burghardt R, Smith R, et al. Peroxisome proliferator-activated receptor γ (PPAR γ) agonists induce proteasome-dependent degradation of cyclin D1 and estrogen receptor α in MCF-7 breast cancer cells. *Cancer Res* 2003;63:958–64.
- Motomura W, Okumura T, Takahashi N, Obara T, Kohgo Y. Activation of peroxisome proliferator-activated receptor γ by troglitazone inhibits cell growth through the increase of p27^{KIP1} in human pancreatic carcinoma cells. *Cancer Res* 2000;60:5558–64.
- Chang TH, Szabo E. Induction of differentiation and apoptosis by ligands of peroxisome proliferator-activated receptor γ in non-small cell lung cancer. *Cancer Res* 2000;60:1129–38.
- Elstner E, Muller C, Koshizuka K, et al. Ligands for peroxisome proliferator-activated receptor γ and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells *in vitro* and in BNX mice. *Proc Natl Acad Sci U S A* 1998;95:8806–11.
- Gupta RA, Sarraf P, Mueller E, et al. Peroxisome proliferator-activated receptor γ -mediated differentiation: a mutation in colon cancer cells reveals divergent and cell type-specific mechanisms. *J Biol Chem* 2003;278:22669–77.
- Brockman JA, Gupta RA, DuBois RN. Activation of PPAR γ leads to inhibition of anchorage independent growth of human colorectal cancer cells. *Gastroenterology* 1998;115:1049–55.
- Kato M, Kusumi T, Tsuchida S, et al. Induction of differentiation and peroxisome proliferator-activated receptor γ expression in colon cancer cell lines by troglitazone. *J Cancer Res Clin Oncol* 2004;130:73–9.
- Gupta RA, Brockman JA, Sarraf P, Willson TM, DuBois RN. Target genes of peroxisome proliferator-activated receptor γ in colorectal cancer cells. *J Biol Chem* 2001;276:29681–7.
- Kitamura S, Miyazaki Y, Shinomura Y, et al. Peroxisome proliferator-activated receptor γ induces growth arrest and differentiation markers of human colon cancer cells. *Jpn J Cancer Res* 1999;90:75–80.
- Takahashi N, Okumura T, Motomura W, et al. Activation of PPAR γ inhibits cell growth and induces apoptosis in human gastric cancer cells. *FEBS Lett* 1999;455:135–9.
- Clay CE, Monjavez A, Thorburn J, Chilton FH, High KP. 15-Deoxy- Δ (12,14)-prostaglandin J $_2$ -induced apoptosis does not require PPAR γ in breast cancer cells. *J Lipid Res* 2002;43:1818–28.
- Baek SJ, Kim JS, Nixon JB, DiAugustine RP, Eling TE. Expression of NAG-1, a transforming growth factor- β superfamily member, by troglitazone requires the early growth response gene *EGR-1*. *J Biol Chem* 2004;279:6883–92.

28. Inoue K, Kawahito Y, Tsubouchi Y, et al. Expression of peroxisome proliferator-activated receptor γ in renal cell carcinoma and growth inhibition by its agonists. *Biochem Biophys Res Commun* 2001;287:727–32.
29. Baek SJ, Wilson LC, Hsi LC, Eling TE. Troglitazone, a peroxisome proliferator-activated receptor γ (PPAR γ) ligand, selectively induces the early growth response-1 gene independently of PPAR γ . A novel mechanism for its anti-tumorigenic activity. *J Biol Chem* 2003;278:5845–53.
30. Niho N, Takahashi M, Kitamura T, et al. Concomitant suppression of hyperlipidemia and intestinal polyp formation in *Apc*-deficient mice by peroxisome proliferator-activated receptor ligands. *Cancer Res* 2003;63:6090–5.
31. Sarraf P, Mueller E, Jones D, et al. Differentiation and reversal of malignant changes in colon cancer through PPAR γ . *Nat Med* 1998;4:1046–52.
32. Osawa E, Nakajima A, Wada K, et al. Peroxisome proliferator-activated receptor γ ligands suppress colon carcinogenesis induced by azoxymethane in mice. *Gastroenterology* 2003;124:361–7.
33. Niho N, Takahashi M, Shoji Y, et al. Dose-dependent suppression of hyperlipidemia and intestinal polyp formation in Min mice by pioglitazone, a PPAR γ ligand. *Cancer Sci* 2003;94:960–4.
34. Sarkar FH, Li Y. Indole-3-carbinol and prostate cancer. *J Nutr* 2004;134:3493–8S.
35. Shertzer HG, Senft AP. The micronutrient indole-3-carbinol: implications for disease and chemoprevention. *Drug Metabol Drug Interact* 2000;17:159–88.
36. Qin C, Morrow D, Stewart J, et al. A new class of peroxisome proliferator-activated receptor γ (PPAR γ) agonists that inhibit growth of breast cancer cells: 1,1-bis(3'-indolyl)-1-(*p*-substitutedphenyl)methanes. *Mol Cancer Ther* 2004;3:247–59.
37. Chintharlapalli S, Smith III R, Samudio I, Zhang W, Safe S. 1,1-Bis(3'-indolyl)-1-(*p*-substitutedphenyl)methanes induce peroxisome proliferator-activated receptor γ -mediated growth inhibition, transactivation and differentiation markers in colon cancer cells. *Cancer Res* 2004;64:5994–6001.
38. Hong J, Samudio I, Liu S, Abdelrahim M, Safe S. Peroxisome proliferator-activated receptor γ -dependent activation of p21 in Panc-28 pancreatic cancer cells involves Sp1 and Sp4 proteins. *Endocrinology* 2004;145:5774–85.
39. Contractor R, Samudio I, Estrov Z, et al. A novel ring-substituted diindolylmethane 1,1-bis[3'-(5-methoxyindolyl)]-1-(*p*-*t*-butylphenyl)methane inhibits ERK activation and induces apoptosis in acute myeloid leukemia. *Cancer Res* 2005;65:2890–8.
40. Chintharlapalli S, Burghardt R, Papineni S, et al. Activation of Nur77 by selected 1,1-bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes induces apoptosis through nuclear pathways. *J Biol Chem* 2005;280:24903–14.
41. Chintharlapalli S, Papineni S, Baek SJ, Liu S, Safe S. 1,1-Bis(3'-indolyl)-1-(*p*-substitutedphenyl)methanes are peroxisome proliferator-activated receptor γ agonists but decrease HCT-116 colon cancer cell survival through receptor-independent activation of early growth response-1 and NAG-1. *Mol Pharmacol* 2005;68:1782–92.
42. Abdelrahim M, Newman K, Vanderlaag K, Samudio I, Safe S. 3,3'-Diindolylmethane (DIM) and derivatives induce apoptosis in pancreatic cancer cells through endoplasmic reticulum stress-dependent upregulation of DR5. *Carcinogenesis* 2006;27:717–28.
43. Chintharlapalli S, Papineni S, Konopleva M, et al. 2-Cyano-3,12-dioxoolean-1,9-dien-28-oic acid and related compounds inhibit growth of colon cancer cells through peroxisome proliferator-activated receptor γ -dependent and -independent pathways. *Mol Pharmacol* 2005;68:119–28.
44. Bender FC, Reymond MA, Bron C, Quest AF. Caveolin-1 levels are down-regulated in human colon tumors, and ectopic expression of caveolin-1 in colon carcinoma cell lines reduces cell tumorigenicity. *Cancer Res* 2000;60:5870–8.
45. Burgermeister E, Tencer L, Liscovitch M. Peroxisome proliferator-activated receptor- γ upregulates caveolin-1 and caveolin-2 expression in human carcinoma cells. *Oncogene* 2003;22:3888–900.
46. Shack S, Wang XT, Kokkonen GC, et al. Caveolin-induced activation of the phosphatidylinositol 3-kinase/Akt pathway increases arsenite cytotoxicity. *Mol Cell Biol* 2003;23:2407–14.
47. Ono K, Iwanaga Y, Hirayama M, et al. Contribution of caveolin-1 α and Akt to TNF- α -induced cell death. *Am J Physiol Lung Cell Mol Physiol* 2004;287:L201–9.
48. Baek SJ, Kim KS, Nixon JB, Wilson LC, Eling TE. Cyclooxygenase inhibitors regulate the expression of a TGF- β superfamily member that has proapoptotic and antitumorigenic activities. *Mol Pharmacol* 2001;59:901–8.
49. Baek SJ, Wilson LC, Eling TE. Resveratrol enhances the expression of nonsteroidal anti-inflammatory drug-activated gene (*NAG-1*) by increasing the expression of p53. *Carcinogenesis* 2002;23:425–34.
50. Kim JS, Baek SJ, Sali T, Eling TE. The conventional nonsteroidal antiinflammatory drug sulindac sulfide arrests ovarian cancer cell growth via the expression of *NAG-1/MIC-1/GDF-15*. *Mol Cancer Ther* 2005;4:487–93.
51. Bottone FG, Jr., Baek SJ, Nixon JB, Eling TE. Diallyl disulfide (DADS) induces the antitumorigenic NSAID-activated gene (*NAG-1*) by a p53-dependent mechanism in human colorectal HCT 116 cells. *J Nutr* 2002;132:773–8.
52. Li PX, Wong J, Ayed A, et al. Placental transforming growth factor- β is a downstream mediator of the growth arrest and apoptotic response of tumor cells to DNA damage and p53 overexpression. *J Biol Chem* 2000;275:20127–35.
53. Liu T, Bauskin AR, Zaunders J, et al. Macrophage inhibitory cytokine 1 reduces cell adhesion and induces apoptosis in prostate cancer cells. *Cancer Res* 2003;63:5034–40.

Molecular Cancer Therapeutics

1,1-Bis(3'-indolyl)-1-(*p*-substituted phenyl)methanes inhibit colon cancer cell and tumor growth through PPAR γ -dependent and PPAR γ -independent pathways

Sudhakar Chintharlapalli, Sabitha Papineni and Stephen Safe

Mol Cancer Ther 2006;5:1362-1370.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/5/5/1362>

Cited articles This article cites 52 articles, 31 of which you can access for free at:
<http://mct.aacrjournals.org/content/5/5/1362.full#ref-list-1>

Citing articles This article has been cited by 15 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/5/5/1362.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.