

## Research Article

# Cannabidiol Enhances the Inhibitory Effects of $\Delta^9$ -Tetrahydrocannabinol on Human Glioblastoma Cell Proliferation and Survival

Jahan P. Marcu<sup>1</sup>, Rigel T. Christian<sup>1</sup>, Darryl Lau<sup>1</sup>, Anne J. Zielinski<sup>1</sup>, Maxx P. Horowitz<sup>1</sup>, Jasmine Lee<sup>1</sup>, Arash Pakdel<sup>1</sup>, Juanita Allison<sup>1</sup>, Chandani Limbad<sup>1</sup>, Dan H. Moore<sup>1,2</sup>, Garret L. Yount<sup>1</sup>, Pierre-Yves Desprez<sup>1</sup>, and Sean D. McAllister<sup>1</sup>

## Abstract

The cannabinoid 1 (CB<sub>1</sub>) and cannabinoid 2 (CB<sub>2</sub>) receptor agonist  $\Delta^9$ -tetrahydrocannabinol (THC) has been shown to be a broad-range inhibitor of cancer in culture and *in vivo*, and is currently being used in a clinical trial for the treatment of glioblastoma. It has been suggested that other plant-derived cannabinoids, which do not interact efficiently with CB<sub>1</sub> and CB<sub>2</sub> receptors, can modulate the actions of  $\Delta^9$ -THC. There are conflicting reports, however, as to what extent other cannabinoids can modulate  $\Delta^9$ -THC activity, and most importantly, it is not clear whether other cannabinoid compounds can either potentiate or inhibit the actions of  $\Delta^9$ -THC. We therefore tested cannabidiol, the second most abundant plant-derived cannabinoid, in combination with  $\Delta^9$ -THC. In the U251 and SF126 glioblastoma cell lines,  $\Delta^9$ -THC and cannabidiol acted synergistically to inhibit cell proliferation. The treatment of glioblastoma cells with both compounds led to significant modulations of the cell cycle and induction of reactive oxygen species and apoptosis as well as specific modulations of extracellular signal-regulated kinase and caspase activities. These specific changes were not observed with either compound individually, indicating that the signal transduction pathways affected by the combination treatment were unique. Our results suggest that the addition of cannabidiol to  $\Delta^9$ -THC may improve the overall effectiveness of  $\Delta^9$ -THC in the treatment of glioblastoma in cancer patients.

*Mol Cancer Ther*; 9(1): 180-9. ©2010 AACR.

## Introduction

$\Delta^9$ -Tetrahydrocannabinol ( $\Delta^9$ -THC) and other cannabinoids can act as direct anticancer agents in multiple types of cancer in culture and *in vivo* (1). Specifically, activation of the two cloned cannabinoid receptors, CB<sub>1</sub> and CB<sub>2</sub>, by  $\Delta^9$ -THC can lead to the inhibition of cell proliferation, invasion, and induction of apoptosis in cancer cell lines, resulting in the reduction of tumor burden *in vivo* (2-4). The promising preclinical therapeutic potential of  $\Delta^9$ -THC, as an inhibitor of glioblastoma, has prompted a human clinical trial (5).

The CB<sub>1</sub> and CB<sub>2</sub> receptors are members of the G-protein coupled receptor (GPCR) superfamily, and can inter-

act with five structurally distinct classes of compounds. These include the plant-derived classical cannabinoids, such as  $\Delta^9$ -THC; the nonclassical bicyclic cannabinoids, such as CP55,940; the endogenous cannabinoids, such as anandamide; the aminoalkylindoles, such as WIN55,212-2; and the antagonist/inverse agonists, such as SR141716A (6). Interaction sites, independent of CB<sub>1</sub> and CB<sub>2</sub> receptors, also seem to be responsible for the anticancer activity of cannabinoids (7-10). There are >60 cannabinoids in *Cannabis sativa*. In addition to  $\Delta^9$ -THC, cannabidiol, cannabinol, and cannabigerol are also present in the plant (11). Cannabinol has low affinity for CB<sub>1</sub> and CB<sub>2</sub> receptors, whereas the nonpsychotropic cannabinoids, cannabidiol and cannabigerol, have negligible affinity for the cloned receptors (12-14). Whereas cannabinol and cannabigerol have not been tested for their ability to inhibit human brain cancer, cannabidiol has been reported to inhibit the growth of a human glioblastoma in a xenograft model (7, 15, 16).

There are conflicting reports as to what extent other cannabinoids can modulate the activity of  $\Delta^9$ -THC, and it has been suggested that nonpsychoactive cannabinoids can either potentiate or inhibit the actions of  $\Delta^9$ -THC (17-20). Cooperative effects have also been observed with endogenous cannabinoids (21). The potential benefits of using a cannabinoid-based medicine comprising multiple cannabinoids has been a driving force in recent

**Authors' Affiliations:** <sup>1</sup>California Pacific Medical Center Research Institute and <sup>2</sup>Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, California

**Note:** Supplementary material for this article is available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

J.P. Marcu and R.T. Christian contributed equally.

**Corresponding Author:** Sean D. McAllister, California Pacific Medical Center Research Institute, 475 Brannan Street, Suite 220, San Francisco, CA 94107. Phone: 415-600-5926; Fax: 415-600-5970. E-mail: [mcallis@cpmcri.org](mailto:mcallis@cpmcri.org)

doi: 10.1158/1535-7163.MCT-09-0407

©2010 American Association for Cancer Research.

human clinical trials (20, 22, 23). Investigations have shown that nonpsychoactive cannabinoids can alter the physiologic response to  $\Delta^9$ -THC, potentially by altering its metabolism (17–19, 24, 25). However, no investigation to date has provided the molecular mechanisms to explain how cannabinoids, acting through distinct pathways, could converge onto a shared pathway resulting in a modulation of activity unique to the combination.

In this study, we sought to determine whether cannabidiol, the plant-derived cannabinoid, would modulate the ability of  $\Delta^9$ -THC to inhibit glioblastoma cell proliferation and survival. We found that cannabidiol enhanced the ability of  $\Delta^9$ -THC to inhibit glioblastoma cell growth and induce apoptosis. The molecular mechanisms associated with these specific effects are presented.

## Material and Methods

### Cell Culture and Treatments

The human glioblastoma cell lines used were SF126, U251, and U87. The cell lines were maintained at 37°C and 5% CO<sub>2</sub>. In all experiments, the different cell populations were first cultured in RPMI media containing 10% fetal bovine serum. Glioblastoma cells were then seeded into 96-well plates in 10% fetal bovine serum and on the first day of treatment the media were replaced with vehicle control or drug in RPMI and 0.1% fetal bovine serum as previously reported (8). The media with the appropriate compounds were replaced every 24 h.  $\Delta^9$ -THC and cannabidiol were obtained from NIH through the National Institute of Drug Abuse.

### MTT Assay

Assays were done as previously described (26). Percent control was calculated as the MTT absorbance of the treated cells/control cells  $\times 100$ .

### Apoptosis Analysis

Cells were grown in 6-well culture dishes and treated with the appropriate compounds every 24 h for 3 d. Cells attached to the plate and the cells in media were collected, pelleted, washed once with PBS, and processed for labeling with FITC-tagged annexin and propidium iodide (PI) by use of an Apo-Direct apoptosis kit obtained from Phoenix Flow Systems. Briefly, the cell pellet was resuspended in 300  $\mu$ L of the supplied reaction buffer along with 3  $\mu$ L of both PI and FITC-tagged annexin. After a 15-min incubation period at room temperature, the labeled cells were analyzed by flow cytometry using a FITC detector (FL1) and a PI emission signal detector (FL2). Cell flow cytometry in combination with PI and annexin staining was used to quantify the percentage of cells undergoing apoptosis in control and treatment groups. Percent control was calculated as annexin-positive staining in treated cells/control cells  $\times 100$ . PI staining was used to distinguish necrotic cells from those undergoing apoptosis.

### Cell Cycle Analysis

U251 cells were grown in Petri dishes (100 mm  $\times$  15 mm) and received drug treatments for 2 d. On the third day, the cells were harvested and centrifuged at 1,200 rpm for 5 min. The pellet was washed once with PBS plus 1% bovine serum albumin, and centrifuged again. The pellet was resuspended in 0.5 mL of 2% paraformaldehyde and fixed overnight at room temperature. The next day the cells were pelleted and resuspended in 0.5 mL 0.3% Triton in PBS and incubated for 5 min at room temperature. The cells were then washed twice with PBS plus 1% bovine serum albumin. The cells were finally suspended in PBS (0.1% bovine serum albumin) with 10  $\mu$ g/mL PI and 100  $\mu$ g/mL RNase. The cells were incubated for 30 min at room temperature before being stored at 4°C. Cell cycle was measured using a Fluorescence Activated Cell Sorting (FACS) Calibur, Cell Quest Pro and Modfit software.

### Boyden Chamber Invasion Assay

Assays were done in modified Boyden Chambers (BD Biosciences) as previously described (26). Data were presented as relative invasiveness of the cells through the Matrigel, where the respective controls were set as 100%.

### Reactive Oxygen Species Measurements

The production of cellular reactive oxygen species (ROS)/H<sub>2</sub>O<sub>2</sub> was measured using 2'-7'-Dichlorodihydrofluorescein (DCFH-DA; Sigma Aldrich). DCFH-DA is deacylated intracellularly into a nonfluorescent product, which reacts with intracellular ROS to produce 2'-7'-Dichlorofluorescein, which remains trapped inside the cell, and can be measured quantitatively. Cells were plated onto 6-well dishes and received drug treatments for 3 d. On the third day, 10 mmol/L DCFH-DA was added to the media (RPMI with 0.1% fetal bovine serum) and the cells were incubated with DCFH-DA overnight. The next day, the cells were trypsinized, washed with PBS, and the fluorescent intensity was measured using FACS and Cell Quest Pro software.

### Western Analysis

Western analysis was done as previously described (26). Anti-phospho-JNK, anti-phospho-p38, anti-phospho-ERK1/2, and anti-ERK1/2 were obtained from Millipore. Anti-cleaved caspase 3, 7, 9 and poly(ADP-ribose) polymerase (PARP) were obtained from Cell Signaling. Antibodies were added according to the manufacturer's protocol.

### PCR

Total cellular RNA was isolated from glioblastoma cancer cells treated with vehicle control or with cannabidiol. Transcripts for *p8* and for  $\beta$ -*actin* were reverse-transcribed using SuperscriptII Reverse TranscriptaseII (Gibco-BRL), and PCR was done. The 5' and 3' PCR primers were GAAGAGAGGCAGGGAAGACA and CTGCCGTGCGTGTCTATTTA for *p8*; and GCGGAAATCGTGCGTGACATT and GATGGAGTTGAAGGTAGTTTCGTG

for  $\beta$ -actin. PCR was done in buffer containing 1  $\mu$ mol/L of each of the 5' and 3' PCR primers and 0.5 U of Taq polymerase using 18 cycles for amplification of *p8* and  $\beta$ -actin cDNAs. The cycle conditions were 45 s denaturation at 94°C, 45 s annealing at 55°C, and 1 min extension at 72°C.

### Pharmacologic and Statistical Analyses

In the proliferation assays,  $IC_{50}$  values with corresponding 95% confidence limits were calculated using nonlinear analysis of logged data (GraphPad Prism). When just the confidence limits of the  $IC_{50}$  values overlapped significant differences were determined using unpaired Student's *t*-test. Significant differences were also determined using one-way ANOVA where suitable. Bonferroni's multiple comparison posthoc analyses were conducted when appropriate. *P* values <0.05 defined statistical significance. Positive and negative aspects of constituent interaction were determined in a 2 × 2 design using two-way ANOVA as previously described (27).  $IC_{20}$  and  $IC_{80}$  values were calculated using the equation  $IC_F = (F/100-F)^{1/H} \times IC_{50}$ , where *F* is the fractional response expressed as a percentage,  $IC_F$  is the quantity of drug needed to inhibit an *F* percentage response, and *H* is the hillslope. Treatment groups were divided into (a) no treatment (control), (b)  $\Delta^9$ -THC alone, (c) cannabidiol alone, and (d)  $\Delta^9$ -THC and cannabidiol combined. Data were analyzed using a two-way ANOVA with an interaction term that was used to test for whether the combination of  $\Delta^9$ -THC and cannabidiol differed from the additive effects of each alone (GraphPad Prism).

To further test for synergism, the combination index (CI) was calculated where  $CI < 1$ ,  $= 1$ , and  $> 1$  indicate synergism, additive effect, and antagonism, respectively (28, 29). Based on the classic isobologram for mutually exclusive effects relative to the end point of measurement, the CI value for *x*% inhibition is calculated as:  $CI = (D)_1 / (Dx)_1 + (D)_2 / (Dx)_2$ ; where (D)<sub>1</sub> and (D)<sub>2</sub> represent  $\Delta^9$ -THC and CBD, respectively, and (Dx)<sub>1</sub> and (Dx)<sub>2</sub> are the doses for *x*% growth that can be obtained using the  $IC_F$  equation described above, and (D)<sub>1</sub> and (D)<sub>2</sub> are the concentrations in the combination that also inhibit cell growth by *x*% (29).

A cell cycle analysis program was used to estimate the proportions of cells in each of three compartments:  $G_0$ - $G_1$ , *S*, and  $G_2$ - $G_M$ . The experiments were conducted on four different dates and there were two replicates for each date. It was noticed that in the control (vehicle) experiments the percentage of cells in each compartment varied significantly from day to day, therefore each treatment compartment percentage estimate was standardized by dividing it by the average percentage for the vehicle on that date. This procedure was carried out for data from each experiment on each day. Multivariate analysis of variance (MANOVA) on the vector of cell cycle compartment standardized ratios ( $G_0$ - $G_1$ , *S*, and  $G_2$ - $G_M$ ) was done to control the multiple comparison type 1 (false positive) error rate. Because this produced a sig-

nificant result (at *P* < 0.05), it was concluded that there were differences due to treatments. Standardized ratios for each compartment were then tested separately using univariate one-way ANOVA with treatment as the explanatory factor. These tests were also significant at *P* < 0.05 for each compartment except  $G_2$ - $G_M$ . It was concluded that there was evidence for a treatment effect in each of the three cell cycle compartments  $G_0$ - $G_1$ , *S*, and  $G_2$ - $G_M$ . We then tested the ratios for each treatment within a compartment to determine if they significantly differed from 1.0, indicating a treatment effect for that particular treatment. Finally, we tested for a significant interaction of cannabidiol and  $\Delta^9$ -THC, each at its lowest dose by ANOVA with an interaction term. This was carried out after transforming the standardized ratios to logarithms so a test for additive interactions could be done.


## Results

### $\Delta^9$ -THC and Cannabidiol Inhibit the Growth of Multiple Glioblastoma Cell Lines

The  $CB_1$  and  $CB_2$  receptor agonist  $\Delta^9$ -THC can inhibit glioblastoma cell proliferation in culture and *in vivo*, and is currently being used in a clinical trial (4). Cannabidiol, a cannabinoid constituent with negligible affinity for  $CB_1$  and  $CB_2$  receptors, can also inhibit the proliferation of glioblastoma in culture and *in vivo* (7, 16). SF126, U251, and U87 cells were treated for three days with a range of concentrations of either  $\Delta^9$ -THC or cannabidiol. The antiproliferative activity of the compounds was assessed using the MTT assay and the corresponding  $IC_{50}$  values were calculated as previously described (26). The  $IC_{50}$  values for  $\Delta^9$ -THC in SF126, U251, and U87 cells were 2.5  $\mu$ mol (1.8–3.4), 3.3  $\mu$ mol/L (2.4–4.6), and 3.3  $\mu$ mol/L (2.3–4.8), respectively. The  $IC_{50}$  values for cannabidiol in SF126, U251, and U87 cells were 1.2  $\mu$ mol/L (1.1–1.3), 0.6  $\mu$ mol/L (0.5–1.0), and 0.6  $\mu$ mol/L (0.5–0.7), respectively. Cannabidiol was therefore a more potent inhibitor of cell growth than  $\Delta^9$ -THC in the three cell lines studied.

### Cannabidiol Enhances the Inhibitory Effects of $\Delta^9$ -THC on Glioblastoma Cell Growth

It has been suggested that nonpsychoactive cannabinoid constituents can either potentiate or inhibit the actions of  $\Delta^9$ -THC (11, 17–19). Therefore, the glioblastoma cell lines that were originally used to test the antiproliferative activity of individual cannabinoids were used to determine the effects of combination treatments. The positive and negative aspects of constituent interaction were tested by analyzing the activity of different combinations of  $\Delta^9$ -THC and cannabidiol in a 2 × 2 design (Fig. 1). The concentrations used for the treatments were  $IC_{80}$  or  $IC_{20}$  values calculated from the  $IC_{50}$  values as described in Materials and Methods. When applied in combination at the predicted  $IC_{80}$  concentration,  $\Delta^9$ -THC and cannabidiol produced a greater than additive inhibition of cell growth in SF126 and U251 cells. This was not observed in U87 cells



**Figure 1.** Cannabidiol (CBD) enhances the inhibitory effects of  $\Delta^9$ -THC on glioblastoma cell growth. To test for positive and negative interactions, a  $2 \times 2$  factorial design using specific  $\mu\text{mol/L}$  concentrations of drug was used as described in Materials and Methods. Cell proliferation was measured using the MTT assay. SF126 (A) and U251 (B) cells were treated for 3 d with vehicle/no drug,  $\Delta^9$ -THC, cannabidiol, or a combination of  $\Delta^9$ -THC and cannabidiol. Concentrations of  $\Delta^9$ -THC and cannabidiol that produce only minimal effects on cell proliferation were also tested in  $2 \times 2$  factorial design in SF126 (C) and U251 (D) cells. Percent control was calculated as the MTT product absorbance in the treated cells/control cells  $\times 100$ . Data are the mean of at least three independent experiments; bars,  $\pm$  SE. Data were analyzed using two-way ANOVA (GraphPad Prism). \*, statistically significant interaction ( $P < 0.01$ ). D, inset, representative light microscope image of the effects of the combination treatment on U251 cells ( $\times 40$ ).

(data not shown). In SF126 cells, cell viability was  $26 \pm 9\%$  in the presence of  $\Delta^9$ -THC ( $3.9 \mu\text{mol/L}$ ),  $40 \pm 4\%$  in cannabidiol ( $1.4 \mu\text{mol/L}$ ), and  $8 \pm 4\%$  in  $\Delta^9$ -THC ( $3.9 \mu\text{mol/L}$ ) plus cannabidiol ( $1.4 \mu\text{mol/L}$ ; Fig. 1A). In U251 cells, cell viability was  $55 \pm 3\%$  in the presence of  $\Delta^9$ -THC ( $5.4 \mu\text{mol/L}$ ),  $69 \pm 4\%$  in cannabidiol ( $0.9 \mu\text{mol/L}$ ), and  $2 \pm 2\%$  in  $\Delta^9$ -THC ( $5.4 \mu\text{mol/L}$ ) plus cannabidiol ( $0.9 \mu\text{mol/L}$ ; Fig. 1B). Predicted  $\text{IC}_{20}$  concentrations of  $\Delta^9$ -THC and cannabidiol that alone produce only minimal effects on cell growth were combined and further tested in a  $2 \times 2$  factorial design in the positive responding cell lines (SF126 and U251). Again, greater than additive effects were observed in SF126 and U251 cells. In SF126 cells, cell viability was  $90 \pm 4\%$  in the presence of  $\Delta^9$ -THC ( $1.6 \mu\text{mol/L}$ ),  $63 \pm 4\%$  in cannabidiol ( $1.1 \mu\text{mol/L}$ ), and  $25 \pm 6\%$  in  $\Delta^9$ -THC ( $1.6 \mu\text{mol/L}$ ) plus cannabidiol ( $1.1 \mu\text{mol/L}$ ; Fig. 1C). In U251 cells, cell viability was  $71 \pm 4\%$  in the presence of  $\Delta^9$ -THC ( $1.7 \mu\text{mol/L}$ ),  $83 \pm 5\%$  in cannabidiol ( $0.4 \mu\text{mol/L}$ ), and  $7 \pm 4\%$  in  $\Delta^9$ -THC ( $1.7 \mu\text{mol/L}$ ) plus cannabidiol ( $0.4 \mu\text{mol/L}$ ; Fig. 1D). In the cell lines showing significant interactions (SF126 and U251 cells), we further tested for synergism using the CI described in Materials and Methods. As shown in Supplementary Table S1, a synergistic increase in the anti-

proliferative activity of the cannabinoids was observed in both U251 and SF126 cells. CI values of  $<1$ ,  $1$ , and  $>1$  indicate synergism, additivity, and antagonism, respectively (29). Synergistic activity was observed at all the concentration ranges tested in U251 cells; therefore, this cell line was used primarily in the remaining experiments.


#### Cannabidiol Does Not Enhance the Inhibitory Effects of $\Delta^9$ -THC on Glioblastoma Cell Invasiveness

In addition to uncontrolled cell growth, a hallmark of the aggressive phenotype of glioblastoma cells is their ability to migrate away from the primary tumor of origin and invade into neighboring central nervous system tissue (30). Thus, we sought to determine whether the addition of cannabidiol to  $\Delta^9$ -THC would improve the activity of the compound to inhibit migration and invasion through a reconstituted basement membrane in a Boyden chamber assay (Fig. 2). Both  $\Delta^9$ -THC and cannabidiol could significantly inhibit the invasiveness of U251 cells. The predicted  $\text{IC}_{50}$  values for  $\Delta^9$ -THC and cannabidiol to inhibit U251 cell invasiveness were  $85 \text{ nmol/L}$  (range,  $49$ – $150$ ) and  $126 \text{ nmol/L}$  (range,  $20$ – $796$ ), respectively. Concentrations of  $100 \text{ nmol/L}$   $\Delta^9$ -THC and cannabidiol

(their approximate  $IC_{50}$  values) were used to test for positive or negative interactions. These concentrations were chosen, as compared with the predicted  $IC_{80}$  values, to ensure significant increases in cell death would not be produced, because this could confound the results of the invasion assay, i.e., dead cells would not migrate and invade. Whereas both  $\Delta^9$ -THC and cannabidiol were able to inhibit U251 cell invasiveness, cannabidiol did not enhance the activity of  $\Delta^9$ -THC when the compounds were combined. In U251 cells, invasiveness was  $48 \pm 3\%$  in the presence of  $\Delta^9$ -THC (0.1  $\mu\text{mol/L}$ ),  $72 \pm 3\%$  in cannabidiol (0.1  $\mu\text{mol/L}$ ), and  $36 \pm 3\%$  in  $\Delta^9$ -THC (0.1  $\mu\text{mol/L}$ ) plus cannabidiol (0.1  $\mu\text{mol/L}$ ). Because  $\Delta^9$ -THC and cannabidiol acted synergistically to inhibit glioblastoma cell growth, but not to inhibit cell invasiveness, mechanistic experiments were focused on understanding the reduction in cell viability produced by the combination treatment. The 4:1 ratio of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ; as described above in Fig. 1D) was primarily used as the combination treatment for the remaining experiments.

### The Combination Treatment of $\Delta^9$ -THC and Cannabidiol Leads to the Modulation of Specific Mitogen-Activated Protein Kinases

The regulation of extracellular signal-regulated kinase (ERK), *c-jun*-N-terminal kinase (JNK), and p38 mitogen-activated protein kinase (MAPK) activity plays a critical




**Figure 2.**  $\Delta^9$ -THC in combination with cannabidiol does not produce a greater overall inhibition of glioma invasiveness. To test for positive and negative interactions, a  $2 \times 2$  factorial design was used as described in Materials and Methods. The Boyden chamber invasion assay was used to determine the effects of treatment on the invasiveness of U251 cells. U251 cells were treated for 3 d with  $\Delta^9$ -THC (0.1  $\mu\text{mol/L}$ ), cannabidiol (0.1  $\mu\text{mol/L}$ ), or a combination of  $\Delta^9$ -THC (0.1  $\mu\text{mol/L}$ ) and cannabidiol (0.1  $\mu\text{mol/L}$ ). Data are presented as relative invasiveness of the cells through the Matrigel, where the respective controls are set as 100%. Data are the mean of at least three independent experiments; bars,  $\pm$  SE. \*, statistically significant differences from control ( $P < 0.05$ ).

role in controlling cell growth and apoptosis (31). Modulation of these pathways has been indicated in cannabinoid control of cancer cell growth and survival (9, 32–34). We used U251 cells to determine whether modulation of ERK, JNK, and p38 MAPK activity occurred. Treatment with the combination of cannabinoids led to a substantial downregulation of phosphorylated ERK (pERK), but produced no significant change in total ERK (Fig. 3A). Additionally, no inhibition of p38 MAPK or JNK activity was observed. When U251 cells were treated with individual concentrations of  $\Delta^9$ -THC and cannabidiol, instead of the combination, no changes in pERK were observed (Fig. 3B). These data show that the modulation of pERK was specific for the combination treatment. Downregulation of pERK in the presence of the combination treatment was first observed after two days treatment in U251 cells (Fig. 3C). The downregulation of pERK was also observed in SF126 cells using a combination treatment of  $\Delta^9$ -THC and cannabidiol (Fig. 3D). Therefore, the modulation of pERK by the combination treatment of THC and cannabidiol seems to represent a common mechanism shared by different glioblastoma cell lines.

### The Combination Treatment of $\Delta^9$ -THC and Cannabidiol Inhibits Cell Cycle and Induces Apoptosis

Significant reductions in ERK activity have been shown to lead to growth arrest and induction of apoptosis (31). The large reduction in glioblastoma cell viability and ERK activity, observed in the presence of the combination treatment, led us to hypothesize there would be a corresponding modulation of the cell cycle and programmed cell death. Therefore, U251 cells were treated with  $\Delta^9$ -THC and cannabidiol alone or with the combination of the two drugs, and cell cycle was analyzed using cell flow cytometry (Table 1). When administered separately,  $\Delta^9$ -THC and cannabidiol both produced increases in the population of cells in  $G_0$ - $G_1$  phase, but not in the S and  $G_2$ - $G_M$  phases. The combination of  $\Delta^9$ -THC and cannabidiol produced a greater than additive increase in the population of cells in  $G_0$ - $G_1$  phase and  $G_2$ - $G_M$  phase and a decrease in cells in S phase.

In addition to producing cell cycle arrest, the combination treatment may reduce cell viability through induction of apoptosis. We therefore measured apoptosis using annexin staining in combination with cell flow cytometry (Fig. 4A). There was a minor increase in apoptosis produced with 1.7  $\mu\text{mol/L}$   $\Delta^9$ -THC, but it was not found to be significantly different from control ( $n = 7$ ). No increase in apoptosis was observed in the presence of 0.4  $\mu\text{mol/L}$  cannabidiol. However, when  $\Delta^9$ -THC and cannabidiol were combined, a greater than additive increase in apoptosis was observed. In a time course analysis studying the induction of apoptosis produced by the combination treatment, we observed only a small increase with the combination treatment after two days (Supplementary Fig. S1A and B), whereas a strong induction of apoptosis was observed by day 3 (Fig. 4A).



**Figure 3.** The combination treatment of  $\Delta^9$ -THC and cannabidiol specifically inhibits ERK activity. The effects of cannabinoids on mitogen-activated protein kinases were analyzed using Western analysis. **A**, U251 cells were treated with vehicle or a combination of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ) for 3 d. Proteins were then extracted and analyzed for pERK, total ERK, pJNK1/2, and pP38 MAPK. **B**, U251 cells were treated with  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) or cannabidiol (0.4  $\mu\text{mol/L}$ ) alone for 3 d and analyzed for pERK and total ERK. **C**, U251 cells were treated with vehicle or a combination of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ) for 1 and 2 d. **D**, SF126 cells were treated with vehicle or a combination of  $\Delta^9$ -THC (1.6  $\mu\text{mol/L}$ ) and cannabidiol (1.1  $\mu\text{mol/L}$ ) for 12 h or 1 d. Either  $\alpha$ -tubulin or  $\beta$ -actin was used as a loading control (LC). Blots are representative of at least three independent experiments.

### The Inhibitory Effects of the Combination Treatment Are the Result of CB<sub>2</sub> Receptor Activation and Production of ROS

Depending on the glioblastoma cell line used, studies have linked the inhibitory activity of plant-derived cannabinoids to activation of CB<sub>1</sub> and/or CB<sub>2</sub>, modulation of MAPKs, and induction of cellular stress through increases in ROS and additional stress-related proteins, leading to activation of caspases (15, 32, 35). We used the measure of apoptosis to investigate mechanisms by which cannabidiol enhanced the activity of  $\Delta^9$ -THC.

Apoptosis produced by the combination of  $\Delta^9$ -THC and cannabidiol was partially blocked by the CB<sub>2</sub> receptor antagonist SR144528, but almost complete reversal was observed in the presence of the antioxidant  $\alpha$ -tocopherol (TCP; Fig. 4B). The cannabinoid receptor antagonists and TCP had no effect on apoptosis on their own at 0.5  $\mu\text{mol/L}$  and 20  $\mu\text{mol/L}$ , respectively (data not shown). As predicted by TCP blockade, the combination of  $\Delta^9$ -THC and cannabidiol produced a significant increase in the formation of ROS as assessed by DCDHF-DA

oxidation using FACS analysis (Supplementary Fig. S1C and D). A small increase in ROS was observed after one day of treatment, with a major induction observed by day 2, preceding the majority of the observed apoptotic cell death.

In an attempt to match levels of apoptosis produced by the combination treatment, the concentrations of the individual cannabinoids ( $\Delta^9$ -THC and cannabidiol) were next increased. The purpose of these experiments was to determine whether the compounds alone recruited similar pathways as compared with the combination of  $\Delta^9$ -THC and cannabidiol. When U251 cells were treated with  $\Delta^9$ -THC alone, the induction of apoptosis was almost completely blocked by TCP and partially blocked by the CB<sub>2</sub> antagonist SR144528 (Fig. 4C). However,  $\Delta^9$ -THC alone could not produce the level of apoptosis observed with the combination treatment (Fig. 4B and C). This finding was not simply an issue of the treatment concentration used because application of  $\Delta^9$ -THC up to 5  $\mu\text{mol/L}$  did not produce a greater induction of apoptosis (data not shown). When U251 cells were treated with cannabidiol alone, the induction of apoptosis was blocked by TCP but no reversal was observed with SR144528 (Fig. 4D). This result was expected because cannabidiol does not interact efficiently with either CB<sub>1</sub> or CB<sub>2</sub> receptors.

The ability of the higher concentrations of  $\Delta^9$ -THC (2.5  $\mu\text{mol/L}$ ) and cannabidiol (2.0  $\mu\text{mol/L}$ ) to inhibit pERK was also studied and compared with the combination treatment (Supplementary Fig. S2). Again, the combination treatment produced a substantial downregulation of pERK. However, the higher concentration of  $\Delta^9$ -THC alone had no effect on pERK. The higher concentration of cannabidiol produced only a small inhibition of pERK. Taken together, these data suggest that a unique pathway was activated by the combined administration of  $\Delta^9$ -THC and cannabidiol, which led to the downregulation of pERK.

### Cannabinoids Mediate Apoptosis through p8 and Caspases

The induction of the stress-associated gene, *p8*, has been shown to be a specific event in THC-induced apoptosis, but its involvement in cannabidiol-induced apoptosis has not been determined (35, 36). This pathway was evaluated to determine the role *p8* plays in the observed increase in apoptosis during cannabinoid treatments (Fig. 5A and B). Treatment of U251 cells with cannabidiol led to a small reduction in *p8* expression compared with control, but this change was not statistically significant ( $n = 4$ ). Treatment with  $\Delta^9$ -THC alone or the combination of  $\Delta^9$ -THC and cannabidiol led to an upregulation of *p8* expression. The magnitude of the effect was similar between treatment groups. These data show that the modulation of *p8* was not specific for the combination treatment.

Multiple caspase pathways were next evaluated to determine additional mechanisms by which the combination

**Table 1.** Cannabinoid modulation of cell cycle

Treatment, $\mu\text{mol/L}$	Mean ( $G_0$ - $G_1$ )	Mean (S)	Mean ( $G_2$ - $G_M$ )
CBD 0.4	1.08* $\pm$ 0.03	0.90 $\pm$ 0.05	0.97 $\pm$ 0.28
$\Delta^9$ -THC 1.7	1.12* $\pm$ 0.04	0.78 $\pm$ 0.11	1.28 $\pm$ 0.37
$\Delta^9$ -THC 1.7/CBD 0.4	1.23* <sup>†</sup> $\pm$ 0.02	0.49* <sup>†</sup> $\pm$ 0.08	2.69* <sup>†</sup> $\pm$ 0.56

NOTE: Cell cycle was measured using PI staining and FACS analysis, and Modfit was used to determine the percentage of cells in  $G_0$ - $G_1$ , S, and  $G_2$ - $G_M$  phase. U251 cells were treated for 3 d with CBD (0.4  $\mu\text{mol/L}$ ),  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ), or a combination of CBD (0.4  $\mu\text{mol/L}$ ) and  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ). The percentage of cells in each compartment was standardized by dividing it by the average percentage for the vehicle. This procedure was carried out for data from each experiment on each day. Statistical analysis was done as described in the Material and Methods.

Abbreviation: CBD, cannabidiol.


\* $P < 0.05$ .

<sup>†</sup>Significant interaction.


treatment increased apoptosis (Fig. 5C). In the presence of cannabidiol alone, no significant changes in caspase activity were observed. Small increases in the activity of caspase 7, caspase 9, and PARP, but not caspase 3, were observed when U251 cells were treated with  $\Delta^9$ -THC alone. Treatment with the combination of  $\Delta^9$ -THC and cannabidiol led to a substantial upregulation of caspase 3, 7, and 9 activities as well as an increase in PARP expression. These data show that a unique modulation of caspase activity is produced when glioblastoma cells are treated with the combination of  $\Delta^9$ -THC and cannabidiol as opposed to the individual cannabinoids.

## Discussion

We observed that plant-derived cannabinoids inhibit the proliferation of human glioblastoma cell lines. Compared with  $\Delta^9$ -THC, cannabidiol was significantly more potent at inhibiting cancer cell growth. This finding is in agreement with studies using models of aggressive breast cancers (26, 37).  $\Delta^9$ -THC is currently being used in a clinical trial for treatment of recurrent glioblastoma (5). Past studies have suggested that nonpsychoactive cannabinoids can modulate the actions of  $\Delta^9$ -THC (17–20). We hypothesized that the cannabinoid therapy utilizing



**Figure 4.** The effects of the combination treatment are the result of  $CB_2$  receptor activation. The number of U251 cells positive for annexin (apoptosis) staining after 3 d treatment was measured using FACS analysis. Cells were treated with **A**,  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ), cannabidiol (0.4  $\mu\text{mol/L}$ ), or a combination of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ) denoted as THC/CBD; **B**, a combination of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ) denoted as THC/CBD; **C**, 2.5  $\mu\text{mol/L}$   $\Delta^9$ -THC; and **D**, 2.0  $\mu\text{mol/L}$  cannabidiol. In **B**, **C**, and **D**, cells were also treated in the presence of 0.5  $\mu\text{mol/L}$  of the  $CB_1$  antagonist SR141716A (SR1), 0.5  $\mu\text{mol/L}$  of the  $CB_2$  antagonist SR144528 (SR2), or 20  $\mu\text{mol/L}$  TCP. Percent control was calculated as the positive annexin staining of the treated cells minus control cells. Data are the mean of at least three independent experiments; bars,  $\pm$  SE. Data were compared using one-way ANOVA with Bonferroni's multiple comparison posthoc analyses. \*, statistically significant differences from control ( $P < 0.05$ ); #, statistically significant differences from the combination treatment of THC/CBD ( $P < 0.05$ ).



**Figure 5.** When combined,  $\Delta^9$ -THC and cannabidiol produce an increase in activation of *p8* and multiple caspases. The effects of cannabinoids on *p8* and caspase expression were analyzed using semiquantitative reverse transcriptase-PCR and Western analysis, respectively. RNA and protein were collected from U251 cells treated for 3 d with cannabidiol (0.4  $\mu\text{mol/L}$ ),  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ), or a combination of  $\Delta^9$ -THC (1.7  $\mu\text{mol/L}$ ) and cannabidiol (0.4  $\mu\text{mol/L}$ ). **A**, reverse transcriptase-PCR was run on RNA extracted from control-treated and  $\Delta^9$ -THC/cannabidiol-treated samples. Expression of the  $\beta$ -actin gene product was used as a control for equal loading. **B**, data are represented as percentage *p8* expression of the treated cells/control cells  $\times 100$ , and all values were normalized against  $\beta$ -actin. Blots and PCR reactions are representative of at least three independent experiments. Data were compared using one-way ANOVA with Dunnett's multiple comparison posthoc analyses. \*, statistically significant differences from control ( $P < 0.05$ ). **C**, proteins were extracted from treated cells and analyzed for cleaved caspase 3, 7, 9, and PARP expression.

$\Delta^9$ -THC alone could be improved using a strategy of combination treatments.

We discovered that cannabidiol enhanced the ability of  $\Delta^9$ -THC to inhibit cell proliferation and induce cell cycle arrest and apoptosis. This activity occurred in two of three glioblastoma cell lines tested. Treatment of U251 cells with the combination led to a substantial downregulation of ERK activity, but not p38 MAPK and JNK1/2. The reduction in the phosphorylation of ERK was specific for the combination treatment and occurred in more than one glioblastoma cell line. Importantly, continuing to increase the concentration of  $\Delta^9$ -THC alone did not result in the inhibition of phosphorylated ERK. These data indicate that the enhanced effects observed were not solely due to an increase in the potency of  $\Delta^9$ -THC in U251 cells on coapplication with cannabidiol. Further support for this conclusion was observed when studying the activity of  $\Delta^9$ -THC and cannabidiol on U251 cell invasiveness. Both compounds were effective at inhibiting the invasiveness of U251 cells, but there was no evidence that cannabidiol improved the activity of  $\Delta^9$ -THC on coapplication.

In human glioblastoma cells, the ability of  $\Delta^9$ -THC to inhibit growth and induce apoptosis has been linked to the initial activation of CB<sub>1</sub> and CB<sub>2</sub> receptors (35). Similar effects produced by cannabidiol have been linked in part to CB<sub>2</sub> receptor activation, but the initial interaction site for the additional activity of cannabidiol remains to be clarified (7). We observed that increases in apoptosis produced by  $\Delta^9$ -THC alone, or the combination of  $\Delta^9$ -THC and cannabidiol, were partially dependent on CB<sub>2</sub> receptor activation. Apoptosis produced by cannabidiol alone was not dependent on CB<sub>2</sub> receptor activation. Importantly, the induction of apoptosis in the presence of the combination treatment was significantly greater than

that observed with  $\Delta^9$ -THC alone. Apoptosis produced by the combination of  $\Delta^9$ -THC and cannabidiol was dependent on the production of oxidative stress and resulted in a unique activation of both intrinsic and extrinsic caspases.

Studies have shown that the inhibitory activity of cannabinoids in glioblastoma is dependent on activation of CB<sub>1</sub> and CB<sub>2</sub> receptors, modulation of MAPKs, and induction of multiple types of cellular stresses leading to apoptosis (32, 35, 38, 39). In the case of  $\Delta^9$ -THC, upregulation of *p8* seems to be a specific event that leads to apoptosis in multiple types of cancers (35, 36). Treatment of U251 cells with the combination of  $\Delta^9$ -THC and cannabidiol led to an upregulation of *p8* expression, but similar activity was seen with  $\Delta^9$ -THC alone. This is in contrast to what we observed when studying modulation of caspase activity, and suggests that the enhanced apoptotic activity produced by the combination treatment was not the result of an interaction with the *p8* pathway.

The ability of cannabidiol to inhibit growth and induce apoptosis in glioblastoma and additional cancers has been primarily associated with the upregulation of ROS and multiple caspases, and has been linked to alterations in NADPH oxidases (9, 15). A link between ROS production and modulation of the LOX pathway has been hypothesized as a potential mechanism for the antitumor activity of cannabidiol in glioblastoma (16). In this study, an initial increase in ROS was clearly linked to a latter induction of apoptosis. Individually, both  $\Delta^9$ -THC and cannabidiol could increase apoptosis through the production of ROS, but  $\Delta^9$ -THC was significantly less efficient at inducing this process as a single agent as compared with when it was used in combination with cannabidiol. Although the concentration of cannabidiol used in the combination treatment did not significantly



stimulate ROS, it may have primed this pathway for  $\Delta^9$ -THC through a convergence on shared signal transduction pathways. A similar hypothesis could explain the unique downregulation of phosphorylated ERK that was produced by the combination treatment. Alternatively, cannabidiol may have potentiated the activity of  $\Delta^9$ -THC by inhibiting pathways that impart drug resistance in glioblastoma. For instance, a recent study showed that amphiregulin expression was associated with increased ERK activation, which mediated resistance to THC-induced apoptosis in gliomas (40). Therefore, cannabidiol may have potentiated the activity of THC-induced apoptosis by inhibiting amphiregulin-regulated increases in ERK activation. Future studies will be needed to elucidate the detailed mechanism associated with the unique effects of the  $\Delta^9$ -THC and cannabidiol combination treatment.

Individually,  $\Delta^9$ -THC and cannabidiol can activate distinct pathways in glioblastoma cells that ultimately culminate in inhibition of cancer cell growth and invasion as well as induction of cell death (2, 4, 41). We hypothesized that, if the individual agents were combined, a convergence on shared pathways may ensue, leading to an enhanced ability of the combination treatment to inhibit certain cancer cell phenotypes. We found this to be true in this investigation. Cannabidiol significantly improved the inhibitory effects of  $\Delta^9$ -THC on glioblastoma cell proliferation and survival, but not on cell invasiveness. The data suggest that the improved activity observed with the combination treatment is the result of a specific modulation of ERK and ROS activity leading to inhibition of cell cycle and induction of apoptosis.

Combinations, compared with individual drug treatments with specific cannabinoid-based compounds,

may represent an improvement for the treatment of patients with glioblastoma and perhaps additional cancers. It is also possible that other constituents of *C. sativa* that are not structurally related to cannabinoids could improve antitumor activity when combined. An important next step will be to carry out studies testing for synergistic antitumor activity of cannabinoids in additional preclinical models of glioblastoma. Even if synergism is not evident, combination treatments may allow for increased dosing due to nonoverlapping toxicities and decrease the development of resistance to the activity of  $\Delta^9$ -THC or cannabidiol when administered alone. With the growing evidence showing cannabinoids are effective inhibitors of multiple types of cancer, it is likely that additional clinical trials will be carried out. Combination treatments with cannabinoids may improve overall efficacy in these future clinical trials.

### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

### Grant Support

NIH (grants CA102412, CA111723, and DA09978), and the SETH Group.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 5/11/09; revised 11/12/09; accepted 11/15/09; published OnlineFirst 1/6/10.

### References

- Bifulco M, Di Marzo V. Targeting the endocannabinoid system in cancer therapy: a call for further research. *Nat Med* 2002;8:547–50.
- Blazquez C, Carracedo A, Salazar M, et al. Down-regulation of tissue inhibitor of metalloproteinases-1 in gliomas: a new marker of cannabinoid antitumoral activity? *Neuropharmacology* 2008;54:235–43.
- Preet A, Ganju RK, Groopman JE.  $\Delta(9)$ -Tetrahydrocannabinol inhibits epithelial growth factor-induced lung cancer cell migration *in vitro* as well as its growth and metastasis *in vivo*. *Oncogene* 2008;27:339–46.
- Velasco G, Carracedo A, Blazquez C, et al. Cannabinoids and gliomas. *Mol Neurobiol* 2007;36:60–7.
- Guzman M, Duarte MJ, Blazquez C, et al. A pilot clinical study of  $\Delta 9$ -tetrahydrocannabinol in patients with recurrent glioblastoma multiforme. *Br J Cancer* 2006;95:197–203.
- Pertwee RG. Pharmacology of cannabinoid CB1 and CB2 receptors. *Pharmacol Ther* 1997;74:129–80.
- Massi P, Vaccani A, Ceruti S, Colombo A, Abbraccio MP, Parolaro D. Antitumor effects of cannabidiol, a nonpsychoactive cannabinoid, on human glioma cell lines. *J Pharmacol Exp Ther* 2004;308:838–45.
- McAllister SD, Chan C, Taft RJ, et al. Cannabinoids selectively inhibit proliferation and induce death of cultured human glioblastoma multiforme cells. *J Neurooncol* 2005;74:31–40.
- McKallip RJ, Jia W, Schlomer J, Warren JW, Nagarkatti PS, Nagarkatti M. Cannabidiol-induced apoptosis in human leukemia cells: a novel role of cannabidiol in the regulation of p22phox and Nox4 expression. *Mol Pharmacol* 2006;70:897–908.
- Ruiz L, Miguel A, Diaz-Laviada I.  $\Delta 9$ -tetrahydrocannabinol induces apoptosis in human prostate PC-3 cells via a receptor-independent mechanism. *FEBS Lett* 1999;458:400–4.
- McPartland JM, Russo EB. Cannabis and cannabis extract: greater than the sum of the parts? *J Cannabis Therapeut* 2001;1:103–32.
- Howlett AC. Cannabinoid inhibition of adenylate cyclase: relative activity of the constituents and metabolites of marijuana. *Neuropharmacology* 1987;26:507–12.
- Devane WA, Dysarz I, Johnson FA, Melvin MR, Howlett LS. Determination and characterization of a cannabinoid receptor in rat brain. *Mol Pharmacol* 1988;34:605–13.
- Showalter VM, Compton DR, Martin BR, Abood ME. Evaluation of binding in a transfected cell line expressing a peripheral cannabinoid receptor (CB2): Identification of cannabinoid receptor subtype selective ligands. *J Pharmacol Exp Ther* 1996;278:989–99.
- Massi P, Vaccani A, Bianchessi S, Costa B, Macchi P, Parolaro D. The non-psychoactive cannabidiol triggers caspase activation and oxidative stress in human glioma cells. *Cell Mol Life Sci* 2006;63:2057–66.
- Massi P, Valenti M, Vaccani A, et al. 5-Lipoxygenase and anandamide hydrolase (FAAH) mediate the antitumor activity of cannabidiol, a non-psychoactive cannabinoid. *J Neurochem* 2008;104:1091–100.

17. Jones G, Pertwee RG. A metabolic interaction *in vivo* between cannabidiol and 1-tetrahydrocannabinol. *Br J Pharmacol* 1972;45:375–7.
18. Krantz JC, Berger HJ, Welch BL. Blockade of (-)-trans- $\delta$ -9-tetrahydrocannabinol depressant effect by cannabidiol in mice. *Amer J Pharm* 1971;143:149–52.
19. Poddar MK, Bhattacharya KC, Ghosh JJ. Potentiating effects of cannabidiol on  $\delta$ -9-tetrahydrocannabinol-induced changes in hepatic enzymes. *Biochem Pharmacol* 1974;23:758–9.
20. Russo E, Guy GW. A tale of two cannabinoids: the therapeutic rationale for combining tetrahydrocannabinol and cannabidiol. *Med Hypotheses* 2006;66:234–46.
21. Ben-Shabat S, Fride E, Sheskin T, et al. An entourage effect: inactive endogenous fatty acid glycerol esters enhance 2-arachidonoyl-glycerol cannabinoid activity. *Eur J Pharmacol* 1998;353:23–31.
22. Nurmikko TJ, Serpell MG, Hoggart B, Toomey PJ, Morlion BJ, Haines D. Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical trial. *Pain* 2007;133:210–20.
23. Rog DJ, Nurmikko TJ, Young CA. Oromucosal  $\delta$ 9-tetrahydrocannabinol/cannabidiol for neuropathic pain associated with multiple sclerosis: an uncontrolled, open-label, 2-year extension trial. *Clin Ther* 2007;29:2068–79.
24. Bornheim LM, Grillo MP. Characterization of cytochrome P450 3A inactivation by cannabidiol: possible involvement of cannabidiol-hydroxyquinone as a P450 inactivator. *Chem Res Toxicol* 1998;11:1209–16.
25. Bornheim LM, Kim KY, Li J, Perotti BY, Benet LZ. Effect of cannabidiol pretreatment on the kinetics of tetrahydrocannabinol metabolites in mouse brain. *Drug Metab Dispos* 1995;23:825–31.
26. McAllister SD, Christian RT, Horowitz MP, Garcia A, Desprez PY. Cannabidiol as a novel inhibitor of Id-1 gene expression in aggressive breast cancer cells. *Mol Cancer Ther* 2007;6:2921–7.
27. Slinker BK. The statistics of synergism. *J Mol Cell Cardiol* 1998;30:723–31.
28. Chou TC. Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. *Pharmacol Rev* 2006;58:621–81.
29. Chou TC, Tan QH, Sirotnak FM. Quantitation of the synergistic interaction of edatrexate and cisplatin *in vitro*. *Cancer Chemother Pharmacol* 1993;31:259–64.
30. Visted T, Enger PO, Lund-Johansen M, Bjerkvig R. Mechanisms of tumor cell invasion and angiogenesis in the central nervous system. *Front Biosci* 2003;8:e289–304.
31. Chang F, Steelman LS, Lee JT, et al. Signal transduction mediated by the Ras/Raf/MEK/ERK pathway from cytokine receptors to transcription factors: potential targeting for therapeutic intervention. *Leukemia* 2003;17:1263–93.
32. Galve-Roperh I, Sanchez C, Cortes ML, del Pulgar TG, Izquierdo M, Guzman M. Anti-tumoral action of cannabinoids: involvement of sustained ceramide accumulation and extracellular signal-regulated kinase activation. *Nat Med* 2000;6:313–9.
33. Gustafsson K, Christensson B, Sander B, Flygare J. Cannabinoid receptor-mediated apoptosis induced by R(+)-methanandamide and Win55,212–2 is associated with ceramide accumulation and p38 activation in mantle cell lymphoma. *Mol Pharmacol* 2006;70:1612–20.
34. Melck D, Rueda D, Galve-Roperh I, De Petrocellis L, Guzman M, Di Marzo V. Involvement of the cAMP/protein kinase A pathway and of mitogen-activated protein kinase in the anti-proliferative effects of anandamide in human breast cancer cells. *FEBS Lett* 1999;463:235–40.
35. Carracedo A, Lorente M, Egia A, et al. The stress-regulated protein p8 mediates cannabinoid-induced apoptosis of tumor cells. *Cancer Cell* 2006;9:301–12.
36. Carracedo A, Gironella M, Lorente M, et al. Cannabinoids induce apoptosis of pancreatic tumor cells via endoplasmic reticulum stress-related genes. *Cancer Res* 2006;66:6748–55.
37. Ligresti A, Moriello AS, Starowicz K, et al. Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on human breast carcinoma. *J Pharmacol Exp Ther* 2006;318:1375–87.
38. Jacobsson SO, Wallin T, Fowler CJ. Inhibition of rat C6 glioma cell proliferation by endogenous and synthetic cannabinoids. Relative involvement of cannabinoid and vanilloid receptors. *J Pharmacol Exp Ther* 2001;299:951–9.
39. Sanchez C, Galve-Roperh I, Canova C, Brachet P, Guzman M.  $\Delta$ 9-tetrahydrocannabinol induces apoptosis in C6 glioma cells. *FEBS Lett* 1998;436:6–10.
40. Lorente M, Carracedo A, Torres S, et al. Amphiregulin is a factor for resistance of glioma cells to cannabinoid-induced apoptosis. *Glia* 2009;57:1374–85.
41. Parolaro D, Massi P. Cannabinoids as potential new therapy for the treatment of gliomas. *Expert Rev Neurother* 2008;8:37–49.

# Molecular Cancer Therapeutics

## Cannabidiol Enhances the Inhibitory Effects of $\Delta^9$ -Tetrahydrocannabinol on Human Glioblastoma Cell Proliferation and Survival

Jahan P. Marcu, Rigel T. Christian, Darryl Lau, et al.

*Mol Cancer Ther* 2010;9:180-189. Published OnlineFirst January 12, 2010.

**Updated version** Access the most recent version of this article at:  
doi:[10.1158/1535-7163.MCT-09-0407](https://doi.org/10.1158/1535-7163.MCT-09-0407)

**Supplementary Material** Access the most recent supplemental material at:  
<http://mct.aacrjournals.org/content/suppl/2010/01/06/1535-7163.MCT-09-0407.DC1>

**Cited articles** This article cites 41 articles, 11 of which you can access for free at:  
<http://mct.aacrjournals.org/content/9/1/180.full#ref-list-1>

**Citing articles** This article has been cited by 8 HighWire-hosted articles. Access the articles at:  
<http://mct.aacrjournals.org/content/9/1/180.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://mct.aacrjournals.org/content/9/1/180>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.