

Curcumin Inhibition of the Functional Interaction between Integrin $\alpha6\beta4$ and the Epidermal Growth Factor Receptor

Young Hwa Soung and Jun Chung

Abstract

The functional interaction between integrin $\alpha6\beta4$ and growth factor receptors has been implicated in key signaling pathways important for cancer cell function. However, few attempts have been made to selectively target this interaction for therapeutic intervention. Previous studies showed that curcumin, a yellow pigment isolated from turmeric, inhibits integrin $\alpha6\beta4$ signaling important for breast carcinoma cell motility and invasion, but the mechanism is not currently known. To address this issue, we tested the hypothesis that curcumin inhibits the functional interaction between $\alpha6\beta4$ and the epidermal growth factor receptor (EGFR). In this study, we found that curcumin disrupts functional and physical interactions between $\alpha6\beta4$ and EGFR, and blocks $\alpha6\beta4$ /EGFR-dependent functions of carcinoma cells expressing the signaling competent form of $\alpha6\beta4$. We further showed that curcumin inhibits EGF-dependent mobilization of $\alpha6\beta4$ from hemidesmosomes to the leading edges of migrating cells such as lamellipodia and filopodia, and thereby prevents $\alpha6\beta4$ distribution to lipid rafts where functional interactions between $\alpha6\beta4$ and EGFR occur. These data suggest a novel paradigm in which curcumin inhibits $\alpha6\beta4$ signaling and functions by altering intracellular localization of $\alpha6\beta4$, thus preventing its association with signaling receptors such as EGFR. *Mol Cancer Ther*; 10(5); 883–91. ©2011 AACR.

Introduction

Curcumin is a polyphenolic component of turmeric (Fig. 1A), which is widely used as a spice and as a traditional ancient medicine in Indian culture (1). Curcumin has garnered a lot of attention as anticancer agent because of its nontoxic chemopreventive effect against virtually all known human cancers (2–3). There is accumulating evidence that curcumin inhibits cancer initiation and progression through regulation of multiple cellular pathways including EGFR/PDGFR (platelet-derived growth factor receptor; refs. 4, 5), AKT/mTOR (6, 7), NF- κ B (8), MAPK (9), and STAT pathways (10). Most of these curcumin targets are either signaling partners or downstream effectors of integrin $\alpha6\beta4$ (11–13), and our recent study showed that curcumin indeed inhibits $\alpha6\beta4$ signaling and functions associated with cancer cell motility and invasion (14). However, the mechanism by which curcumin inhibits $\alpha6\beta4$ signaling in aggressive cancer cells remains to be elucidated.

$\alpha6\beta4$ integrin is a laminin receptor and it was thought that its primary role was to maintain the integrity of epithelia (15). However, recent evidence has established that $\alpha6\beta4$ also plays a pivotal role in functions associated with tumorigenesis and carcinoma progression, suggesting that $\alpha6\beta4$ may switch its functions depending on the surrounding microenvironment (15, 16). In aggressive carcinoma cells, the host-tumor microenvironment induces growth factor-dependent mobilization of $\alpha6\beta4$ from hemidesmosomes (HD) into the leading edges of migrating cells such as lamellipodia and filopodia (17). The localization of $\alpha6\beta4$ to actin filament-rich structures such as lamellipodia and filopodia has an important implication for cancer cell motility and invasion, because integrins can harness traction forces through association with F-actin and their ability to engage the extracellular matrix (18, 19). In addition, localization of $\alpha6\beta4$ in the leading edge is thought to increase the level of $\alpha6\beta4$ in lipid rafts (20). Lipid rafts are sphingolipid and cholesterol-rich microdomains of the plasma membrane (21, 22). Lipid rafts can act as "signaling platforms," in which signaling initiation and amplification occur more efficiently by recruiting signaling receptors into close proximity (20–23). Once $\alpha6\beta4$ localizes in lipid rafts, it is assumed that its signaling function is enhanced through interaction with other signaling receptors such as EGFR (20, 21).

In this study, we assessed the mechanism by which curcumin inhibits $\alpha6\beta4$ signaling. Specifically, we tested the hypothesis that curcumin disrupts functional and physical interactions between $\alpha6\beta4$ and growth factor

Authors' Affiliation: Department of Biochemistry and Molecular Biology, Louisiana State University Health Science Center, Shreveport, Louisiana

Corresponding Author: Jun Chung, Department of Biochemistry and Molecular Biology, Louisiana State University Health Sciences Center, 1501 Kings Highway, Shreveport, LA 71103. Phone: 318-675-8797; Fax: 318-675-5180. E-mail: jchung@lsuhsc.edu

doi: 10.1158/1535-7163.MCT-10-1053

©2011 American Association for Cancer Research.

receptors. EGFR was chosen as a model to address this hypothesis because the interaction between EGFR and $\alpha 6\beta 4$ has been implicated in carcinoma cell functions important for progression (17, 24). We found that curcumin inhibits $\alpha 6\beta 4$ /EGFR-dependent proliferation in A431 squamous carcinoma and MDA-MB-231 breast carcinoma cells, both of which express signaling competent form of $\alpha 6\beta 4$. Additional studies have shown that curcumin reduces $\alpha 6\beta 4$ -dependent EGFR phosphorylation and the physical association between these 2 receptors. Curcumin inhibition of $\alpha 6\beta 4$ /EGFR signaling is likely because of its prevention of EGF-induced mobilization of $\alpha 6\beta 4$ from HDs to lamellipodia and filopodia, where $\alpha 6\beta 4$ can functionally interact with EGFR. Consistent with this observation, curcumin decreases the ratio of raft/nonraft $\alpha 6\beta 4$. Altogether, our studies provide a basis for a novel paradigm in which curcumin inhibits cancer cell functions by disrupting the interaction between integrin and growth factor receptors.

Materials and Methods

Cell lines and reagents

A431 squamous carcinoma cells, MDA-MB-231 breast carcinoma cells, and MCF10A cells were characterized by and purchased from American Type Culture Collections in January 2009. These cell lines have not been tested and authenticated by us. A431 and MDA-MB-231 cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) with 1 g/L glucose, L-glutamine, and sodium pyruvate formulation, supplemented with 10% FBS, 100 units/mL penicillin, and 100 g/mL streptomycin. MCF-10A cells were maintained in MEGM containing 13 mg/mL bovine pituitary extract, 0.5 mg hydrocortisone, 10 μ g/mL hEGF, 5 mg/mL insulin, and 100 ng/mL cholera toxin (Lonza). All cells used were always less than 20 passages from the stock and cultured in humidified incubators at 37°C in 5% CO₂. To stably reduce $\beta 4$ integrin expression, A431 cells were infected with lentiviruses that expressed short hairpin RNA (shRNA) targeted against either green fluorescent protein (GFP) or the $\beta 4$ integrin subunit. MDA-MB-231 cells expressing shRNA against either GFP or the $\beta 4$ integrin were used as previously described (14). To evaluate the effect of curcumin, cells were treated with the indicated doses (see figure legends) of curcumin (Sigma) under serum starvation for 24 hours, followed by stimulation with 10 ng/mL EGF for 15 to 20 minutes.

The following reagents were obtained commercially. EGF was purchased from Sigma-Aldrich (Sigma). Integrin $\beta 4$ (clone H-101) and actin (clone C-11) antibodies were purchased from Santa Cruz Biotechnology; Akt, p-Akt (Ser473), EGFR, p-EGFR(Tyr1068), and p-EGFR (Tyr1045) antibodies were obtained from Cell Signaling Technology. Phospho- $\beta 4$ integrin (Y1494) was purchased from ECM Biosciences. Flotillin-1 antibody was obtained from BD Transduction Laboratories and used to detect isolation of lipid rafts. Transferrin receptor antibody was

purchased from Invitrogen. Integrin $\beta 4$ (CD104) antibody was purchased from BD Pharmingen and used to visualize localization of $\beta 4$ and EGFR. EGFR (2E9) antibody was purchased from Santa Cruz Biotechnology. Alexa-fluor 546 goat anti-mouse immunoglobulin G (IgG) and Alexa-fluor 488 goat anti-rat IgG were purchased from Invitrogen.

Cell proliferation assay

The cells were plated in 6-well plates at 1×10^4 cells per well. Once stabilized, the cells were treated with the indicated doses of curcumin and then stimulated with or without 10 ng/mL of EGF. At the times indicated, cells were trypsinized and counted by a hemacytometer.

Soft agar growth assay

MDA-MB-231 cells (1×10^3) were suspended in a top layer of DMEM (1 mL) containing 0.35% low melt agarose (ISC BioExpress) with or without 10 μ mol/L curcumin and plated on a bottom layer of DMEM (2 mL) containing 0.75% agar with or without 10 ng/mL of EGF in 6-well plates. The cells were fed twice per week with 0.5 mL DMEM added with each indicated dose of EGF and curcumin. After 3 weeks, the total number of colonies was quantified by counting 50 fields per well by using bright-field optics. The average total number of colonies was obtained from counting triplicate wells.

Isolation of detergent-free lipid rafts

To isolate lipid raft fractions, we used the procedure described by Macdonald and Pike (25) with minor modifications. A431 cells were maintained in 150-nm dishes until the cells were 70% confluent and treated with or without curcumin overnight. The cells were scraped into cold base buffer [20 mmol/L Tris-HCl (pH 7.8), 250 mmol/L sucrose] to which 1 mmol/L CaCl₂ and 1 mmol/L MgCl₂ had been added. Cells pelleted by centrifugation were resuspended in 1 mL lysis buffer containing protease inhibitors and then lysed by repeated passage through a 22-gauge needle (20 times). After centrifugation at $1000 \times g$ for 10 minutes, the resulting postnuclear supernatants were collected into 12 mL centrifuge tubes. The pellets were lysed once more with 1 mL lysis buffer, and the second postnuclear supernatants were combined with the first. Equal volumes (2 mL) of base buffer containing 50% OptiPrep (Sigma) were mixed with the postnuclear supernatants in the bottoms of 12 mL centrifuge tubes. An 8 mL gradient of 0% to 20% OptiPrep in base buffer was poured on top and the tubes were centrifuged for 3 hours at $52,000 \times g$ by using the SW41 rotor in a Beckman ultracentrifuge. Gradients were fractionated into 1 mL fractions obtained from the top. The distribution of various proteins was determined by Western blotting.

Western blot analysis

Total cell extracts were prepared according to the methods described in our previous study (14), and separated on

4% to 20% gradient SDS-PAGE and transferred to polyvinylidene fluoride membranes (Millipore). The blots were incubated with primary antibodies in TBS-T or TBS-T with 5% (w/v) nonfat dry milk, then with appropriate secondary antibodies conjugated to IgG-horseradish peroxidase. Proteins were detected by using the pierce ECL Western blotting substrate (Thermo).

Immunoprecipitation assay

Cells were counted and plated into 10-cm or 6-well culture dishes. After reaching 50% to 60% of confluence, cells were treated with curcumin (15 $\mu\text{mol/L}$ in A431 cells and 10 $\mu\text{mol/L}$ in MDA-MB-231 cells) under serum starvation overnight and then stimulated with or without 10 ng/mL EGF for 20 minutes, before cell lysis. Cells were then lysed by using lysis buffer [50 mmol/L Tris-HCl (pH 7.4), 150 mmol/L NaCl, 1% NP-40, 5 mmol/L EDTA, NaF, Na_3VO_4 , $\text{Na}_4\text{P}_2\text{O}_7$, β -glycerolphosphate, aprotinin, leupeptin, pepstatin A, 100 mmol/L phenylmethylsulfonyl fluoride]. The lysates containing 1 mg of protein were precleared with protein A-conjugated agarose beads (Sigma) for 2 hours and then incubated with the appropriate primary antibodies overnight, followed by incubation with protein A-conjugated agarose beads. The beads were washed 4 times with PBS, and then boiled for 5 minutes in Laemmli sample buffer (BioRad Life Science). The immunoprecipitated proteins were separated by SDS-PAGE and probed with specific antibodies against EGFR, $\beta 4$, and IgG.

Immunofluorescence staining and confocal microscopy

A431 cells and MDA-MB-231 cells seeded on cover glasses at a density of 5×10^3 per well were treated with or without the indicated doses of curcumin overnight before EGF (100 ng/mL) stimulation for 15 minutes. After these treatments, cells were fixed in 4% paraformaldehyde for 15 minutes, permeabilized with 0.5% Triton X-100 for 15 minutes, and then incubated with primary antibody against EGFR and $\beta 4$ integrin overnight at 4°C. After washing 3 times with PBS, cells were incubated with Alexa Fluor 488 or Alexa Fluor 558-labeled secondary antibodies (Invitrogen). Fluorescent images were observed under a Zeiss LSM 510 laser-scanning confocal microscope. Colocalization of $\alpha 6\beta 4$ and EGFR was observed by merging the 2 fluorescence images in green (488nm) and red (546nm), indicating similar distribution patterns within merged images in yellow.

Results

Curcumin inhibits $\alpha 6\beta 4$ /EGFR-dependent functions of carcinoma cells that express signaling competent $\alpha 6\beta 4$

On the basis of the previous studies that curcumin inhibits carcinoma cell functions by inhibiting integrin

$\alpha 6\beta 4$ signaling (14), we tested the hypothesis that curcumin's inhibitory effect is mainly observed in cells that express the signaling competent form of $\alpha 6\beta 4$. To address this issue, we used A431 squamous carcinoma, MDA-MB-231 breast carcinoma, and MCF-10A normal epithelial cells. Both A431 and MDA-MB-231 carcinoma cells express the signaling competent form of $\alpha 6\beta 4$ (14, 17), as confirmed by high levels of $\beta 4$ phospho-Y1494 signal (Fig. 1B; an indicator of $\alpha 6\beta 4$ signaling competency; ref. 26). Consistent with our previous report (14), curcumin effectively blocked $\beta 4$ Y1494 phosphorylation in A431 cells and MDA-MB-231 cells in a dose-dependent manner (Fig. 1B). In contrast, MCF-10A cells express levels of $\beta 4$ integrin comparable to those of A431 and MDA-MB-231 cells, but the phospho-Y1494 signal was undetectable, suggesting that $\alpha 6\beta 4$ in MCF-10A cells is signaling incompetent (Fig. 1B).

On the basis of previous reports that $\alpha 6\beta 4$ synergizes with growth factor receptors to enhance carcinoma cell functions (16, 27–28), we assessed the effect of curcumin on $\alpha 6\beta 4$ -dependent cell proliferation in response to growth factor (EGF) stimulation in these 3 cell lines. These cells were treated with curcumin before stimulation with or without EGF. Proliferation of signaling competent $\alpha 6\beta 4$ expressing A431 cells and MDA-MB-231 cells was significantly enhanced by EGF treatment (Fig. 1C). EGF-dependent carcinoma cell proliferation requires the presence of $\alpha 6\beta 4$, as knockdown of $\beta 4$ integrin expression by shRNA effectively blocks EGF stimulation of carcinoma cell proliferation (data not shown). Curcumin also blocks EGF-dependent carcinoma cell proliferation in a dose-dependent manner, mimicking the loss of $\beta 4$ integrin expression (Fig. 1C). In contrast, signaling-incompetent $\alpha 6\beta 4$ expressing MCF-10A cells responded neither to EGF stimulation nor to curcumin treatment. These results suggest that the presence of the signaling competent form of $\alpha 6\beta 4$ sensitizes carcinoma cells to curcumin inhibition.

We then assessed the effect of curcumin treatment on $\alpha 6\beta 4$ /EGFR-dependent anchorage-independent growth, which is another important component of carcinoma progression, by measuring colony formation on soft agar (Fig. 2). As shown in Fig. 2, colony formation in soft agar requires the presence of $\alpha 6\beta 4$, as knockdown of $\beta 4$ expression by shRNA reduced colony formation by 60% compared with that of GFP shRNA expressing control cells. EGF stimulation in MDA-MB-231 cells further increased the colony formation and size in soft agar (Fig. 2). Curcumin effectively blocked colony formation in soft agar, with levels similar to those of MDA-MB-231 cells whose $\beta 4$ integrin is knocked down by shRNA in spite of EGF treatment (Fig. 2). No colony formation in soft agar was observed in MCF-10A cells, which express the signaling incompetent form of $\alpha 6\beta 4$ (data not shown). These results suggest that curcumin prevents anchorage-independent growth mediated by cooperative signaling between $\alpha 6\beta 4$ and EGFR.




Figure 1. Curcumin inhibits $\alpha 6\beta 4$ /EGFR-dependent cancer cell proliferation. **A**, chemical structure of curcumin. **B**, MDA-MB-231, A431, and MCF-10A cells were treated with the indicated doses of curcumin for 24 hours before lysis with radioimmunoprecipitation assay buffer. Equal amounts of extracts from each sample were used for Western blot analysis by using antibodies against $\beta 4$ integrin, phospho- $\beta 4$ integrin (Y1494), and β -actin. **C**, MDA-MB-231 cells, A431 cells, and MCF10A cells were serum starved, followed by treatment with different doses of curcumin (5 and 10 $\mu\text{mol/L}$; Cu5 and Cu10, respectively) and then stimulated with or without EGF (10 ng/mL). The cell numbers were counted by using a hemacytometer over the following 1-, 3- and 5-day periods. The statistical analysis was done by 2-tailed Student's *t* test. *, $P < 0.01$; **, $P < 0.05$ compared with the results of the control cell lines. Cont, control.

Curcumin blocks $\alpha 6\beta 4$ -dependent EGFR activation

Curcumin inhibition of $\alpha 6\beta 4$ /EGFR-mediated proliferation and anchorage-independent growth indicates its possible intervention in the cross-talk between these 2

receptors. To test this hypothesis, we examined whether curcumin blocks $\alpha 6\beta 4$ -dependent EGFR activation. As shown in Fig. 3A, the presence of $\alpha 6\beta 4$ in A431 cells and MDA-MB-231 cells contributes to higher steady-state




Figure 2. Curcumin inhibits $\alpha 6\beta 4$ /EGFR-mediated anchorage-independent growth in response to EGF. DMA-MB-231 cell lines (parental, $\beta 4$ shRNA infectants) were grown in 0.7% agar containing low serum media with or without curcumin (10 $\mu\text{mol/L}$) in the absence or presence of 10 ng/mL EGF for 3 weeks. Representative images of soft agar assays were captured at $\times 10$ magnification. Bottom, quantification of the soft agar assays. Columns, mean of 3 representative experiments done in triplicate; bars, SD. Significant differences were evaluated by Student's *t* test. *, $P < 0.01$; **, $P < 0.05$.

levels of EGFR Y1068 and Y1045 phosphorylation (indicator of EGFR activation; refs. 29, 30), as the knockout of $\alpha 6\beta 4$ expression by shRNA significantly reduced the phosphorylation of these residues. These data suggest that $\alpha 6\beta 4$ amplifies EGFR signaling through cross-talk even in the absence of ligand stimulation. EGF stimulation further increases the levels of EGFR Y1068 and Y1045 phosphorylation, but not in carcinoma cells deficient in $\alpha 6\beta 4$ expression by shRNA (data not shown). We next examined whether curcumin inhibits $\alpha 6\beta 4$ -dependent EGFR activation. As shown in Fig. 3B, curcumin effectively blocked phosphorylation of Y1068 and Y1045 of EGFR, and Y1494 of $\beta 4$ integrin, in MDA-MB-231 cells and A431 cells when these cells were treated with curcumin before EGF stimulation for 20 minutes. The inhibitory effect of curcumin on phosphorylation of these tyrosine residues was similar to that resulting from knockdown of $\beta 4$ integrin expression (Fig. 3). In addition, there is a nice correlation of phosphorylation levels between EGFR's Y1068/Y1045 and $\beta 4$ integrin's Y1494 (Fig. 3B), suggesting that these 2 receptors are function-

ally linked. These findings indicate that the inhibitory effect of curcumin on $\alpha 6\beta 4$ signaling could occur at least in part as a result of disruption of the functional interaction between these 2 receptors.

Curcumin blocks EGF-dependent mobilization of $\alpha 6\beta 4$ from hemidesmosomes into filopodia and lamellipodia

To gain additional evidence of curcumin disruption of the interaction between $\alpha 6\beta 4$ and EGFR, we assessed the effect of curcumin treatment on the intracellular localization of $\alpha 6\beta 4$ and EGFR in MDA-MB-231 cells and A431 cells by using confocal microscopy (Fig. 4). Under normal conditions (no EGF stimulation and no curcumin treatment; Fig. 4A and D), a significant portion of $\alpha 6\beta 4$ localized in the HDs (see arrow), whereas some portion of $\alpha 6\beta 4$ localized at the leading edges. As reported previously (17), EGF stimulation for 15 minutes destabilized the HDs and induced the mobilization of $\alpha 6\beta 4$ into the leading edge of MDA-MB-231 cells and A431 cells (Fig. 4B and E). Overall, there is a good correlation of localization between $\alpha 6\beta 4$ and EGFR. Their colocalization pattern is more obvious at the leading edges on EGF stimulation (Fig. 4B and E). In contrast, treatment of curcumin for 24 hours before EGF stimulation effectively blocks EGF-induced $\alpha 6\beta 4$ translocation from the HDs into the leading edges of MDA-MB-231 cells and A431 cells (Fig. 4C and F). Curcumin also blocks EGF-dependent disassembly of HDs as the majority of $\alpha 6\beta 4$ remains localized at HDs in spite of EGF treatment (Fig. 4C and F).

We then examined whether prevention of $\alpha 6\beta 4$ localization at the leading edges by curcumin has any impact on the physical association between $\alpha 6\beta 4$ and EGFR. We carried out coimmunoprecipitation assays with specific $\beta 4$ and EGFR antibodies. As shown in Fig. 5, curcumin treatment reduced the amount of coimmunoprecipitated $\alpha 6\beta 4$ with EGFR in MDA-MB-231 cells (Fig. 5A) and in A431 cells (Fig. 5B). Taken together, these results indicate that curcumin prevents the interaction between $\alpha 6\beta 4$ and EGFR, possibly by altering the intracellular location of $\alpha 6\beta 4$.

Curcumin prevents localization of $\alpha 6\beta 4$ /EGFR to lipid rafts

Mobilization of $\alpha 6\beta 4$ from HDs to leading edges of migrating carcinoma cells is thought to increase the localization of $\alpha 6\beta 4$ into lipid rafts (16), where lots of other signaling receptors are located in the near vicinity so that $\alpha 6\beta 4$ acts as a signal amplifier of these receptors (20,21,31). Therefore, we tested whether curcumin-mediated $\alpha 6\beta 4$ sequestration in HDs affects localization of $\alpha 6\beta 4$ in lipid rafts (Fig. 6). We monitored the distribution of $\alpha 6\beta 4$ and EGFR in the raft versus nonraft fractions of A431 carcinoma cells by using the detergent-free fractionation method as previously described (25). Fractions 1 and 2 represent the lipid raft fractions as confirmed by the lipid raft marker, flotillin-1, whereas fractions 3 or higher represent nonlipid raft fractions as indicated by




Figure 3. Curcumin disrupts $\alpha 6\beta 4$ integrin-dependent EGFR activation. **A**, equal extracts from the lysates of GFP and $\beta 4$ shRNA infectants of A431 and MDA-MB-231 cells were used for Western blot analysis by using antibodies against phospho- $\beta 4$ integrin (Y1494), phospho-EGFR (Y1068), phospho-EGFR (Y1045), or β -actin. **B**, A431 cells and MDA-MB-231 cells were pretreated with 15 and 10 $\mu\text{mol/L}$ curcumin, respectively, under serum starvation before EGF stimulation (10 ng/mL) for 20 minutes and then lysed by radioimmunoprecipitation assay buffer. Equal amounts of protein were isolated from extracts of these cells for Western blot analysis by using antibodies against phospho- $\beta 4$ integrin (Y1494), phospho-EGFR (Y1068), phospho-EGFR (Y1045), or β -actin. The Western blot results presented were carried out at least 3 times.

transferrin receptor signals (Fig. 6A). Densitometric analysis from Fig. 6A Western blot analysis showed that the steady-state raft/nonraft ratios of $\alpha 6\beta 4$ and EGFR are 1.17 and 1.32, respectively, in A431 cells, whereas curcumin treatment effectively lowered this ratio to 0.66 and 0.53, respectively (Fig. 6A). These results suggest that curcumin-mediated prevention of subcellular movement of $\alpha 6\beta 4$ in HDs negatively affects lipid raft localization of this integrin.

We then investigated the effect of curcumin on the physical association between $\alpha 6\beta 4$ and EGFR in raft versus nonraft fractions. Coimmunoprecipitation assays showed that physical association between these 2 recep-

tors is detected equally in both raft and nonraft fractions in unstimulated A431 cells, whereas EGF stimulation significantly increased their physical association, especially in raft fractions (Fig. 6B). Curcumin treatment effectively reduced the amount of overall coimmunoprecipitation between these 2 receptors by selectively affecting their association in raft fractions regardless of EGF stimulation (Fig. 6B). It is interesting to note that physical association between $\alpha 6\beta 4$ and EGFR in nonraft fractions is not affected by curcumin treatment (Fig. 6B). These data suggest that curcumin inhibition of the physical association between these 2 receptors is likely because of its prevention of their localization at the lipid rafts.




Figure 4. Curcumin blocks EGF-dependent mobilization of $\alpha 6\beta 4$ from HDs to the leading edge of cancer cells. MDA-MB-231 cells and A431 cells were plated on cover glasses and fixed with 0.5% Triton-X and double stained by using the anti- $\beta 4$ antibodies (green) and anti-EGFR antibodies (red) followed by Alexa-labeled secondary antibodies. A and D, control. B and E, stimulation with EGF in serum-free media for 15 minutes. C and F, pretreatment with curcumin (15 $\mu\text{mol/L}$ in A431 cells and 10 $\mu\text{mol/L}$ in MDA-MB-231 cells) for 24 hours before stimulation with 100 ng/mL EGF for 15 minutes. The localization of $\beta 4$ and EGFR was determined by the confocal microscopy. Arrows indicate HDs. Scale bar, 10 μm .

Discussion

It is widely assumed that $\alpha 6\beta 4$ plays a pivotal role in carcinoma progression through functional and physical interactions with specific growth factor receptors (16, 17). However, therapeutic intervention targeting these interactions has not been explored. On the basis of our previous report that curcumin inhibits $\alpha 6\beta 4$ -dependent breast cancer cell motility and invasion (14), we tested the hypothesis that curcumin inhibits $\alpha 6\beta 4$ signaling by intervening in the interaction between $\alpha 6\beta 4$ and growth factor receptors in this study. We used 2 carcinoma cell lines, MDA-MB-231 and A431, which overexpress both $\alpha 6\beta 4$ and EGFR (14, 17, 32). Interactions between these 2 receptors have been shown to be contributed to the malignant behaviors of these carcinoma cells (29, 33). Using these 2 carcinoma cell lines, we showed for the




Figure 5. Curcumin interrupts the physical association between $\beta 4$ and EGFR. MDA-MB-231 cells (A) and A431 cells (B) were treated with or without curcumin overnight before stimulation with 10 ng/mL EGF. The cell lysates were immunoprecipitated with EGFR, $\beta 4$, or control IgG antibodies. Western blotting was carried out on samples with integrin $\beta 4$, EGFR antibodies, and IgG antibodies. IP, immunoprecipitation.

first time that curcumin inhibits the functional and physical interactions between integrin and growth factor receptors. More specifically, curcumin effectively blocked EGFR/ $\alpha 6\beta 4$ -induced carcinoma cell proliferation and anchorage-independent growth. The mechanism involves prevention of $\alpha 6\beta 4$ mobilization from the HDs into the leading edges so that $\alpha 6\beta 4$ no longer effectively interacts with EGFR in the membrane. Additional studies have shown that curcumin reduced the amount of $\alpha 6\beta 4$ localized in lipid rafts where signaling molecules such as EGFR are concentrated. Altogether, our studies provide the evidence for a novel paradigm by which curcumin selectively inhibits carcinoma functions by altering subcellular localization of $\alpha 6\beta 4$ and therefore blocking its signaling competency.

The detailed mechanism(s) by which curcumin inhibits interaction between $\alpha 6\beta 4$ and EGFR remains to be determined. Our study suggests that either sequestration of $\alpha 6\beta 4$ in the HDs or prevention of its lipid raft localization by curcumin could be the major mechanisms. Earlier studies showed that, in response to EGF of cancer cells, $\alpha 6\beta 4$ in HDs mobilizes to lamellipodia and filopodia. This is where $\alpha 6\beta 4$ interacts with growth factor receptors, leading to interaction with the actin cytoskeleton important to cancer cell invasion and metastasis (17, 31). Phosphorylation of key serine residues (S1356, 1360, 1364, 1424) of $\beta 4$ integrin by PKC- α is involved in HD disassembly and the mobilization of $\alpha 6\beta 4$ from HDs (34–36). Therefore, it is possible that curcumin may affect one of the signaling pathways responsible for the phosphorylation of these Ser residues. Alternatively, palmitoylation of membrane




Figure 6. Curcumin prevents $\alpha 6\beta 4$ distribution to lipid rafts. **A**, A431 cells were treated with or without curcumin (15 $\mu\text{mol/L}$). Isolation of lipid rafts from the cells was done by the detergent-free method as described in Materials and Methods. Twelve fractions were collected by an Optiprep gradient producer. Each fraction was analyzed by Western blotting by using the indicated antibodies. Flotillin-1 (Flot) was used as a lipid raft marker and transferrin receptors (Trf) was used as a nonlipid raft marker. The Western blotting presented here was carried out at least 3 times. **B**, A431 cells were treated with or without 15 $\mu\text{mol/L}$ curcumin before stimulation with or without 10 ng/mL EGF for 20 minutes. The lipid raft fractions (1–3) and nonlipid raft fractions (4–6) from these cells were immunoprecipitated with $\beta 4$ antibodies and probed with EGFR antibodies. Densitometric analysis was done to measure the relative intensity of the bands from Western blotting analysis. Cur, curcumin; Non, nonraft.

proximal Cys residues of $\beta 4$ integrin is required for lipid raft localization of $\alpha 6\beta 4$ (20). On the basis of our data, which show that curcumin shifts $\alpha 6\beta 4$ from lipid raft to nonlipid raft fractions (Fig. 6), it is also possible that curcumin may block palmitoylation of the $\beta 4$ integrin subunit. These possibilities are currently under investigation.

The selective inhibitory effects of curcumin, as well as those of other phytochemicals, on cancer cell functions without toxicity to normal cells are well known, but there has been little explanation of the mechanisms. Our data,

which show that carcinoma cells expressing the signaling competent form of $\alpha 6\beta 4$ are more sensitive to curcumin, may provide the clue that explains the selective inhibitory effects of curcumin. This could be due to the fact that only the signaling competent form of $\alpha 6\beta 4$ localizes into lipid rafts where the functional interaction with growth factor receptors such as EGFR occurs. The selective inhibitory effect of curcumin on carcinoma cells could result from carcinoma cells depending more on lipid raft signaling as curcumin blocks trafficking of signaling molecules such as $\alpha 6\beta 4$ into the lipid rafts. In contrast, the signaling incompetent form of $\alpha 6\beta 4$ is mainly localized in HDs and simply provides tissue integrity in normal epithelia (37). Our studies show that curcumin does not affect HD stability or $\alpha 6\beta 4$ localization in HDs. Therefore, signaling-incompetent $\alpha 6\beta 4$ expressing cells would be less sensitive to curcumin treatment. Although this hypothesis needs to be tested on a larger scale in future studies, it is highly encouraging that this study provides the mechanistic basis for understanding the selective inhibitory effect of curcumin on cancer cells, but not normal cells.

The current paradigm for the most effective treatment of cancer is to combine multiple drugs against multiple targets. In this regard, our finding that curcumin has an inhibitory effect on the interaction of the 2 receptors suggests a novel possibility, that a multimodality approach such as one involving curcumin and EGFR inhibitors (i.e., gefitinib and eriotinib; ref. 38) could provide the design for effective multitarget agents for clinical chemotherapy. In conclusion, our study showed that the chemopreventive effect of curcumin in carcinoma cells derives from its ability to prevent $\alpha 6\beta 4$ mobilization into the leading edge and lipid rafts, which blocks cooperative interaction between $\alpha 6\beta 4$ and EGFR. On the basis of previous reports that functional interactions between $\alpha 6\beta 4$ and growth factor receptors are characteristic of aggressive carcinoma cells, our studies provide the basis for a novel paradigm that explains how curcumin selectively inhibits cancer cell function without affecting normal cells.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Dr. John Clifford for the critical reading of this manuscript.

Grant Support

This work is supported by the American Cancer Society, Life Savers Ball proceeds award, and Wendy Will Case Cancer Fund.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received November 18, 2010; revised February 8, 2011; accepted February 21, 2011; published OnlineFirst March 9, 2011.

References

1. Strimpakos AS, Sharma RA. Curcumin: preventive and therapeutic properties in laboratory studies and clinical trials. *Antioxid Redox Signal* 2008;10:511–45.
2. Inano H, Onoda M, Inafuku N, Kubota M, Kamada Y, Osawa T, et al. Chemoprevention by curcumin during the promotion stage of tumorigenesis of mammary gland in rats irradiated with gamma-rays. *Carcinogenesis* 1999;20:1011–8.
3. Dhillon N, Aggarwal BB, Newman RA, Wolff RA, Kunnumakkara AB, Abbruzzese JL, et al. Phase II trial of curcumin in patients with advanced pancreatic cancer. *Clin Cancer Res* 2008;14:4491–9.
4. Zhou Y, Zheng S, Lin J, Zhang QJ, Chen A. The interruption of the PDGF and EGF signaling pathways by curcumin stimulates gene expression of PPARgamma in rat activated hepatic stellate cell in vitro. *Lab Invest* 2007;87:488–98.
5. Chen A, Xu J, Johnson AC. Curcumin inhibits human colon cancer cell growth by suppressing gene expression of epidermal growth factor receptor through reducing the activity of the transcription factor Egr-1. *Oncogene* 2006;25:278–87.
6. Yu S, Shen G, Khor TO, Kim JH, Kong AN. Curcumin inhibits Akt/mammalian target of rapamycin signaling through protein phosphatase-dependent mechanism. *Mol Cancer Ther* 2008;7:2609–20.
7. Chaudhary LR, Hruska KA. Inhibition of cell survival signal protein kinase B/Akt by curcumin in human prostate cancer cells. *J Cell Biochem* 2003;89:1–5.
8. Aggarwal S, Takada Y, Singh S, Myers JN, Aggarwal BB. Inhibition of growth and survival of human head and neck squamous cell carcinoma cells by curcumin via modulation of nuclear factor-kappaB signaling. *Int J Cancer* 2004;111:679–92.
9. Squires MS, Hudson EA, Howells L, Sale S, Houghton CE, Jones JL, et al. Relevance of mitogen activated protein kinase (MAPK) and phosphatidylinositol-3-kinase/protein kinase B (PI3K/PKB) pathways to induction of apoptosis by curcumin in breast cells. *Biochem Pharmacol* 2003;65:361–76.
10. Bharti AC, Donato N, Aggarwal BB. Curcumin (diferuloylmethane) inhibits constitutive and IL-6-inducible STAT3 phosphorylation in human multiple myeloma cells. *J Immunol* 2003;171:3863–71.
11. Chung J, Bachelder RE, Lipscomb EA, Shaw LM, Mercurio AM. Integrin (alpha 6 beta 4) regulation of eIF-4E activity and VEGF translation: a survival mechanism for carcinoma cells. *J Cell Biol* 2002;158:165–74.
12. Zahir N, Lakins JN, Russell A, Ming W, Chatterjee C, Rozenberg GI, et al. Autocrine laminin-5 ligates alpha6beta4 integrin and activates RAC and NFkappaB to mediate anchorage-independent survival of mammary tumors. *J Cell Biol* 2003;163:1397–407.
13. Mercurio AM. Invasive skin carcinoma–Ras and alpha6beta4 integrin lead the way. *Cancer Cell* 2003;3:201–2.
14. Kim HI, Huang H, Cheepala S, Huang S, Chung J. Curcumin inhibition of integrin (alpha6beta4)-dependent breast cancer cell motility and invasion. *Cancer Prev Res* 2008;1:385–91.
15. Mercurio AM, Rabinovitz I, Shaw LM. The alpha 6 beta 4 integrin and epithelial cell migration. *Curr Opin Cell Biol* 2001;13:541–5.
16. Lipscomb EA, Mercurio AM. Mobilization and activation of a signaling competent alpha6beta4 integrin underlies its contribution to carcinoma progression. *Cancer Metastasis Rev* 2005;24:413–23.
17. Rabinovitz I, Toker A, Mercurio AM. Protein kinase C-dependent mobilization of the alpha6beta4 integrin from hemidesmosomes and its association with actin-rich cell protrusions drive the chemotactic migration of carcinoma cells. *J Cell Biol* 1999;146:1147–60.
18. Ridley AJ, Schwartz MA, Burridge K, Firtel RA, Ginsberg MH, Borisy G, et al. Cell migration: integrating signals from front to back. *Science* 2003;302:1704–9.
19. de Pereda JM, Lillo MP, Sonnenberg A. Structural basis of the interaction between integrin alpha6beta4 and plectin at the hemidesmosomes. *EMBO J* 2009;28:1180–90.
20. Gagnoux-Palacios L, Dans M, van't Hof W, Mariotti A, Pepe A, Meneguzzi G, et al. Compartmentalization of integrin alpha6beta4 signaling in lipid rafts. *J Cell Biol* 2003;162:1189–96.
21. De Laurentis A, Donovan L, Arcaro A. Lipid rafts and caveolae in signaling by growth factor receptors. *Open Biochem J* 2007;1:12–32.
22. Simons K, Toomre D. Lipid rafts and signal transduction. *Nat Rev Mol Cell Biol* 2000;1:31–9.
23. Patra SK. Dissecting lipid raft facilitated cell signaling pathways in cancer. *Biochim Biophys Acta* 2008;1785:182–206.
24. Jorissen RN, Walker F, Pouliot N, Garrett TP, Ward CW, Burgess AW. Epidermal growth factor receptor: mechanisms of activation and signaling. *Exp Cell Res* 2003;284:31–53.
25. Macdonald JL, Pike LJ. A simplified method for the preparation of detergent-free lipid rafts. *J Lipid Res* 2005;46:1061–7.
26. Dutta U, Shaw LM. A key tyrosine (Y1494) in the beta4 integrin regulates multiple signaling pathways important for tumor development and progression. *Cancer Res* 2008;68:8779–87.
27. Gambaletta D, Marchetti A, Benedetti L, Mercurio AM, Sacchi A, Falcioni R. Cooperative signaling between alpha(6)beta(4) integrin and ErbB-2 receptor is required to promote phosphatidylinositol 3-kinase-dependent invasion. *J Biol Chem* 2000 Apr 7;275:10604–10.
28. Mainiero F, Pepe A, Yeon M, Ren Y, Giancotti FG. The intracellular functions of alpha6beta4 integrin are regulated by EGF. *J Cell Biol* 1996;134:241–53.
29. Rojas M, Yao S, Lin YZ. Controlling epidermal growth factor (EGF)-stimulated Ras activation in intact cells by a cell-permeable peptide mimicking phosphorylated EGF receptor. *J Biol Chem* 1996;271:27456–61.
30. Frey MR, Dise RS, Edelblum KL, Polk DB. p38 kinase regulates epidermal growth factor receptor downregulation and cellular migration. *EMBO J* 2006;25:5683–92.
31. Mariotti A, Kedeshian PA, Dans M, Curatola AM, Gagnoux-Palacios L, Giancotti FG. EGF-R signaling through Fyn kinase disrupts the function of integrin alpha6beta4 at hemidesmosomes: role in epithelial cell migration and carcinoma invasion. *J Cell Biol* 2001;155:447–58.
32. Gilcrease MZ, Zhou X, Welch K. Adhesion-independent alpha6beta4 integrin clustering is mediated by phosphatidylinositol 3-kinase. *Cancer Res* 2004;64:7395–8.
33. Gilcrease MZ, Zhou X, Lu X, Woodward WA, Hall BE, Morrissey PJ. Alpha6beta4 integrin crosslinking induces EGFR clustering and promotes EGF-mediated Rho activation in breast cancer. *J Exp Clin Cancer Res* 2009;26:28:67.
34. Rabinovitz I, Tsomo L, Mercurio AM. Protein kinase C-alpha phosphorylation of specific serines in the connecting segment of the beta 4 integrin regulates the dynamics of type II hemidesmosomes. *Mol Cell Biol* 2004;24:4351–60.
35. Wilhelmsen K, Litjens SH, Kuikman I, Margadant C, van Rheeën J, Sonnenberg A. Serine phosphorylation of the integrin beta4 subunit is necessary for epidermal growth factor receptor induced hemidesmosome disruption. *Mol Biol Cell* 2007;18:3512–22.
36. Germain EC, Santos TM, Rabinovitz I. Phosphorylation of a novel site on the (beta)4 integrin at the trailing edge of migrating cells promotes hemidesmosome disassembly. *Mol Biol Cell* 2009;20:56–67.
37. Borradori L, Sonnenberg A. Structure and function of hemidesmosomes: more than simple adhesion complexes. *J Invest Dermatol* 1999;112:411–8.
38. Modjtahedi H, Essapen S. Epidermal growth factor receptor inhibitors in cancer treatment: advances, challenges and opportunities. *Anticancer Drugs* 2009;20:851–5.

Molecular Cancer Therapeutics

Curcumin Inhibition of the Functional Interaction between Integrin $\alpha6\beta4$ and the Epidermal Growth Factor Receptor

Young Hwa Soung and Jun Chung

Mol Cancer Ther 2011;10:883-891. Published OnlineFirst March 9, 2011.

Updated version Access the most recent version of this article at:
doi:[10.1158/1535-7163.MCT-10-1053](https://doi.org/10.1158/1535-7163.MCT-10-1053)

Cited articles This article cites 38 articles, 19 of which you can access for free at:
<http://mct.aacrjournals.org/content/10/5/883.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/10/5/883.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/10/5/883>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.