

Regulation of *Vinca* alkaloid-induced apoptosis by NF- κ B/I κ B pathway in human tumor cells

Yi Huang,¹ Yong Fang,^{1,2} Jinmin Wu,²
Jennifer M. Dziadyk,¹ Xueming Zhu,¹
Meihua Sui,¹ and Weimin Fan^{1,2}

¹ Department of Pathology and Laboratory Medicine, Medical University of South Carolina, Charleston, SC and ² Department of Oncology, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China

Abstract

Antimicrotubule *Vinca* alkaloids, such as vinblastine and vincristine, interfere with the dynamics of microtubules and have shown significant cell killing activity in a variety of tumor cells through induction of apoptosis. The mechanism by which *Vinca* alkaloids induce apoptosis is not entirely clear. In this study, we found that glucocorticoids inhibit *Vinca* alkaloid-induced apoptosis without affecting G₂-M arrest in human breast cancer BCap37 cells and human epidermoid tumor KB cells, suggesting that *Vinca* alkaloid-induced apoptosis may occur via a pathway independent of cell cycle arrest. Further analyses indicated that *Vinca* alkaloids cause significant degradation of I κ B α , which in turn results in nuclear factor- κ B (NF- κ B) activation. Transfection of antisense I κ B α in BCap37 cells sensitizes *Vinca* alkaloid-induced apoptosis. Moreover, *in vitro* kinase assays show that the activity of I κ B kinase (IKK) was activated by *Vinca* alkaloids and was not affected by glucocorticoids. Stable transfection of dominant-negative deletional mutant I κ B α , which is insensitive to IKK-mediated phosphorylation and degradation, resulted in the inhibition of *Vinca* alkaloid-induced NF- κ B activation and reduced sensitivity of tumor cells to *Vinca* alkaloid-induced apoptosis. These findings suggest that the NF- κ B/I κ B signaling pathway may contribute to the mediation of *Vinca* alkaloid-induced apoptosis in human tumor cells. [Mol Cancer Ther. 2004;3(3):271–277]

Introduction

The *Vinca* alkaloid antimicrotubule agents, including vinblastine and vincristine, have been widely used as clinical anticancer agents for the treatment of leukemia, lymphomas, and some solid tumors (1). Unlike other classes of antimicrotubule agents such as taxanes, *Vinca*

alkaloids induce the destabilization of polymerized tubulin by blocking the region involved in tubulin dimer attachment, thus preventing polymerization of microtubules (2). It has generally been believed that the antitumor effects of *Vinca* alkaloids mainly depend on interference with the normal function of microtubules and blockage of cell cycle progression in the G₂-M phase. In recent years, several laboratories demonstrated that, at clinically relevant concentrations, *Vinca* alkaloids are able to induce apoptotic cell death in several solid tumor cells (2, 3).

Previous studies have revealed that glucocorticoids could selectively inhibit paclitaxel-induced apoptotic cell death but did not affect the ability of paclitaxel to induce microtubule bundling and mitotic arrest (4, 5). This phenomenon suggested that antimicrotubule agent-induced apoptosis might take place via a separate pathway independent of cell cycle arrest. Further studies demonstrated that paclitaxel and glucocorticoids possess opposite regulatory effects on I κ B α degradation and activation of nuclear factor- κ B (NF- κ B; Ref. 6). NF- κ B, a member of Rel transcription factor family, and its specific intracellular inhibitor, I κ B α , participate in the mediation or regulation of many biological processes including inflammation, immune response, cell proliferation, and apoptotic cell death (7, 8). NF- κ B normally resides in the cytoplasm as an inactivated form by forming a complex with its inhibitory protein I κ B α . On certain stimuli, I κ B α is rapidly phosphorylated and degraded, allowing NF- κ B to translocate to the nucleus, where it participates in transcriptional regulation of numerous genes (8, 9). A key player in this cascade of events is the I κ B kinase complex (IKK α and IKK β), which is responsible for the phosphorylation and degradation of I κ B α (10). In recent years, increasing evidence indicates that activation of NF- κ B plays an important role in coordinating the control of apoptotic cell death, which either promotes or inhibits apoptosis, depending on different apoptotic stimulus and cell types (9, 11–15).

In this study, through characterization of the inhibitory effect of glucocorticoids on vinblastine- and vincristine-induced apoptosis in human breast cancer BCap37 cells and human epidermoid tumor KB cells, we obtained evidence that *Vinca* alkaloids cause the degradation of I κ B α , which in turn promotes nuclear translocation and activation of NF- κ B. We also found that glucocorticoids inhibit *Vinca* alkaloid-induced apoptosis via antagonizing the ability of *Vinca* alkaloids in the activation of NF- κ B. These findings suggest that the NF- κ B/I κ B α signaling pathway plays an important role in the mediation of *Vinca* alkaloid-induced tumor cell apoptosis.

Materials and Methods

Drugs and Cell Culture

Vinblastine and vincristine were purchased from Sigma Chemical Co. (St. Louis, MO) and dissolved in 100% DMSO

Received 8/12/03; revised 11/19/03; accepted 12/9/03.

Grant support: NIH grants CA 92880 and CA 82440 (W. Fan).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for Reprints: Weimin Fan, Department of Pathology and Laboratory Medicine, Medical University of South Carolina, 171 Ashley Avenue, Charleston, SC 29425. Phone: (843) 792-5108; Fax: (843) 792-0368. E-mail: fanw@muscc.edu

to make a stock solution with various concentrations. Triamcinolone acetonide (TA) was also purchased from Sigma Chemical and dissolved in 100% ethanol as 10^{-2} to 10^{-5} M stock solution. Wild-type breast tumor BCap37 cells, stable BCap37 cells transfected with antisense I κ B α (6) and mutant I κ B α (16), and human epidermoid tumor KB cells were cultured in RPMI 1640 (Mediatech, Herndon, VA) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin (Hyclone Laboratories, Logan, UT).

Preparation of Glutathione S-Transferase-I κ B α Fusion Protein

pGEX-I κ B α fusion protein expression vectors were constructed by subcloning I κ B α cDNA restriction enzyme fragments from pCR2.1-I κ B α vectors. Glutathione S-transferase (GST)-I κ B α fusion proteins were purified from *Escherichia coli* cells transformed with pGEX-I κ B α expression vectors by using glutathione-agarose affinity chromatography (Amersham Biosciences, Piscataway, NJ).

Determination of Internucleosomal DNA Fragmentation

After incubation with the various concentration of drug, $\sim 1 \times 10^6$ cells were harvested and suspended in lysis buffer [5 mM Tris-HCl (pH 8.0), 20 mM EDTA, and 0.5% (v/v) Triton X-100] for 20 min on ice. The remaining steps for DNA fragmentation were performed as described previously (6). DNA samples were analyzed by electrophoresis on a 1.2% agarose gel containing ethidium bromide (0.2 μ g/ml) and visualized under UV illumination.

Western Blotting

After cells were treated with the various concentrations of drugs, cellular proteins were isolated and fractionated on a 12.5% SDS-PAGE gel and transferred to polyvinylidene difluoride (PVDF) membranes using the methods as described previously (6). The membranes were incubated with anti-I κ B α or anti-IKK α primary antibodies (1:2000; Santa Cruz Biotechnology, Santa Cruz, CA). The membranes were then incubated with peroxidase-conjugated secondary antibody (1:5000; ImmunoResearch, West Grove, PA) followed by enhanced chemiluminescence staining system (Amersham Bioscience, Inc.). β -actin proteins were used to normalize protein loading.

Flow Cytometry Analysis

Cells were treated with various increasing concentration of drugs for 48 h and then harvested and fixed in 70% ethanol in PBS. After washing in rinse buffer (0.5% BSA, 0.1% Triton X-100 in PBS) twice, cells were incubated in PBS containing RNase (50 μ g/ml), EDTA (0.1 M), and propidium iodide (50 μ g/ml) at room temperature for 1 h. Cell cycle distribution was determined using a Coulter Epics V instrument (Coulter Corp., Fullerton, CA) with an argon laser set to excite at 488 nm. The results were analyzed using Elite 4.0 software (Phoenix Flow System, San Diego, CA).

Light Microscope Examination

The cytospin preparations and light microscope examination were performed as described previously (17). Briefly, cells were plated and treated with 100 nM

vinblastine or vincristine with or without TA treatment. After 48 h of culture, cells were harvested by trypsinization and washed with PBS twice. About $5\text{--}10 \times 10^4$ cells were used for cytospin preparation. Slides were air-dried and fixed in methanol for 5 min prior to Wright-Giemsa staining. The DNA contents of G₂-M phase of cells were counted under bright-field microscopy. Data presented represent three independent experiments.

MTT Assays

Growth inhibition was assessed by using MTT assays as described previously (16). Briefly, 2×10^5 cells/well were plated in 96-well dishes and treated with the various drug regimes for the indicated times. All of the experiments were plated in triplicate and the results of assays were presented as means \pm SD.

Immunoprecipitation and IKK Assays

Cells were washed twice in PBS buffer and resuspended in 500 μ l of immunoprecipitation lysis buffer [50 mM Tris-HCl (pH 7.6), 150 mM NaCl, 10% glycerol, 1% NP40, 5 mM EDTA, 1 mM DTT, 100 mM NaF, 2 mM sodium pyrophosphate, 2 mM sodium orthovanadate, 1 mM phenylmethylsulfonyl fluoride (PMSF), 10 μ g/ml of aprotinin, and leupeptin] and stored on ice for 20 min before centrifugation (14,000 $\times g$, 20 min, 4°C). IKK complex was immunoprecipitated by incubation for 2 h at 4°C with IKK α rabbit polyclonal antibodies (Santa Cruz Biotechnology) bound to protein A-Sepharose (Pharmacia/Biotech). The immunoprecipitates were washed twice with immunoprecipitation buffer and twice with kinase buffer [20 mM HEPES (pH 7.4), 20 mM β -glycerophosphate, 20 mM MgCl₂, 2 mM DTT, and 0.1 mM sodium orthovanadate]. The kinase assays were initiated by the addition of GST-I κ B α fusion protein (1 mg) and [γ -³²P] ATP (10 Ci/mmol). Reaction mixtures were incubated for 30 min at 30°C and stopped by the addition of 2 \times SDS-PAGE sample buffer. The phosphorylation of the I κ B α proteins was examined by SDS-PAGE followed by autoradiography.

Nuclear Extraction Preparation and Electrophoretic Mobility Shift Assays

After BCap37 cells were transfected with empty pcDNA3 vectors or mutant I κ B α , cells were treated with different concentrations of vinblastine or vincristine for 24 h, harvested, and resuspended in 1 ml of hypotonic lysis buffer [10 mM HEPES (pH 7.8), 10 mM KCl, 2 mM MgCl₂, 1 mM DTT, 0.1 mM EDTA, and 0.1 mM PMSF]. Cells were then incubated on ice for 15 min. Ten-percent NP40 (25 μ l) was then added and vigorously mixed and centrifuged. The nuclear pellets were suspended in 50 μ l of extraction buffer (50 mM KCl, 300 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 0.1 mM PMSF, and 10% glycerol), mixed for 20 min, and centrifuged at 12,000 $\times g$ for 20 min to produce supernatant containing nuclear proteins. Protein concentrations were determined using the Bio-Rad Dc Protein Assay (Bio-Rad, San Diego, CA). Electrophoretic mobility shift assays (EMSA) were performed using ³²P-labeled double-stranded oligonucleotide probes, which contain a specific consensus sequence (5'-AGTTGAGGGGAGTTTCCCAGGC-3';

Santa Cruz Biotechnology) recognized by NF- κ B. Probes were labeled with T4 polynucleotide kinase (Promega, Madison, WI) and [γ - 32 P]ATP and purified using G-25 spin columns (Amersham Bioscience). The remaining steps for EMSAs were performed as described previously (16).

Statistical Analysis

The Student's *t* test was used to determine the statistical differences between various experimental and control groups. *P* < 0.01 was considered significant.

Results

Glucocorticoids Inhibit *Vinca* Alkaloid-Induced Apoptosis in BCap37 and KB Cells

An important hallmark of apoptotic cell death is the fragmentation of genomic DNA into integer multiples of 180-bp units, producing a characteristic ladder on agarose gel electrophoresis. To examine the cytotoxicity of *Vinca* alkaloids to tumor cells as well as the possible effects of glucocorticoids on antitumor activity of *Vinca* alkaloids, human breast tumor BCap37 and epidermoid tumor KB cells were treated with a variety of concentrations of vinblastine and vincristine with or without pretreatment of glucocorticoids followed by internucleosomal DNA fragmentation assays. As shown in Fig. 1, DNA fragmentation ladders were clearly observed following treatment of BCap37 cells and KB cells with 20 nM or higher concentrations of vinblastine and vincristine. These results demonstrate that BCap37 and KB cells are sensitive to *Vinca* alkaloid-induced apoptosis. However, when the tumor cells were pretreated with glucocorticoids (10^{-7} M TA) for 24 h, *Vinca* alkaloid-induced DNA fragmentation was dramatically inhibited in both BCap37 and KB cells. This result suggests that glucocorticoids inhibit *Vinca* alkaloid-induced apoptosis in human tumor cell lines.




Figure 1. Glucocorticoids inhibit *Vinca* alkaloid-induced DNA fragmentation. BCap37 cells and KB cells were treated with different concentrations of vinblastine (*Vinb.*) or vincristine (*Vinc.*) for 48 h with or without pretreatment of 10^{-7} M TA (*Gluco.*). Cells were then harvested and fragmented DNA was extracted and analyzed by electrophoresis in 1.2% agarose gel containing 0.1% ethidium bromide.

Glucocorticoids Do Not Affect *Vinca* Alkaloid-Induced Mitotic Arrest

To determine the possible influence of glucocorticoids on *Vinca* alkaloid-induced mitotic arrest, a cytospin method was used by which we examined the exact number of mitotically arrested cells. The results in Fig. 2 show that the percentage of mitotically arrested cells (with clear condensed chromosomes) due to *Vinca* alkaloid treatment at 24 h were essentially the same in the absence (61% for vinblastine, 58% for vincristine) or presence (57% for vinblastine, 60% for vincristine) of glucocorticoids. These results clearly suggest that glucocorticoids selectively inhibit *Vinca* alkaloid-induced apoptosis but do not affect the effect of *Vinca* alkaloids on cell cycle arrest. Similar results were also observed in the KB cell line (data not shown).

Vinca Alkaloids and Glucocorticoids Possess Different Regulatory Effects on I κ B α

Our previous studies revealed that paclitaxel induces protein degradation of I κ B α , the cytoplasmic inhibitor of NF- κ B in both BCap37 and KB cells (6). To examine whether *Vinca* alkaloids also affect I κ B α , BCap37 cells were treated with different concentrations of vinblastine or vincristine. As shown in Fig. 3, both agents caused significant degradation of I κ B α . Further, we determined that pretreatment of glucocorticoids could prevent the degradation of I κ B α induced by either vinblastine or vincristine in Bcap37 cells. This result implies that glucocorticoids might inhibit *Vinca* alkaloid-induced apoptosis by interfering with the activation of NF- κ B/I κ B pathway.

Glucocorticoids Do Not Affect *Vinca* Alkaloid-Stimulated IKK Activation

Recent studies have revealed that signal-induced serine phosphorylation and degradation of I κ B α are primarily mediated by activation of IKK complex (18). To determine whether IKK is involved in *Vinca* alkaloid-mediated I κ B α degradation and NF- κ B activation, endogenous IKK activity assays were performed. BCap37 cells were treated with 100 nM vinblastine and vincristine with or without the pretreatment of glucocorticoids. Cell extracts were immunoprecipitated by using antibody against IKK α . IKK activities were measured by using GST-I κ B α protein as the substrate. The results show that phosphorylations of GST-I κ B α protein were remarkably stimulated by vinblastine or vincristine (Fig. 4, A and B). However, glucocorticoids did not affect *Vinca* alkaloid-stimulated phosphorylation of GST-I κ B α protein (Fig. 4C). These results indicated that activation of IKK may play a critical role in the *Vinca* alkaloid-induced I κ B α degradation, while glucocorticoid-inhibited NF- κ B activation is more likely due to the up-regulation of I κ B α protein rather than the inhibition of IKK activity.

Suppression of I κ B α Sensitizes *Vinca* Alkaloid-Induced Apoptosis

To further investigate if IKK-mediated I κ B α degradation plays an important role in the regulation of alkaloid-induced apoptosis, we examined the cytotoxic effects of vinblastine and vincristine in BCap37 I κ B α -ANT5 cell line




Figure 2. Glucocorticoids do not affect mitotic arrest induced by *Vinca* alkaloids. Cells were exposed to 100 nM vinblastine (*Vinb.*) and vincristine (*Vinc.*) with or without pretreatment of glucocorticoids (*Gluc.*) for 24 h before they were harvested for preparation of cytospin slides as described in Materials and Methods. Three hundred cells/slide were counted and only those cells with typical morphological features of condensed chromosomes were identified as mitotically arrested cells. *Columns*, mean percentage of mitotic cells (independent experiments performed in triplicate); *bars*, SD.

that was transfected with antisense I κ B α (6). DNA fragmentation assays show that I κ B α -ANT5 cells treated with a series of concentrations of vinblastine (1–10 nM) or vincristine (1–10 nM) exhibited increasing sensitivity in comparison with the BCap37 cells transfected with empty pcDNA3 vectors (Fig. 5A). The minimum concentrations of vinblastine and vincristine required for induction of the typical fragmented DNA ladders in the control cells were about 10 nM, while DNA fragments were detected when antisense I κ B α transfectants were treated with as low as 1 nM vinblastine or vincristine. Subsequently, comparison of *Vinca* alkaloid cytotoxicity on transfectants was analyzed by MTT assay. As shown in Fig. 5B, BCap37 cells




Figure 3. *Vinca* alkaloids and glucocorticoids possess opposite effects on I κ B α . BCap37 cells were incubated with different concentrations of vinblastine (*Vinb.*) or vincristine (*Vinc.*) for 24 h with or without pretreatment of glucocorticoids (10^{-7} M TA, 24 h; *Gluc.*). Equal amounts of cellular proteins (50 μ g/lane) were fractionated on a 12.5% SDS-PAGE gel and transferred to PVDF membranes followed by immunoblotting with anti-I κ B α polyclonal antibody. β -actin proteins were blotted as a control.




Figure 4. Effects of *Vinca* alkaloids and glucocorticoids on IKK activity. BCap37 cells were treated with the indicated concentrations of vinblastine (*Vinb.*) or vincristine (*Vinc.*) for 24 h with or without pretreatment of glucocorticoids (10^{-7} M TA, 24 h; *Gluc.*). IKK complex was immunoprecipitated with an anti-IKK α antibody, and the immune complex was subject to *in vitro* kinase assay (KA) using GST-I κ B α fusion protein as the substrate. Following SDS-PAGE, the portion of the gel containing the substrate was dried and processed for autoradiography. The portion containing IKK was analyzed by Western blotting for IKK α protein.

transfected with antisense I κ B α at different doses of treatment are more susceptible to *Vinca* alkaloid cytotoxicity than empty vector transfectants. These results indicate that introduction of antisense I κ B α increases the sensitivity of tumor cells in response to *Vinca* alkaloid-induced apoptosis.

Mutant I κ B α Lacking Ser³² and Ser³⁶ Suppresses *Vinca* Alkaloid-Induced NF- κ B Activation

Proteolytic degradation of I κ B α is essential for activation of NF- κ B (9, 19). Previous studies have revealed that the degradation of I κ B α protein is mainly due to the inducible phosphorylation of I κ B α at Ser³² and Ser³⁶ by I κ B α kinase complex (20–22). To further confirm that *Vinca* alkaloids down-regulate I κ B α through induction of I κ B α phosphorylation and degradation, we constructed a mutant I κ B α expression vector by deleting 36 amino acids, including Ser³² and Ser³⁶, from the NH₂ terminus of the I κ B α gene. Such a mutant I κ B α protein cannot be degraded by the I κ B α kinase complex but still possesses the ability to bind to NF- κ B through its interior domain and functions as a supersuppressor of NF- κ B molecules (20, 23). As shown in Fig. 6, BCap37 cells with stable transfection of this mutant I κ B α expressed a smaller size of deletional mutant I κ B α protein, which was not degraded by vinblastine or vincristine. Furthermore, we examined the effect of mutant I κ B α on *Vinca* alkaloid-induced DNA binding activity of NF- κ B. By EMSAs, an increased level of DNA binding activity of NF- κ B was clearly detected in empty vector-transfected BCap37 cells exposed to *Vinca* alkaloids, but this elevated DNA binding activity of NF- κ B by *Vinca* alkaloids was markedly inhibited in the cells transfected with the mutant I κ B α (Fig. 7). These findings demonstrated that the mutant I κ B α could interfere with *Vinca* alkaloid-induced NF- κ B activation.




Figure 5. Transfection with antisense IκBα sensitizes *Vinca* alkaloid-induced apoptosis. BCap37 cells transfected with vector (pcDNA3) only and antisense IκBα (IκBα-ANT5) were incubated with the indicated concentrations of vinblastine (*Vinb.*) or vincristine (*Vinc.*) for 48 h. **A**, DNA fragmentation assays. Cells were harvested and fragmented DNA was analyzed by electrophoresis in 1.2% agarose gel containing 0.1% ethidium bromide. **B**, MTT assays. Points, mean of three separate experiments; bars, SD. There is a statistically significant difference between antisense IκBα-transfected cells and vector-transfected cells treated by *Vinca* alkaloids ($P < 0.01$, Student's *t* test).

Transfection of Mutant IκBα Reduces Sensitivity of Tumor Cells to *Vinca* Alkaloid-Induced Apoptosis

Next, the tumor cells with stable expression of the mutant IκBα were compared with their parental cells to determine whether the expression of the mutant IκBα altered the sensitivity of tumor cells to *Vinca* alkaloid-induced apoptosis. As shown in Fig. 8, wild-type BCap37 cells, empty pcDNA3 vector transfectants, and mutant IκBα cDNA transfectants were treated with vinblastine and vincristine for 48 h followed by flow cytometric assays to determine DNA content. As a result, the cells transfected with the mutant IκBα exhibited more resistance to *Vinca* alkaloid-induced apoptosis. These results indicate that the introduction of the dominant-negative mutant IκBα resulted in the decreased sensitivity of tumor cells to *Vinca* alkaloid-induced apoptosis.

Discussion

The *Vinca* alkaloids vinblastine and vincristine have been successfully used in cancer chemotherapy for the treatment of different types of tumors (24). Their mechanism of action involves disruption of the dynamic reorganization of the microtubule network, resulting in aberrant mitotic formation. Thus, cells are unable to transverse successfully from metaphase to anaphase. Ultimately, the prolonged mitotic arrest in many types of tumor cells leads to apoptotic cell death. However, the mechanism by which *Vinca* alkaloids induce apoptotic cell death in tumor cells is not entirely clear.

In this study, pretreatment with glucocorticoids inhibited *Vinca* alkaloid-induced apoptosis without affecting the G₂-M phase arrest caused by *Vinca* alkaloids (Fig. 1), suggesting that glucocorticoids may interrupt the specific downstream events of *Vinca* alkaloid-induced mitotic

arrest. Another possibility is that *Vinca* alkaloid-induced apoptosis may occur via a pathway independent of mitotic arrest. It has been observed that glucocorticoids selectively inhibited apoptotic cell death induced by paclitaxel, a member of the taxane class of antimicrotubule agent (4, 5). Paclitaxel and *Vinca* alkaloids possess different antimicrotubule mechanisms, but both classes of drugs disrupt the normal structure and function of cellular microtubules and cause mitotic arrest. Thus, the inhibitory action of glucocorticoids on apoptosis induced by those antimicrotubule agents may have broad significance and share a similar molecular mechanism.

It has been reported that the regulation of antimicrotubule-induced apoptosis may involve several genes in various signal pathways, such as the Bcl-2 family, p53, p21, Fas/Fas ligand, caspase family, Myc, etc. (25–27). However, many questions need to be clarified further as to the exact role of these proteins or signaling pathways in the regulation of apoptosis. In our previous studies, we discovered that paclitaxel significantly down-regulated IκBα, the cytoplasmic inhibitor of NF-κB, which in turn promoted NF-κB translocation to the nucleus and its DNA binding activity. In the present study, we demonstrated that *Vinca* alkaloids have similar regulatory effects on the NF-κB/IκBα cascade, which include degradation of IκBα protein and activation of NF-κB (Figs. 3 and 7). Further, by *in vitro* IKK assay, IKK activities were found to be significantly stimulated by vincristine or vinblastine (Fig. 4A), suggesting that IKK might be the common target of different kinds of antimicrotubule agents. Meanwhile,




Figure 6. *Vinca* alkaloids do not degrade mutant IκBα. Equal amounts of cellular proteins (50 μg/lane) from wild-type BCap37 cells (WT), cells transfected with empty pcDNA3 vector (Vector), and cells transfected with mutant IκBα (Mutant IκBα) treated with different concentrations of vinblastine (**A**) and vincristine (**B**) for 24 h were fractionated on 12.5% SDS-PAGE gel and transferred to PVDF membranes followed by immunoblotting with anti-IκBα polyclonal antibody. β-actin protein was used as a control.




Figure 7. Overexpression of mutant I κ B α blocks *Vinca* alkaloid-induced NF- κ B activation. BCap37 cells transfected with pcDNA3 vector only (*Vector*) or mutant I κ B α (*Mutant I κ B α*) were treated with different concentrations of vinblastine (*Vinb.*; **A**) and vincristine (*Vinc.*; **B**) for 24 h. Equal amounts of nuclear cell extracts were subjected to EMSAs with [γ - 32 P]-labeled oligonucleotide encompassing the NF- κ B binding site. NS, nonspecific site.

kinase assays also indicated that glucocorticoids did not interrupt *Vinca* alkaloid-activated IKK activity (Fig. 4C), which provided another piece of evidence that glucocorticoids antagonize *Vinca* alkaloid-induced I κ B α degradation by stimulation of I κ B α protein synthesis rather than interference with the phosphorylation and degradation of I κ B α protein. In addition, because glucocorticoids interfere with the I κ B α /NF- κ B pathway without affecting the G₂-M phase arrest caused by *Vinca* alkaloids, it also suggests that activation of I κ B α /NF- κ B pathway by *Vinca* alkaloids occur through a pathway independent of cell cycle arrest.

I κ B α functions as a key inhibitor of the transcription factor NF- κ B. The exact mechanism of NF- κ B in the modulation of apoptosis is not entirely clear. However, it has been reported that the activation of NF- κ B is able to either induce or block apoptosis depending on different stimuli and different cell types (28). To determine the role of activation of NF- κ B in *Vinca* alkaloid-induced apoptosis, we used the previously established antisense I κ B α -transfected BCap37 cell line I κ B α -ANT5 (6) as a study model. The results indicated that BCap37 cells transfected with antisense I κ B α significantly increased their sensitivity to *Vinca* alkaloid-induced apoptosis (Fig. 5). Because exoge-

nous antisense I κ B α blocks the inhibitory binding of endogenous I κ B α to NF- κ B, this finding implies that *Vinca* alkaloids may induce apoptotic cell death through the activation of the NF- κ B/I κ B signaling pathway.

Furthermore, to investigate the mechanism by which *Vinca* alkaloids activate NF- κ B, we constructed a mutant I κ B α expression vector in which a NH₂-terminal fragment containing Ser³² and Ser³⁶ was deleted. Based on current knowledge, the degradation of I κ B α is mainly a result of the inducible phosphorylation of Ser³² and Ser³⁶. Deletion or substitution of these two amino acids with other residues has been reported to prevent I κ B α from signal-induced phosphorylation (20, 23). Through stable transfection of this mutant I κ B α into wild-type BCap37 cells, we demonstrated that the mutant I κ B α protein was insensitive to IKK-mediated phosphorylation and degradation but still possessed the ability to interact with cytoplasmic NF- κ B and inhibit *Vinca* alkaloid-induced NF- κ B activation (Figs. 6 and 7). Meanwhile, the results from flow cytometric assays and MTT assay revealed that the expression of the mutant I κ B α significantly inhibited *Vinca* alkaloid-induced apoptotic cell death (Fig. 8). These findings further suggest that blockage of NF- κ B activation by the mutant I κ B α disrupts the signaling pathway leading to *Vinca* alkaloid-induced apoptotic cell death. Furthermore, because glucocorticoids antagonize *Vinca* alkaloid-induced I κ B α degradation through similar mechanism as mutant I κ B α , it also suggests that glucocorticoids inhibit *Vinca* alkaloid-induced apoptosis through inhibiting NF- κ B activation. Based on these experimental results, the activation of NF- κ B likely plays the role of promoter in *Vinca* alkaloid-induced apoptosis. It is currently unclear how the activated NF- κ B triggers the apoptotic machinery. NF- κ B has been revealed to regulate the transcription of more than 150 target genes (29). Many of these NF- κ B target genes are believed to be proapoptotic genes, such as Fas/Apo-1 ligand (*FasL*), *ICE*, *c-myc*, and *p53* (13, 30–32). NF- κ B may mediate *Vinca* alkaloid-induced apoptosis through the regulation of the activities of specific genes that eventually trigger the downstream signaling pathway, leading to apoptotic cell death in tumor cells.

In summary, through characterization of the inhibitory effect of glucocorticoids on *Vinca* alkaloid-induced apoptosis in BCap37 and KB cells, we found that *Vinca* alkaloids




Figure 8. Overexpression of mutant I κ B α suppresses vinblastine- and vincristine-induced apoptosis. Wild-type BCap37 cells (*WT*) and cells transfected with pcDNA3 vector only (*Vector*) or mutant I κ B α (*Mutant I κ B α*) were treated with 50 or 100 nM vinblastine (**A**) and vincristine (**B**) for 48 h. Cells were harvested and stained with propidium iodide for flow cytometric analysis. The peaks corresponding to G₁ and G₂-M phases of the cell cycle are indicated. The sub-G₁ peaks (*AP*) represent apoptotic cells.

may induce apoptotic cell death through activation of NF- κ B/I κ B signaling pathway. On contrary, glucocorticoids inhibit *Vinca* alkaloid-induced apoptosis without affecting *Vinca* alkaloid-induced cell cycle arrest through up-regulating I κ B α , which antagonizes NF- κ B activation. These findings suggest that the NF- κ B/I κ B signaling pathway may contribute to the mediation of *Vinca* alkaloid-induced apoptosis in human breast cancer cells.

References

- Rowinsky EK, Donehower RC. Systemic chemotherapy. In Perry MC, editor. The chemotherapy source book. 2nd ed. Baltimore: Williams and Wilkins; 1996. p. 116–40.
- Haskell CM. Antineoplastic agents: cancer treatment. In Haskell CM, editor. Cancer treatment. 4th ed. Philadelphia: W. B. Saunders Co.; 1995. p. 78–165.
- Fan M, Goodwin ME, Birrer MJ, Chambers TC. The c-Jun NH₂-terminal protein kinase/AP-1 pathway is required for efficient apoptosis induced by vinblastine. *Cancer Res*, 2001;61:4450–8.
- Fan W, Cheng LR, Norris JS, Willingham MC. Glucocorticoids selectively inhibit paclitaxel-induced apoptosis in a human breast cancer cell line. *Cell Pharmacol*, 1996;3:435–40.
- Fan W, Schandl CA, Cheng LR, Norris JS, Willingham MC. Glucocorticoids modulate Taxol cytotoxicity in human solid tumor cells. *Cell Pharmacol*, 1996;3:343–8.
- Huang Y, Johnson KR, Norris JS, Fan W. Nuclear factor- κ B/I κ B signaling pathway may contribute to the mediation of paclitaxel-induced apoptosis in solid tumor cells. *Cancer Res*, 2000;60:4426–32.
- Baldwin AS Jr. The NF- κ B and I κ B proteins: new discoveries and insights. *Annu Rev Immunol*, 1996;14:649–83.
- Brown K, Park S, Kanno T, Franzoso G, Siebenlist U. Mutual regulation of the transcriptional activator NF- κ B and its inhibitor, I κ B- α . *Proc Natl Acad Sci USA*, 1993;90:2532–6.
- Baeuerle PA. The inducible transcription factor NF- κ B: regulation by distinct protein subunits. *Biochim Biophys Acta*, 1991;12:141–79.
- Zandi E, Rothwarf DM, Delhase MH, Karin M. The I κ B kinase complex (IKK) contains two kinase subunits, IKK α and IKK β , necessary for I κ B phosphorylation and NF- κ B activation. *Cell*, 1997;91:243–52.
- Grimm S, Bauer MK, Baeuerle PA, Schulze-Osthoff K. Bcl-2 down-regulates the activity of transcription factor NF- κ B induced upon apoptosis. *J Cell Biol*, 1996;134:13–23.
- Beg AA, Baltimore D. An essential role for NF- κ B in preventing TNF- α -induced cell death. *Science*, 1996;274:782–4.
- Wu H, Lozano G. NF- κ B activation of p53: a potential mechanism for suppressing cell growth in response to stress. *J Biol Chem*, 1994;269:20067–74.
- Lin YZ, Yao SY, Veach RA, Torgerson TR, Hawiger J. Inhibition of nuclear translocation of transcription factor NF- κ B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence. *J Biol Chem*, 1995;270:14255–8.
- Ryan KM, Ernst MK, Rice NR, Vousden KH. Role of NF- κ B in p53-mediated programmed cell death. *Nature*, 2000;404:892–6.
- Huang Y, Fan W. I κ B kinase activation is involved in regulation of paclitaxel-induced apoptosis in human tumor cell lines. *Mol Pharmacol*, 2002;61:105–13.
- Johnson KR, Wang L, Miller MC, Willingham MC, Fan W. 5-Fluorouracil interferes with paclitaxel cytotoxicity against human solid tumor cells. *Clin Cancer Res*, 1997;3:1739–45.
- May MJ, Ghosh S. I κ B kinase: kinsmen with different crafts. *Science*, 1999;284:271–3.
- Sun S, Elwood J, Greene WC. Both amino- and carboxyl-terminal sequences within I κ B α regulate its inducible degradation. *Mol Cell Biol*, 1995;16:1058–65.
- Brown K, Gerstberg S, Carlson L, Franzoso G, Siebenlist U. Control of I κ B α proteolysis by site-specific, signal-induced phosphorylation. *Science*, 1995;267:1485–8.
- Traenckner EBM, Pahl HL, Henkel T, Schmidt KN, Wilk S, Baeuerle PA. Phosphorylation of human I κ B- α on serines 32 and 36 controls I κ B- α proteolysis and NF- κ B activation in response to diverse stimuli. *EMBO J*, 1995;14:2876–83.
- DiDonato JA, Hayakawa M, Rothwarf DM, Zandi E, Karin M. A cytokine responsive I κ B kinase that activates the transcription factor NF- κ B. *Nature*, 1997;388:548–54.
- Shinohara T, Miki T, Nishimura N, et al. Nuclear factor- κ B-dependent expression of metastasis suppressor KA11/CD82 gene in lung cancer cell lines expressing mutant p53. *Cancer Res*, 2001;61:673–8.
- Kreis W. Concepts, mechanisms, and new targets for chemotherapy. In: Muggia FM, editor. Boston: IRL Press; 1995. p. 163–82.
- Wang LG, Liu XM, Kreis W, Budman DR. The effect of antimicrotubule agents on signal transduction pathways of apoptosis: a review. *Cancer Chemother Pharmacol*, 1999;44:355–61.
- Simizu S, Takada M, Umezawa K, Imoto M. Requirement of caspase-3 (-like) protease-mediated hydrogen peroxide production for apoptosis induced by various anticancer drugs. *J Biol Chem*, 1998;273:26900–7.
- Fan M, Goodwin M, Vu T, Brantley-Finley C, Gaarde WA, Chambers TC. Vinblastine-induced phosphorylation of Bcl-2 and Bcl-XL is mediated by JNK and occurs in parallel with inactivation of the Raf-1/MEK/ERK cascade. *J Biol Chem*, 2000;275:29980–5.
- Barkett M, Gilmore TD. Control of apoptosis by Rel/NF- κ B transcription factors. *Oncogene*, 1999;18:6910–24.
- Chan H, Bartos DP, Owen-Schaub LB. Activation-dependent transcriptional regulation of the human Fas promoter requires NF- κ B p50-p65 recruitment. *Mol Cell Biol*, 1999;19:2098–108.
- Ibrado AM, Liu L, Bhalla K. Bcl-xL overexpression inhibits progression of molecular events leading to paclitaxel-induced apoptosis of human acute myeloid leukemia HL-60 cells. *Cancer Res*, 1997;57:1109–15.
- La Rosa FA, Pierce JW, Sonenshein GE. Differential regulation of the c-myc oncogene promoter by the NF- κ B rel family of transcription factors. *Mol Cell Biol*, 1994;14:1039–44.
- Pahl H. Activators and target genes of Rel/NF- κ B transcription factors. *Oncogene*, 1999;18:6853–66.

Molecular Cancer Therapeutics

Regulation of *Vinca* alkaloid-induced apoptosis by NF- κ B/I κ B pathway in human tumor cells

Yi Huang, Yong Fang, Jinmin Wu, et al.

Mol Cancer Ther 2004;3:271-277.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/3/3/271>

Cited articles This article cites 29 articles, 17 of which you can access for free at:
<http://mct.aacrjournals.org/content/3/3/271.full#ref-list-1>

Citing articles This article has been cited by 6 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/3/3/271.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/3/3/271>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.