

# Inhibition of insulin-like growth factor-I receptor (IGF-IR) signaling and tumor cell growth by a fully human neutralizing anti-IGF-IR antibody

Yan Wang,<sup>1</sup> Judith Hailey,<sup>1</sup> Denise Williams,<sup>4</sup> Yaolin Wang,<sup>1</sup> Philip Lipari,<sup>1</sup> Michael Malkowski,<sup>1</sup> Xiaoying Wang,<sup>1</sup> Lei Xie,<sup>2</sup> Guanghua Li,<sup>2</sup> Deba Saha,<sup>2</sup> Wai Lam W. Ling,<sup>2</sup> Susan Cannon-Carlson,<sup>2</sup> Robert Greenberg,<sup>2</sup> Robert A. Ramos,<sup>1</sup> Robert Shields,<sup>3</sup> Leonard Presta,<sup>3</sup> Peter Brams,<sup>4</sup> W. Robert Bishop,<sup>1</sup> and Jonathan A. Pachter<sup>1</sup>

Departments of <sup>1</sup>Tumor Biology and <sup>2</sup>Biotechnology Development, Schering-Plough Research Institute, Kenilworth, New Jersey; <sup>3</sup>Department of Biological Research, DNAX Research Institute of Molecular and Cellular Biology, Palo Alto, California; and <sup>4</sup>Medarex, Milpitas, California

## Abstract

Insulin-like growth factor-I receptor (IGF-IR) plays an important role in tumor cell growth and survival. On ligand stimulation, IGF-IR, a receptor tyrosine kinase, phosphorylates tyrosine residues on two major substrates, IRS-1 and Shc, which subsequently signal through the Ras/mitogen-activated protein kinase and phosphatidylinositol 3-kinase/AKT pathways. Here, we describe the characterization of a fully human anti-IGF-IR monoclonal antibody 19D12 that inhibits IGF binding and autophosphorylation of both IGF-IR/IGF-IR homodimers and IGF-IR/insulin receptor heterodimers. 19D12 does not recognize insulin receptor homodimers. In addition to inhibiting IGF-IR autophosphorylation, 19D12 also inhibits IRS-1 phosphorylation and activation of the major downstream signaling molecules AKT and extracellular signal-regulated kinase 1/2. Furthermore, the antibody down-regulates the total IGF-IR protein level and can exhibit antibody-dependent cellular cytotoxicity activity against a non-small cell adenocarcinoma cell line *in vitro* in the presence of isolated human natural killer cells. 19D12 binds tightly to the receptor, with an affinity of 3.8 pmol/L as measured by KinExA. In cell culture, 19D12 inhibits proliferation and soft agar growth of various tumor

cell lines. *In vivo*, 19D12 inhibits the tumor growth of a very aggressive human ovarian tumor xenograft model A2780. These data support the development of this anti-IGF-IR monoclonal antibody as a promising anticancer agent. [Mol Cancer Ther 2005;4(8):1214–21]

## Introduction

Insulin-like growth factor-I receptor (IGF-IR) plays an important role in cell growth and differentiation (1). Overexpression of IGF-IR results in aberrant cell proliferation and malignant transformation (2–4). Cells with IGF-IR deleted are resistant to oncogenic transformation (5). In addition to its role in cell proliferation, IGF-IR plays an important role in tumor cell survival. Overexpressed IGF-IR can protect cells from apoptosis induced by serum withdrawal, tumor necrosis factor, etoposide, or interleukin-3 withdrawal *in vitro* (6–10). Epidemiologic data have linked elevated plasma IGF-I level with prostate, breast, lung, and colon cancer risk (11–14). IGF-II is frequently overexpressed in colorectal, neuroblastoma, liver, adrenocortical, and Wilms' tumors (15–20). IGF-IR is the signaling receptor for both IGF-I and IGF-II ligands and is overexpressed in some human cancers as well (21–26). Therefore, IGF-IR is an attractive antitumor target.

Inhibition of IGF-IR signaling by various methods, including antisense, dominant-negative truncation, small-molecule kinase inhibitors or neutralizing antibodies, has resulted in growth inhibition and apoptosis of tumor cells (27–40). We show here that a fully human anti-IGF-IR neutralizing antibody inhibits IGF-IR signaling in cell culture and inhibits tumor growth in xenograft models. The antibody was obtained using the Medarex's HuMab transgenic mouse technology (41). Therefore, it is a fully human antibody and is expected to show minimal immunogenicity in humans. Expression of IGF-IR on many tumor cell types suggests that this anti-IGF-IR monoclonal antibody has the potential to treat patients for many different types of cancer.

## Materials and Methods

### Cells and Reagents

Rat1/hIR $\beta$  cells are a generous gift from Dr. Byung-Hak Jhun (Pusan National University, Busan, Korea). Anti-IGF-IR antibody C-20 and anti-insulin receptor antibody C-19 are purchased from Santa Cruz (Santa Cruz, CA). Anti-phosphotyrosine antibody 4G10 is from Upstate Biotechnology (Charlottesville, VA). Antibodies for detecting AKT, phospho-AKT, mitogen-activated protein kinase (MAPK), phospho-MAPK, and phospho-IRS-1 are from Cell Signaling Technology (Beverly, MA). CellTiter-Glo reagents are

Received 2/17/05; revised 4/14/05; accepted 6/1/05.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Note:** Current address of J. A. Pachter is Cancer Biology, OSI Pharmaceuticals, Inc., Farmingdale, New York.

**Requests for reprints:** Yan Wang, Department of Oncology, Schering-Plough Research Institute, 2015 Galloping Hill Road, K15-4600, Kenilworth, NJ 07033. Phone: 908-740-7326; Fax: 908-740-3918. E-mail: yan.wang@spcorp.com

Copyright © 2005 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-05-0048

purchased from Promega (Madison, WI). Human natural killer cells were purified with antibody-coupled magnetic bead kit from Miltenyi Biotec (Auburn, CA). Antibody-dependent cellular cytotoxicity was determined with a cytotoxicity detection kit for lactate dehydrogenase from Roche (Indianapolis, IN). [<sup>125</sup>I]IGF-I is from Amersham (Piscataway, NJ).

#### Receptor Autophosphorylation Assay

Antibodies were added to cells for 30 minutes. Cells were then stimulated with 20 ng/mL IGF-I for 5 to 10 minutes at 37°C, washed twice with cold PBS containing 0.1 mmol/L sodium vanadate, and lysed in lysis buffer [50 mmol/L HEPES (pH 7.4), 150 mmol/L NaCl, 10% glycerol, 1% Triton X-100, 1.5 mmol/L MgCl<sub>2</sub>, protease inhibitors, 2 mmol/L sodium vanadate]. Lysates were incubated on ice for 30 minutes and then centrifuged at 13,000 rpm for 10 minutes at 4°C. Protein concentrations of the lysates were measured with a Coomassie Plus Protein Assay (Pierce, Rockford, IL). Lysates were then subjected to immunoprecipitation and Western blot analysis.

#### Immunoprecipitation and Western Blot Analysis

IGF-IR was immunoprecipitated using 1 μg antibody (C-20) and 15 μL of 50% protein G agarose. The precipitates were washed four times with lysis buffer. Precipitated proteins were separated on 10% SDS-PAGE, transferred to a nitrocellulose filter, probed with either anti-IGF-IR or anti-phosphotyrosine (4G10) antibodies, and visualized by enhanced chemiluminescence (Amersham).

For direct Western blot analysis, equal amounts of cell lysates were separated on 10% SDS-PAGE, transferred to nitrocellulose filters, probed with a desired antibody, and visualized by enhanced chemiluminescence.

#### Ligand and Receptor Binding

Millipore 96-well filter plates (1.2 μm Durapore, Bedford, MA) were prewet with 0.5% bovine serum albumin/PBS for 2 hours at 4°C. The buffer was then removed with a vacuum manifold. Various concentrations of 6× control or test antibody were added to the wells (25 μL). The [<sup>125</sup>I]IGF-I ligand was then added to the wells at a final concentration of 0.375 nmol/L in bovine serum albumin/PBS. Cells were harvested with cell dissociation solution, counted with trypan blue, and resuspended in 0.5% bovine serum albumin/PBS to a cell density of 1 × 10<sup>5</sup> to 3 × 10<sup>5</sup>/mL. Cells (10,000-30,000; 100 μL) were added to each well. The plate was shaken at 4°C for 1 hour. The plate was then aspirated and washed thrice with ice-cold PBS. The filters were punched out and counted on an ICN Micromedex gamma counter (Aurora, OH).

#### Antibody-Dependent Cellular Cytotoxicity

NCI-H322 cells were trypsinized for 3 minutes, resuspended in assay medium (RPMI 1640 containing 1% fetal bovine serum) at 4 × 10<sup>5</sup>/mL, and transferred to 96-well round-bottomed plate containing 100 ng/mL 19D12. Cells were opsonized for 30 minutes at room temperature. Serial 2-fold dilutions of purified natural killer cells were prepared in a separate plate in assay medium and transferred to antibody-treated tumor cell targets. Cells were incubated at

37°C for 4 hours. Supernatants were harvested and lactate dehydrogenase content was measured. Percent lactate dehydrogenase release by antibody-dependent cellular cytotoxicity was calculated as a percentage of total release determined by Triton X-100 lysis.

#### Proliferation Assay

Cells were trypsinized, counted, and resuspended at 25,000 cells/mL in 10% HI-FBS RPMI 1640 containing nonessential amino acids, L-glutamine, MEM vitamins, and PS. Cell suspension (2,500 cells; 100 μL) was added to each well of BD Falcon 96-well black, clear-bottomed TC-treated plate (Bedford, MA). Cells were allowed to attach and spread overnight at 37°C. The 10% RPMI 1640 was replaced with 100 μL of 2% RPMI 1640 the following day. All treatments were prepared in 2% RPMI 1640 at 20× concentration and serially diluted. Every test point was prepared in triplicate on separate assay plates. Cell proliferation was measured using the CellTiter-Glo Luminescent Cell Viability Assay 96 hours after treatment. Luminescence was detected on a Wallac (Boston, MA) 420 plate reader with stacker.

#### Xenograft Model in Mice


Four million A2780 human ovarian carcinoma cells in 50% growth factor-reduced Matrigel were inoculated s.c. into each nude mouse. 19D12 treatment was initiated when the tumor size reached ~50 mm<sup>3</sup> (defined as day 0). Two additional doses were given at days 3 and 7. Tumor volumes were measured by Labcat.

## Results

### 19D12 Inhibits IGF-IR Autophosphorylation and Competes with Ligand Binding to the Receptor

Medarex's HuMab mice (41) were alternately immunized with two different human IGF-IR antigens: (a) HEK-293 cells stably overexpressing full-length IGF-IR and (b) purified soluble IGF-IR receptor. Mouse serum was titered based on the presence of antigen-specific activity measured by ELISA. Splens from high-titer mice were fused and the resulting hybridomas were screened for anti-IGF-IR activity by both a cell-based ELISA and flow cytometric analysis. Antigen-positive supernatants were further evaluated in a cell-based autophosphorylation assay to select neutralizing antibodies. Among the 13 neutralizing antibodies identified, 19D12 is the most potent and has a binding affinity of 3.8 pmol/L determined by KinExA (data not shown).

Figure 1A shows the effect of 19D12 on IGF-IR autophosphorylation in the human ovarian tumor cell line A2780. After preincubation with 19D12 for 30 minutes, cells were stimulated with IGF-I (20 ng/mL) for 5 minutes. Cell lysates were made, and IGF-IR was immunoprecipitated with an anti-IGF-IR-specific antibody. The blot was probed with the anti-phosphotyrosine antibody 4G10. Therefore, the band on *top* represents the level of the phospho-IGF-IR in cell lysates. 19D12 completely inhibited the phosphorylation of IGF-IR on ligand stimulation in a dose-dependent manner. The blot was then stripped and reprobed with anti-IGF-IR antibody to ensure equivalent loading of IGF-IR in all lanes (*bottom*).



**Figure 1.** Inhibition of receptor autophosphorylation and ligand binding to the receptor. **A**, A2780 cells were treated with various concentrations of 19D12 antibody (lane 4, 20 nmol/L; lane 5, 2 nmol/L; lane 6, 0.2 nmol/L) or control IgG (lane 3, 20 nmol/L) for 0.5 h before stimulation with IGF-I (20 ng/mL). Each cell lysate (400  $\mu$ g) was subjected to immunoprecipitation with an IGF-IR-specific antibody. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with anti-phosphotyrosine antibody 4G10. Lane 1, untreated cells; lanes 2 to 6, IGF-I (20 ng/mL)-treated cells. The blot was then stripped and reprobed with anti-IGF-IR antibody (bottom). **B**, A2780 cells were treated with various concentrations of 19D12 antibody (lane 3, 20 nmol/L; lane 4, 2 nmol/L; lane 5, 0.2 nmol/L) or control IgG (lane 6, 20 nmol/L) for 0.5 h before stimulation with human plasma. Each cell lysate (400  $\mu$ g) was subjected to immunoprecipitation with an IGF-IR-specific antibody. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with the anti-phosphotyrosine antibody 4G10. Lane 1, untreated cells; lanes 2 to 6, IGF-I (20 ng/mL)-treated cells. The blot was then stripped and reprobed with anti-IGF-IR antibody (bottom). **C**, various concentrations of 19D12 and [ $^{125}$ I]IGF-I were added to a 96-well filter plate. IGF-IR-transfected HEK-293 cells (20,000 per well) were added to each well to initiate binding. The plate was shaken for 1 h, aspirated, washed, and counted on a gamma counter.

In addition to the inhibition of IGF-IR autophosphorylation stimulated by IGF-I ligand, 19D12 also inhibited IGF-IR autophosphorylation stimulated by human plasma (Fig. 1B). Human plasma has a natural balance of IGF-I, IGF-II, and IGF-binding proteins.

To determine whether 19D12 inhibits IGF-I binding to the receptor, ligand-receptor binding was measured using [ $^{125}$ I]IGF-I as the ligand. Data in Fig. 1C show that 19D12 displaces IGF-I binding to the IGF-IR receptor with an  $IC_{50}$  value of 2.7 nmol/L.


### 19D12 Binds and Inhibits the IGF-IR/Insulin Receptor Heterodimeric Receptors but not the Insulin Receptor Homodimeric Receptors

Insulin receptor is a closely related receptor tyrosine kinase and shares high sequence homology with IGF-IR (42, 43). To test the selectivity of the anti-IGF-IR antibody 19D12, direct Western blotting experiments were done with 19D12 using lysates from cells expressing a high endogenous level of human IGF-IR as well as lysates from stably transfected Rat1 cells that overexpress a high level of human insulin receptor (hIR $\beta$  cells). 19D12 only recognized a protein of ~135,000 dalton molecular weight ( $\alpha$ -subunit of IGF-IR) in cells containing human IGF-IR but did not recognize the overexpressed human insulin receptor (data not shown). To further show the functional selectivity of 19D12, an autophosphorylation assay in hIR $\beta$  cells was done. Cells were stimulated with insulin (20 ng/mL) after preincubation with 19D12 or an anti-insulin receptor neutralizing antibody for 30 minutes. Cell lysates were made, and insulin receptor was immunoprecipitated with an anti-insulin receptor-specific antibody. The blot was probed with the anti-phosphotyrosine antibody 4G10. As shown in Fig. 2A, 19D12 did not inhibit the phosphorylation of insulin receptor on insulin stimulation, whereas the anti-insulin receptor-positive control neutralizing antibody inhibited insulin receptor autophosphorylation.

Both the IGF-IR and the insulin receptor are tetrameric complexes consisting of two extracellular  $\alpha$ -subunits that bind the ligands and two  $\beta$ -subunits that have ligand-dependent tyrosine kinase activity. In cells that coexpress IGF-IR and insulin receptor, heterodimeric receptors can form between one half of the receptor consisting of one IGF-IR  $\alpha$ -subunit and one IGF-IR  $\beta$ -subunit and the other half of the receptor consisting of one insulin receptor  $\alpha$ -subunit and one insulin receptor  $\beta$ -subunit (44). The IGF-IR/insulin receptor heterodimeric receptor behaves more like IGF-IR than insulin receptor, exhibiting affinities for IGF-I and IGF-II similar to the IGF-IR homodimer. The heterodimeric receptor binds insulin with much lower affinity (44). To examine whether 19D12 can bind the IGF-IR/insulin receptor heterodimeric receptor and inhibit its signaling, we did immunoprecipitation experiments using 19D12 as the immunoprecipitating antibody in A2780 cells that endogenously express both IGF-IR (150,000 IGF-IR/cell) and insulin receptor (30,000 insulin receptor/cell). As shown in Fig. 2B, both insulin receptor (left) and IGF-IR (right) were detected in 19D12 immunoprecipitates, suggesting that 19D12 can bind to both the IGF-IR homodimer and the IGF-IR/insulin receptor heterodimer. The phosphorylation status of the heterodimeric receptor was also examined in the autophosphorylation assay. As shown in Fig. 2C, 19D12 was able to completely inhibit IGF-I-stimulated autophosphorylation of the heterodimeric receptors immunoprecipitated with an anti-insulin receptor antibody.

### 19D12 Down-Regulates IGF-IR Level and Blocks IGF-IR Signaling

In addition to inhibiting receptor autophosphorylation, we have also found that 19D12 down-regulates the IGF-IR




**Figure 2.** 19D12 activity against insulin receptor (*IR*) receptors and IGF-IR/insulin receptor hybrid receptors. **A**, Rat1/hIR $\beta$  cells were treated with 20 nmol/L 19D12 (*lane 4*) or neutralizing antibody against insulin receptor (*lane 3*) for 0.5 h before stimulation with insulin (20 ng/mL) for 5 min. Each cell lysate (400  $\mu$ g) was subjected to immunoprecipitation with an insulin receptor – specific antibody. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with the anti-phosphotyrosine antibody 4G10. *Lane 1*, untreated cells; *lane 2*, insulin (20 ng/mL) – treated cells. The blot was then stripped and re-probed with anti – insulin receptor antibody (*bottom*). **B**, 19D12 (*lane 2*) or protein G agarose alone (*lane 1*) was used to immunoprecipitate 400  $\mu$ g each of A2780 cell lysate. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with an anti – insulin receptor antibody (*left*). The blot was then stripped and re-probed with anti – IGF-IR antibody (*right*). **C**, A2780 cells were treated with 20 nmol/L 19D12 (*lane 3*) or control IgG antibody (*lane 4*) for 0.5 h before stimulation with IGF-I (20 ng/mL) for 5 min. Each cell lysate (400  $\mu$ g) was subjected to immunoprecipitation with an insulin receptor – specific antibody. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with the anti-phosphotyrosine antibody, 4G10. *Lane 1*, untreated cells; *lane 2*, IGF-I (20 ng/mL) – only treated cells. The blot was stripped and re-probed with anti-insulin receptor antibody (*middle*) and subsequently stripped again and re-probed with anti – IGF-IR (*bottom*).


protein level. As shown in Fig. 3A, 19D12 down-regulated IGF-IR protein expression on 4 hours of treatment in a dose-dependent manner. The down-regulation was observed starting at 1-hour incubation, and the extent of down-regulation was maximal at 4 hours (data not shown). The isotype control antibody or IGF-I ligand did not affect the IGF-IR level with up to overnight incubation (data not shown).

To further evaluate the inhibitory effects of 19D12 in the IGF signaling pathway, the phosphorylation status of downstream signaling molecules of the IGF pathway was

investigated. Activation status of IRS-1, the key immediate downstream signaling molecule of IGF-IR, and activation status of AKT and MAPK, key effector molecules of the two major IGF-IR signaling pathways, were examined using antibodies that are specific for the phosphorylated forms of the respective target molecules after 19D12 treatment. As shown in Fig. 3B and C, 19D12 potently inhibited IGF-I-stimulated phosphorylation of IRS-1, AKT, and MAPK.



**Figure 3.** Down-modulation of IGF-IR and inhibition of IRS-1, AKT, and MAPK activation. **A**, MCF7 cells were treated with various concentrations of 19D12 antibody (*lane 2*, 20 nmol/L; *lane 3*, 4 nmol/L; *lane 4*, 0.8 nmol/L; *lane 5*, 0.16 nmol/L; *lane 6*, 0.032 nmol/L) for 4 h. Each cell lysate (50  $\mu$ g) was separated by 10% SDS-PAGE and transferred to a nitrocellulose blot. The blot was probed with anti-IGF-IR antibody (*top*), stripped, and re-probed with anti-actin antibody (*bottom*). **B**, A2780 cells were treated with various concentrations of 19D12 antibody (*lanes 2* and *4*, 20 nmol/L; *lane 5*, 2 nmol/L; *lane 6*, 0.2 nmol/L) for 4 h. Each cell lysate (50  $\mu$ g) was separated by a 10% SDS-PAGE and transferred to a nitrocellulose blot. The blot was probed with anti-phospho-IRS-1 antibody (*top*). *Bottom*, loading control where the blot was probed with anti-IRS-1 antibody. *Lane 1*, untreated cells; *lane 3*, IGF-I (20 ng/mL) – treated cells. **C**, A2780 cells were treated with various concentrations of 19D12 antibody (*lane 4*, 20 nmol/L; *lane 5*, 2 nmol/L; *lane 6*, 0.2 nmol/L; *lane 7*, 0.02 nmol/L) for 4 h. Each cell lysate (50  $\mu$ g) was separated by a 10% SDS-PAGE and transferred to a nitrocellulose blot. The blot was probed with anti-phospho-AKT antibody first (*top*), then re-probed with anti-phospho – extracellular signal-regulated kinase 1/2 (ERK1/2) antibody (*middle*). The blot was subsequently stripped and probed with anti-AKT and anti-ERK1/2 antibodies (*bottom*). *Lane 1*, untreated cells; *lanes 2* to *7*, IGF-I (20 ng/mL) treated cells; *lane 3*, cells treated with 20 nmol/L control IgG.



**Figure 4.** Activities of F(ab) fragment of 19D12. **A**, MCF7 cells were treated with various concentrations of full-length 19D12 antibody (lane 3, 20 nmol/L; lane 4, 4 nmol/L; lane 5, 0.8 nmol/L) or F(ab) fragment of 19D12 (lane 6, 100 nmol/L; lane 7, 20 nmol/L; lane 8, 4 nmol/L) for 0.5 h before stimulation with IGF-I (20 ng/mL). Each cell lysate (400  $\mu$ g) was subjected to immunoprecipitation with an IGF-IR-specific antibody. The precipitated cell lysates were separated on 10% SDS-PAGE and transferred to a nitrocellulose blot. The filter was probed with anti-phosphotyrosine antibody 4G10. Lane 1, untreated cells; lane 2, cells treated with 20 nmol/L control IgG. **B**, MCF7 cells were treated with various concentrations of full-length 19D12 antibody (lane 2, 20 nmol/L; lane 3, 4 nmol/L; lane 4, 0.8 nmol/L) or F(ab) fragment of 19D12 (lane 5, 100 nmol/L; lane 6, 20 nmol/L) for 4 h. Each cell lysate (50  $\mu$ g) was separated by a 10% SDS-PAGE and transferred to a nitrocellulose blot. The blot was probed with anti-IGF-IR antibody. Lane 1, cells treated with 20 nmol/L control IgG.

#### F(ab) Fragment of 19D12 Inhibits Receptor Autophosphorylation but Does Not Down-Regulate Receptor Expression

The full-length 19D12 antibody is a bivalent molecule that consists of two identical heavy chains and two identical light chains. Functionally, the IgG molecule is divided into three domains: F(ab), the antigen-binding domains; the flexible hinge domain; and the Fc effector domain that interacts with complement and immune effector cells. We were interested to determine whether a single F(ab) fragment of 19D12 could induce both antagonist activity and receptor down-regulation as observed with the full-length antibody. A single F(ab) of 19D12 was made, and it was able to inhibit IGF-IR autophosphorylation in response to IGF-I as shown in Fig. 4A. However, the F(ab) fragment failed to down-regulate IGF-IR expression, suggesting that the receptor down-regulation requires a bivalent IgG molecule (Fig. 4B). The F(ab) fragment of 19D12 exhibited very similar IGF-IR affinity to the full-length 19D12 (data not shown).

#### Induction of Antibody-Dependent Cellular Cytotoxicity *In vitro*

The Fc portion of an IgG1 isotype monoclonal antibody can potentially mediate effector functions by selective binding to Fc $\gamma$ RIII (CD16) expressed by natural killer cells to mediate antibody-dependent cellular cytotoxicity of tumor cells. Some therapeutic monoclonal antibody agents in the clinic, such as Rituxan and Herceptin, may owe part of their antitumor efficacy to their ability to induce antibody-dependent cellular cytotoxicity. We investigated the ability of 19D12/IgG1 to induce antibody-dependent


cellular cytotoxicity *in vitro*. Data in Fig. 5 show that 19D12/IgG1 can augment killing of NCI-H322 non-small cell lung carcinoma tumor cells in the presence of isolated human natural killer cells. In contrast, the IgG4 isotype of 19D12 did not induce antibody-dependent cellular cytotoxicity above background as expected.

#### Inhibition of Tumor Cell Proliferation by 19D12


To further investigate the effect of IGF-IR signaling inhibition by 19D12 on cell growth, the CellTiter-Glo Luminescent Cell Viability Assay was used to measure tumor cell proliferation in culture after 19D12 treatment. As shown in Fig. 6, 19D12 dose-dependently inhibited growth of A2780 ovarian tumor cells. A [ $^{14}$ C]thymidine incorporation assay was also used to measure proliferation and similar results were obtained (data not shown).

#### Inhibition of Human Ovarian Xenograft Tumor Growth by 19D12

We tested the antitumor activity of 19D12 in a human tumor xenograft model. Human A2780 ovarian tumor cells were implanted s.c. in nude mice in the presence of Matrigel. Treatment with 19D12 (twice weekly) was initiated after the tumors reached 50 mm<sup>3</sup>. The study was terminated after the tumors in the control group reached >1,000 mm<sup>3</sup>. As shown in Fig. 7A, 19D12 dose-dependently inhibited the tumor growth of this aggressive human ovarian cancer line in mice. Only three injections of 19D12 were given during the study. At the end of the study, tumors were removed and tumor lysates were prepared. Total IGF-IR levels in tumors were measured by Western analysis. As shown in Fig. 7B, the IGF-IR level in treated animals was lower than the IGF-IR in control IgG-treated animals. This shows that the receptor down-regulation by 19D12 occurs both in cell culture and *in vivo*. This feature can potentially be used as a surrogate marker for anti-IGF-IR monoclonal antibody therapy.



**Figure 5.** Induction of antibody-dependent cellular cytotoxicity by 19D12. NCI-H322 cells and various antibodies were added to a 96-well plate and opsonized with 100 ng/mL 19D12 for 30 min at room temperature. Purified human natural killer cells were then added at various ratios to numbers of target H322 cells. Cells were incubated at 37°C for 9 h. Lactate dehydrogenase released was then measured as an indicator of cell cytotoxicity. Each data point was assayed in quadruple in the experiment.



**Figure 6.** Inhibition of tumor cell proliferation by 19D12. Various concentrations of 19D12 were added to A2780 human ovarian carcinoma cells (2,500 cells per well) for 4 d in RPMI 1640 with 2% fetal bovine serum. Cell proliferation was measured using the CellTiter-Glo Luminescent Cell Viability Assay.


## Discussion

IGF-IR plays an important role in both transformation and survival. Cells from IGF-IR knockout mice are resistant to various viral and cellular oncogenes, including SV40 T Ag and Ras (5). Inhibition of the IGF/IGF-IR axis may contribute in part to the efficacies observed with some clinically effective anticancer therapeutics, such as tamoxifen (45) and 17-allylamino geldanamycin (46). In addition to its roles in transformation and survival, the IGF/IGF-IR system also regulates vascular endothelial growth factor

production, suggesting a role in angiogenesis (47). IGF-IR is therefore an attractive anticancer target. We described here the generation and characterization of a fully human neutralizing anti-IGF-IR antibody, 19D12. 19D12 specifically recognizes both IGF-IR homodimers and IGF-IR/insulin receptor heterodimers. However, it is specific and does not recognize insulin receptor homodimers (Fig. 2). This selectivity relative to insulin receptor homodimers may be an advantage over potential IGF-IR kinase inhibitors, because this antibody might be less likely to induce hyperglycemia. 19D12 has a high affinity for IGF-IR and inhibits the receptor activation stimulated by both purified IGFs and human plasma (Fig. 1). In addition to inhibiting receptor activation, 19D12 down-regulates the IGF-IR receptor at the protein level, shuts down the signaling of the receptor as measured by phosphorylation of IRS-1, AKT, and MAPK, and has the ability to induce antibody-dependent cellular cytotoxicity (Figs. 3 and 5).

To further understand the mechanism of the down-regulation, the F(ab) fragment of 19D12 was made. The data in Fig. 4 show that receptor down-regulation requires the bivalency of the full-length antibody or minimally an F(ab')<sub>2</sub> fragment. Both neutralization and down-regulation activities could contribute to the antitumor activities observed *in vivo*. It would be interesting to examine the efficacy of the F(ab) fragment *in vivo* to further understand the functional importance of IGF-IR receptor down-regulation.

The observations that the single F(ab) is able to inhibit receptor autophosphorylation and that it has similar affinity to the full-length antibody suggest that 19D12 interacts with



**Figure 7.** Inhibition of A2780 tumor growth in mice. **A**, four million A2780 human ovarian carcinoma cells in 50% growth factor – reduced Matrigel were inoculated s.c. into nude mice. 19D12 treatment was initiated when the tumor size reached ~50 mm<sup>3</sup> (day 0). Two additional doses were given at days 3 and 7. Tumor volumes were measured by Labcat. **B**, each tumor lysate (1 mg) was separated by a 10% SDS-PAGE (lanes 1-8, tumor lysates from control animals; lanes 9-16, tumor lysates from 19D12-treated animals) and transferred to nitrocellulose blots. The blots were probed with anti-IGF-IR antibody.

an epitope on one arm of the IGF-IR homodimer. Consistent with this hypothesis, 19D12 is able to bind and inhibit the signaling of IGF-IR/insulin receptor heterodimers (Fig. 2B and C). This activity is important because the presence of IGF-IR/insulin receptor heterodimers has been reported in human tumor samples and these heterodimers behave functionally similar to IGF-IR homodimers (44).

In recent years, IGF-IR, as an anticancer target, has been pursued actively due to its importance in tumor cell growth and transformation. Because it is a receptor tyrosine kinase, two strategies have been employed: making a neutralizing antibody against its extracellular domain and developing a small-molecule kinase inhibitor against its cytoplasmic kinase domain (36–40, 48–52). The potential advantages of an antibody approach over the small-molecule approach include selectivity against the insulin receptor and potential for increased antitumor efficacy through receptor down-regulation and antibody-dependent cellular cytotoxicity. In a xenograft model, 19D12 inhibited tumor growth of a very aggressive human ovarian cancer line A2780. In addition to the activities described here with A2780 ovarian cancer, 19D12 also has shown potent activities against a wide variety of other human tumor cell lines *in vivo*, including MCF7 breast cancer cells and NCI-H322 non-small cell lung carcinoma cells (data not shown).

In addition to potent antitumor activity as a single agent, 19D12 also displays an additive effect with several approved chemotherapeutic agents and with some targeted antitumor agents (data not shown). Some experiments conducted recently with both cultured cells and primary tumor cells have suggested that IGF-IR can confer resistance to both anti-HER-2 and anti-epidermal growth factor receptor therapies (53, 54). This anti-IGF-IR neutralizing antibody 19D12 is therefore a promising agent both as monotherapy and for use in combination therapy for a wide range of cancer patients.

#### Acknowledgments

We thank Dr. Richard W. Bond for the stable HEK-293 cells that overexpress IGF-IR, Dr. Mohan Srinivasan for the F(ab) fragment, and Dr. Byung-Hak Jhun for the Rat1 cell line transfected with human insulin receptor (hIR $\beta$  cells).

#### References

- Butler AA, Yakar S, Gewolb IH, Karas M, Okubo Y, LeRoith D. Insulin-like growth factor-I receptor signal transduction: at the interface between physiology and cell biology. *Comp Biochem Physiol* 1998;121:19–26.
- Rubini M, Hongo A, D'Ambrosio C, Baserga R. The IGF-1 receptor in mitogenesis and transformation of mouse embryo cells; role of receptor number. *Exp Cell Res* 1997;230:284–92.
- Reiss K, Valentinis B, Tu X, Xu SQ, Baserga R. Molecular markers of IGF-I mediated mitogenesis. *Exp Cell Res* 1998;242:361–72.
- Butler AA, Blakesley VA, Tsokos M, Pouliki V, Wood TL, LeRoith D. Stimulation of tumor growth by recombinant human insulin-like growth factor 1 (IGF-I) is dependent on the dose and the level of IGF-I receptor expression. *Cancer Res* 1998;58:3021–7.
- Sell C, Dumenil G, Deveaud C, et al. Effect of a null mutation of the type I IGF receptor gene on growth and transformation of mouse embryo fibroblasts. *Mol Cell Biol* 1994;14:3604.
- Rodriguez-Tarduchy G, Collins MK, Garcia I, Lopez-Rivas A. Insulin-like growth factor I inhibits apoptosis in IL3 dependent hemopoietic cells. *J Immunol* 1992;149:535–40.
- Wu Y, Tewari M, Cui S, Rubin R. Activation of the insulin-like growth factor I receptor inhibits tumor necrosis factor induced cell death. *J Cell Physiol* 1996;168:499–509.
- Sell C, Baserga R, Rubin R. Insulin-like growth factor I (IGF-I) and the IGF-I receptor prevent etoposide induced apoptosis. *Cancer Res* 1995;55:303–6.
- Kulik G, Klippel A, Weber MJ. Anti-apoptotic signaling by the insulin-like growth factor I receptor, phosphatidylinositol 3-kinase and Akt. *Mol Cell Biol* 1997;17:1595–606.
- Singleton JR, Dixit VM, Feldman EL. Type I insulin-like growth factor receptor activation regulates apoptotic proteins. *J Biol Chem* 1996;271:31791–4.
- Chan JM, Stampfer MJ, Giovannucci E, et al. Plasma insulin-like growth factor-I and prostate cancer risk: a prospective study. *Science* 1998;279:563–6.
- Ma J, Pollak M, Giovannucci E, et al. Prospective study of colorectal cancer risk in men and plasma levels of insulin-like growth factor (IGF-I) and IGF-binding protein-3. *J Natl Cancer Inst* 1999;91:620–5.
- Vadgama JV, Wu Y, Datta G, Khan H, Chillar R. Plasma insulin-like growth factor-I and serum IGF-binding protein 3 can be associated with the progression of breast cancer, and predict the risk of recurrence and the probability of survival in African-American and Hispanic women. *Oncology* 1999;57:330–40.
- Decensi A, Veronesi U, Miceli R, et al. Relationships between plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 and second breast cancer risk in a prevention trial of Fenretinide. *Clin Cancer Res* 2003;9:4722–9.
- Zhang L, Zhou W, Velculescu VE, et al. Gene expression profiles in normal and cancer cells. *Science* 1997;276:1268–72.
- Sohda T, Iwata K, Soejima H, Kamimura S, Shijo H, Yun K. *In situ* detection of insulin-like growth factor II (IGF2) and H19 gene expression in hepatocellular carcinoma. *J Hum Genet* 1998;43:49–53.
- Takada S, Kondo M, Kumada T, et al. Allelic-expression imbalance of the insulin-like growth factor 2 gene in hepatocellular carcinoma and underlying disease. *Oncogene* 1996;12:1589–92.
- Boulle N, Logie A, Gicquel C, Perin L, Le Bouc Y. Increased levels of insulin-like growth factor II (IGF-II) and IGF-binding protein 2 are associated with malignancy in sporadic adrenocortical tumors. *J Clin Endocrinol Metab* 1998;83:1713–20.
- Martin DM, Singleton JR, Meghani MA, Feldman EL. IGF receptor function and regulation in autocrine human neuroblastoma cell growth. *Regul Pept* 1993;48:225–32.
- Reeve AE, Eccles MR, Wilkins RJ, Bell GI, Millow LJ. Expression of insulin-like growth factor-II transcripts in Wilms' tumour. *Nature* 1985;317:258–62.
- Resnik JL, Reichart DB, Huey K, Webster NJG, Seely BL. Elevated Insulin-like growth factor I receptor autophosphorylation and kinase activity in human breast cancer. *Cancer Res* 1998;58:1159–64.
- Pekonen F, Partanen S, Makinen T, Rutanen E. Receptor for epidermal growth factor and insulin-like growth factor I and their relation to steroid receptors in human breast cancer. *Cancer Res* 1988;48:1343–7.
- Steller MA, Delgado CH, Bartels CJ, Woodworth CD, Zou Z. Overexpression of the insulin-like growth factor 1 receptor and autocrine stimulation in human cervical cancer cells. *Cancer Res* 1996;15:1761–5.
- Werner H, Re GG, Drummond IA, et al. Increased expression of the insulin-like growth factor I receptor gene, IGF1R, in Wilms tumor is correlated with modulation of IGF1R promoter activity by the WT1 Wilms tumor gene product. *Proc Natl Acad Sci U S A* 1993;90:5828–32.
- Kaiser U, Schardt C, Brandscheidt D, Wollmer E, Havemann K. Expression of insulin-like growth factor receptors I and II in normal human lung and in lung cancer. *J Cancer Res Clin Oncol* 1993;119:665.
- Parker AS, Cheville JC, Janney CA, Cerhan JR. High expression levels of insulin-like growth factor-I receptor predict poor survival among women with clear-cell renal cell carcinoma. *Hum Pathol* 2002;33:801–5.
- Burfeind P, Chernicky CL, Rininsland P, Ilan J. Antisense RNA to the type I insulin-like growth factor receptor suppresses tumor growth and prevents invasion by rat prostate cancer cells *in vivo*. *Proc Natl Acad Sci U S A* 1996;93:7263–8.
- Neuenschwander S, Roberts CT, Leroith D. Growth inhibition of MCF7 breast cancer cells by stable expression of an insulin-like growth factor I receptor antisense ribonucleic acid. *Endocrinology* 1995;136:4298–303.

29. Reiss K, D'Ambrosio C, Tu X, Tu C, Baserga R. Inhibition of tumor growth by a dominant negative mutant of the insulin-like growth factor I receptor with a bystander effect. *Clin Cancer Res* 1998;4:2647–55.
30. D'Ambrosio C, Ferber A, Resnicoff M, Baserga RA. Soluble insulin-like growth factor I receptor that induces apoptosis of tumor cells *in vivo* and inhibits tumorigenesis. *Cancer Res* 1996;56:4013–20.
31. Hailey J, Maxwell E, Koukouras K, Bishop WR, Pachtter JA, Wang Y. Neutralizing anti-insulin-like growth factor receptor 1 antibodies inhibit receptor function and induce receptor degradation in tumor cells. *Mol Cancer Ther* 2002;1:1349–53.
32. Arteaga CL, Kitten LJ, Coronado EB, et al. CK. Blockade of the type I somatomedin receptor inhibits growth of human breast cancer cells in athymic mice. *J Clin Invest* 1989;84:1418–23.
33. Gansler T, Furlanetto R, Grambling TS, et al. Antibody to type 1 insulin-like growth factor inhibits growth of Wilm's tumor in culture and in athymic mice. *Am J Pathol* 1989;135:961–6.
34. Prager D, Li H, Asa S, Melmed S. Dominant negative inhibition of tumorigenesis *in vivo* by human insulin-like growth factor 1 receptor mutant. *Proc Natl Acad Sci U S A* 1994;91:2181–5.
35. Liu X, Turbyville T, Fritz A, Whitesell L. Inhibition of insulin-like growth factor 1 receptor expression in neuroblastoma cells induces the regression of established tumors in mice. *Cancer Res* 1998;58:5432–8.
36. Maloney EK, McLaughlin JL, Dagdigian NE, et al. An anti-insulin-like growth factor receptor antibody that is a potent inhibitor of cancer cell proliferation. *Cancer Res* 2003;63:5073–83.
37. Burtrum D, Zhu Z, Lu D, et al. A fully human monoclonal antibody to the insulin-like growth factor I receptor blocks ligand-dependent signaling and inhibits human tumor growth *in vivo*. *Cancer Res* 2003;63:8912–21.
38. Goetsch L, Gonzalez A, Leger O, et al. A recombinant humanized anti-insulin-like growth factor receptor type I antibody (h7C10) enhances the antitumor activity of vinorelbine and anti-epidermal growth factor receptor therapy against human cancer xenografts. *Int J Cancer* 2005;113:316–28.
39. Jackson-Booth P-G, Terry C, Lackey B, Lopaczynska M, Nissley P. Inhibition of the biologic response to insulin-like growth factor I in MCF-7 breast cancer cells by a new monoclonal antibody to the insulin-like growth factor-I receptor. The importance of receptor down-regulation. *Horm Metab Res* 2003;35:850–6, 2003.
40. Sachdev D, Li S, Hartell JS, Fujita-Yamaguchi Y, Miller JS, Yee D. A chimeric humanized single-chain antibody against the type I insulin-like growth factor (IGF) receptor renders breast cancer cells refractory to the mitogenic effects of IGF-I. *Cancer Res* 2003;63:627–35.
41. Lonberg N, Taylor L, Harding F, et al. Antigen-specific human antibodies from mice comprising four distinct genetic modifications. *Nature* 1994;368:856–9.
42. LeRoith D, Werner H, Beitner-Johnson D, Roberts CT Jr. Molecular and cellular aspects of the insulin-like growth factor I receptor. *Endocr Rev* 1995;16:143–63.
43. De Meyts P, Whittaker J. Structural Biology of insulin and IGF-I receptors: implications for drug design. *Nat Rev* 2002;1:769–83.
44. Siddle K, Soos MA, Field CE, Nave BT. Hybrid and atypical insulin/insulin-like growth factor I receptors. *Horm Res* 1994;41:56–65.
45. Corsello SM, Rota CA, Putignano P, et al. Effect of acute and chronic administration of tamoxifen on GH response to GHRH and on IGF-I serum levels in women with breast cancer. *Eur J Endocrinol* 1998;139:309–13.
46. Nielsen TO, Andrews HN, Cheang M, et al. Expression of the insulin-like growth factor I receptor and urokinase plasminogen activator in breast cancer is associated with poor survival: potential for intervention with 17-allylamino geldanamycin. *Cancer Res* 2004;64:286–91.
47. Tang Y, Zhang D, Fallavollita L, Brodt P. Vascular endothelial growth factor C expression and lymph node metastasis are regulated by the type I insulin-like growth factor receptor. *Cancer Res* 2003;63:1166–71.
48. Parrizas M, Gazit A, Levitzki A, Wertheimer E, LeRoith D. Specific inhibition of insulin-like growth factor-1 and insulin receptor tyrosine kinase activity and biological function by typhostins. *Endocrinology* 1997;138:1427.
49. Blum G, Gazit A, Levitzki A. Substrate competitive inhibitors of IGF-1 receptor kinase. *Biochemistry* 2000;39:15705.
50. Blum G, Gazit A, Levitzki A. Development of new insulin-like growth factor-1 receptor kinase inhibitors using catechol mimics. *J Biol Chem* 2003;278:40442.
51. Garcia-Echeverria C, Pearson M, Marti A, et al. *In vivo* anti-tumor activity of NVP-AEW541—a novel, potent and selective inhibitor of the IGF-1R kinase. *Cancer Cell* 2004;5:231–9.
52. Mitsiades CS, Mitsiades NS, McMullan, et al. Inhibition of the insulin-like growth factor receptor 1 tyrosine kinase activity as a therapeutic strategy for multiple myeloma, other hematologic malignancies, and solid tumors. *Cancer Cell* 2004;5:221–30.
53. Lu Y, Zi X, Zhao Y, Pollak M. Insulin-like growth factor-I receptor signaling and resistance to trastuzumab (Herceptin). *J Natl Cancer Inst* 2001;93:1852–7.
54. Chakravarti A, Loeffler JS, Dyson NJ. Insulin-like growth factor receptor I mediates resistance to anti-epidermal growth factor therapy in primary human glioblastoma cells through continued activation of phosphoinositide 3-kinase signaling. *Cancer Res* 2002;62:200–7.

# Molecular Cancer Therapeutics

## Inhibition of insulin-like growth factor-I receptor (IGF-IR) signaling and tumor cell growth by a fully human neutralizing anti-IGF-IR antibody

Yan Wang, Judith Hailey, Denise Williams, et al.

*Mol Cancer Ther* 2005;4:1214-1221.

**Updated version** Access the most recent version of this article at:  
<http://mct.aacrjournals.org/content/4/8/1214>

**Cited articles** This article cites 51 articles, 26 of which you can access for free at:  
<http://mct.aacrjournals.org/content/4/8/1214.full#ref-list-1>

**Citing articles** This article has been cited by 26 HighWire-hosted articles. Access the articles at:  
<http://mct.aacrjournals.org/content/4/8/1214.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://mct.aacrjournals.org/content/4/8/1214>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.