

The antiproliferative agent MLN944 preferentially inhibits transcription

Sarah A. Byers,¹ Blanca Schafer,²
Darshan S. Sappal,³ Jeffery Brown,³
and David H. Price²

¹Molecular Biology Program and ²Department of Biochemistry, University of Iowa, Iowa City, Iowa and ³Department of Molecular and Cellular Oncology, Millennium Pharmaceuticals, Inc., Cambridge, Massachusetts

Abstract

MLN944 is a novel compound currently being codeveloped by Millennium Pharmaceuticals and Xenova Ltd. as a cancer therapeutic and is in a phase I clinical trial for solid tumors. Although MLN944 was originally proposed to function as a topoisomerase I and II inhibitor, more recent data has shown that it is a DNA-intercalating agent that does not inhibit the catalytic activity of topoisomerase I or II. We show here that MLN944 inhibits incorporation of radiolabeled precursors into RNA preferentially over incorporation into DNA and protein in HCT116 and H460 cells. To determine if MLN944 inhibits transcription, a human RNA polymerase II *in vitro* transcription system was used. MLN944 inhibited initiation when added before or after the formation of preinitiation complexes and inhibited elongation at higher concentrations. The preferential inhibition of initiation differentiates MLN944 from actinomycin D, which more strongly inhibits elongation. Transcription of all RNA polymerases was inhibited in nuclei isolated from HeLa cells treated with low concentrations of MLN944. Our data are consistent with transcription as the target of the potent cytotoxic effects of MLN944. [Mol Cancer Ther 2005;4(8):1260–7]

Introduction

MLN944 is a novel bis-phenazine compound that is currently being tested for anticancer activity in phase I clinical trials for solid tumors. MLN944 was developed in a drug design program intended to create topoisomerase inhibitors (1). Potent cytotoxicity of MLN944 has been seen in Jurkat and Chinese hamster ovary cell lines and has

translated to human colon carcinoma xenograft models in mice (2). Recently, MLN944 was shown to be significantly more potent than doxorubicin, topotecan and paclitaxel against a variety of human tumors *ex vivo* (3). MLN944 has also been shown to have an additive potency when added sequentially with 5-fluorouracil and irinotecan to a human colon cancer cell line or xenografts (4).

The mechanism of action of MLN944 cytotoxicity is currently under investigation. MLN944 was originally reported to function as a topoisomerase I and II inhibitor due to its ability to stimulate the isolation of cleavable complexes containing topoisomerase I or II (2). Subsequently, MLN944 was found not to inhibit topoisomerase I or II catalytic activity in DNA supercoiling assays (5). Compared with the topoisomerase I and II inhibitors, camptothecin and etoposide, MLN944 only slightly stimulated the formation of nicked or linear DNA species in *in vitro* DNA cleavage assays (5). The global cellular effect of MLN944 treatment is cell cycle arrest in G₁/G₂, which is different from camptothecin and etoposide that cause a G₂-M arrest (5). Treatment of human colon carcinoma xenografts with MLN944 or the topoisomerase inhibitor CPT-11 were found to modify the expression of different subsets of genes in microarrays (5). These data strongly suggest that the mechanism of action of MLN944 is not through inhibition of either topoisomerase I or II.

Recently, the nuclear magnetic resonance structure of MLN944 binding the B-DNA oligonucleotide, d(ATG-CAT)₂, was determined (6). The two phenazine rings of MLN944 were found to bis-intercalate between T2pG3 and C4pA5 and the carboxamide amino linker interacted with the central G3:C4 base pairs in the major groove of the DNA, causing an unwinding of the DNA at the intercalation sites, resulting in a more shallow major groove and a broader minor groove (6). Electrophoretic shift mobility assays examining the effect of MLN944 on activator protein and nuclear factor- κ B found that the binding of c-Jun to the activator protein binding site was inhibited by MLN944, but nuclear factor- κ B binding to its recognition sequence was not affected (6).

DNA and the processes associated with it are attractive targets for cancer therapies either directly or as the end result of signaling pathways (7). DNA-intercalating agents have inhibitory effects on transcription initiation or elongation. Actinomycin D, a monointercalator that recognizes the sequence d(pGpC), inhibits RNA synthesis (8, 9) and more specifically the elongating polymerase (9). Daunomycin, a monointercalating agent and its bis-intercalating derivative, WP631, inhibit transcription (10). WP631 also inhibits SP-1-activated transcription at lower concentrations (10). Actinomycin D has been used to treat many cancers and daunomycin is used to treat leukemia. The DNA cross-linker cisplatin is also used to treat many

Received 4/7/05; revised 6/7/05; accepted 6/14/05.

Grant support: Millennium Pharmaceuticals, Inc. and NIH grant GM35500 (D.H. Price).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: David H. Price, Department of Biochemistry, University of Iowa, 3130 MERF, Iowa City, IA 52242. Phone: 319-335-7910; Fax: 319-384-4770. E-mail: david-price@uiowa.edu

Copyright © 2005 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-05-0109

tumors, especially those of testicular and ovarian origin (11). Cisplatin bends the DNA, inhibiting transcription elongation by blocking the ability of RNA polymerase II to traverse the DNA (12, 13). Production of full-length mRNAs is also blocked by compounds, such as flavopiridol, that inhibit the kinase activity of positive transcription elongation factor b (P-TEFb; refs. 14, 15). P-TEFb phosphorylates the CTD of the largest subunit of RNA polymerase II, allowing for a transition from abortive to productive elongation (16, 17). Flavopiridol has completed phase I and II clinical trials for various cancers (18).

The DNA intercalating properties of MLN944 make transcription a strong candidate for its target. The ability of MLN944 to inhibit transcription was tested *in vivo* by examining the incorporation of radiolabeled precursors into RNA and DNA and with nuclear run-ons. The effects of MLN944 on initiation and elongation was examined separately in *in vitro* transcription assays and compared with another bis-intercalator, actinomycin D.

Materials and Methods

Compounds

MLN944 dimesylate salt was synthesized at Auckland Cancer Society Research Centre, University of Auckland, Auckland, New Zealand (2). DMSO stocks of MLN944 of 5 or 10 mmol/L were prepared and stored at -80°C . Actinomycin D was purchased from Sigma-Aldrich (St. Louis, MO) and 1 mmol/L stocks in DMSO or 10 mmol/L stocks in methanol were prepared. 5,6-Dichloro-1- β -D-ribofuranosylbenzimidazole (DRB) was purchased from Sigma-Aldrich and a 10 mmol/L stock in ethanol was prepared.

Trichloroacetic Acid Precipitation Assays

HCT116 and H460 cells (American Type Culture Collection, Manassas, VA) were plated overnight at a density of 5×10^5 per well in a 6-well dish. HCT116 cells were maintained in McCoy's 5A media (Life Technologies, Gaithersburg, MD) supplemented with 10% fetal bovine serum (Hyclone, Logan, UT) and 1% L-Glutamine (Life Technologies). H460 cells were maintained in RPMI 1640 (Life Technologies) media supplemented with 10% fetal bovine serum and 1% L-Glutamine. Cells were cultured under standard conditions (37°C in 5% CO_2). Cells were treated with MLN944 or actinomycin D for 1 hour before labeling while maintaining DMSO below 2% of the final volume. Cells were then pulse labeled for 1 hour with [5- ^3H] uridine, [Methyl- ^3H] thymidine, or L-[U- ^{14}C] leucine (Amersham, Arlington Heights, IL) to measure RNA, DNA, and protein synthesis, respectively. The percent incorporation was measured by precipitating each sample in 5 volumes of 20% trichloroacetic acid (Perkin-Elmer, Norwalk, CT) for 30 minutes at 40°C and recovered by filtration onto glass fiber filters (VWR, Chester, PA). Filters were washed twice with 5 mL of 5% trichloroacetic acid followed by a single wash with 5 mL of 95% ethanol, air dried, and counted in 5 mL Optiphase Super Mix scintillation fluid (Perkin-Elmer) on a scintillation counter (Beckman Coulter LS 6500).

Transcription Assays

The RNA polymerase II *in vitro* transcription assays were done as described previously (14, 19). Transcription reactions were carried out by incubating 20 ng/ μL template DNA, a PCR product containing the cytomegalovirus (CMV) promoter, in the presence of HeLa nuclear extract (20, 21) under standard transcription conditions of 60 mmol/L KCl, 20 mmol/L HEPES, and 7 mmol/L MgCl_2 for 10 minutes. Reactions contained 50 $\mu\text{mol/L}$ DRB (Sigma, St. Louis, MO), or indicated amounts of MLN944 while maintaining a final concentration of 0.5% DMSO. Transcription was initiated by the addition of 0.5 mmol/L ATP, CTP, and GTP and 10 μCi α - ^{32}P -UTP (ICN). After 45 seconds, the UTP concentration was raised to 1 mmol/L UTP to generate run-off transcripts. Transcription was stopped by the addition of Sarkosyl stop solution [1% Sarkosyl, 100 mmol/L Tris (pH 7.5), 100 mmol/L NaCl, 10 mmol/L EDTA, and 100 $\mu\text{g/mL}$ tRNA]. The RNA was isolated and resolved on a 6% urea gel. Quantitation was done using an InstantImager (Packard, Meriden, CT) and data was normalized to the DMSO control and curve fit with Table Curve (Jandel Scientific, San Rafael, CA) to determine the IC_{50} . Transcription initiation assays were done in a similar manner, except 45 seconds after the addition of 0.5 mmol/L ATP, CTP, and GTP and 10 μCi α - ^{32}P -UTP (ICN) transcription was stopped by the addition of the Sarkosyl stop solution. The initiation assays were also done with 30 ng/ μL Bgl II-linearized C<6> plasmid (22) DNA containing the SV40 promoter as the template DNA. The amounts of MLN944 or actinomycin D and order of addition are as indicated in the figure legends.

Kinase Assay

The *in vitro* kinase assays were done as previously described (17) except that recombinant P-TEFb, composed of human Cdk9 and cyclin T2a (23) was used, and the substrate was recombinant human DSIF (21). Kinase reactions were carried out with 2.5 μCi α - ^{32}P -ATP (Amersham), 30 $\mu\text{mol/L}$ ATP, 20 mmol/L HEPES, 7 mmol/L MgCl_2 , and 30 mmol/L KCl for 20 minutes at 30°C . The kinase reactions were stopped by the addition of SDS loading buffer and resolved on a 7.5% SDS polyacrylamide gel. Label incorporation into the largest subunit of DSIF was quantitated with an InstantImager (Packard).

Immobilized Template Transcription Assay

The immobilized template *in vitro* transcription assays were done as outlined previously (14, 24). Transcription reactions were carried out in the presence of HeLa nuclear extract at 60 mmol/L KCl, 20 mmol/L HEPES, 7 mmol/L MgCl_2 , and 50 $\mu\text{mol/L}$ DRB for 10 minutes using a biotinylated template DNA that had been conjugated to streptavidin paramagnetic beads. Transcription was initiated by the addition of 0.5 mmol/L ATP, CTP, and GTP and 10 μCi α - ^{32}P -UTP for 45 seconds. Transcription was stopped by addition of EDTA to a final concentration of 30 mmol/L to produce early elongation complexes that were isolated with a magnetic concentrator. The early elongation complexes were washed twice with 1 mol/L KCl and 1% Sarkosyl to remove all factors except for the engaged

polymerase, and equilibrated by washing once with transcription buffer (60 mmol/L KCl, 20 mmol/L HEPES, 7 mmol/L MgCl₂, and 0.2 µg/mL bovine serum albumin) before suspension in transcription buffer. The isolated early elongation complexes were then incubated with increasing amounts of MLN944 or actinomycin D for 10 minutes and then chased for 5 minutes in transcription buffer containing nucleotide triphosphates. The RNA was isolated and resolved on a 6% UREA gel and subjected to autoradiography.

Nuclear Run-ons

HeLa S3 cells were cultured in suspension under standard conditions to a concentration of 3.5 to 4 × 10⁵ cells/mL in DMEM/F-12 media (Invitrogen, San Diego, CA) supplemented with 10% fetal bovine serum (Hyclone) and 1 × penicillin/streptomycin (Invitrogen). Cells were treated with 0, 30, 100, or 300 nmol/L MLN944 for 1 hour. Nuclei were isolated by isotonic homogenization of the cells in the presence of 0.1% Triton X-100 and spinning them through a 1.9 mol/L sucrose cushion (25). The nuclei were stored at -80°C in storage buffer [25% glycerol, 10 mmol/L Tris (pH 7.5), 5 mmol/L Mg(Ac)₂, and 1 mmol/L DTT]. Nuclear run-on experiments were carried out in 18-µL reactions containing 12.5% glycerol, 120 mmol/L KCl, 7.5 mmol/L Mg(Ac)₂, 15 mmol/L Tris (pH 7.5), 0.5 mmol/L DTT, 0.5 mmol/L A/G/CTP, 5 µmol/L UTP, 20 µCi α-³²P-UTP, ± 2 µg/mL α-amanitin (Sigma), and 20 mmol/L HEPES. Reactions were incubated at room temperature for 0, 10, or 20 minutes and stopped with 82 µL of Sarkosyl stop solution, spotted on Whatman DE81 filters and allowed to air dry. Filters were washed four times for 5 minutes with at least 10 mL of 5% K₂HPO₄ and 0.3% Na₄PO₇ per filter, or until no more counts were detected in the wash, and dried under a strong lamp. Counts were analyzed with an InstantImager and by scintillation counting.

Results

MLN944 Inhibits RNA Synthesis

To begin to understand the mechanism of cellular toxicity of MLN944, its effect on RNA, DNA, and protein synthesis was evaluated in two human tumor cell lines. H460 and HCT116 cells were treated with increasing amounts of MLN944 for 1 hour and pulsed with radiolabeled precursors for 1 hour followed by trichloroacetic acid precipitation to monitor their incorporation. The transcription of new RNA, as indicated by [5-6, ³H] uridine incorporation, was inhibited with IC₅₀s of 0.3 and 3 nmol/L for H460 (Fig. 1A) and HCT116 (Fig. 1B) cells, respectively. DNA synthesis, as indicated by [methyl-³H] thymidine incorporation, was inhibited with an IC₅₀ of 300 nmol/L in both cell lines (Fig. 1A and B). Protein synthesis, as indicated by L-[U-¹⁴C] leucine incorporation, remained unaffected in both cell lines at all concentrations of MLN944 that were tested (Fig. 1A and B). The results indicate that the primary target of MLN944 is transcription, because inhibition of RNA synthesis occurs at two to three orders of magnitude lower concentrations than inhibition of DNA synthesis. The time course of this experiment is

such that protein synthesis should be unaffected by inhibition of transcription. The inhibition of RNA synthesis also occurs in the same concentration range in which the potent cytotoxicity of MLN944 was seen previously in Jurkat and Chinese hamster ovary cells lines, with IC₅₀s in the range of 0.04 to 0.4 nmol/L (2).

MLN944 Inhibits RNA Polymerase II Transcription *In vitro*

To develop an understanding of how MLN944 inhibits RNA synthesis, its effect on transcription by RNA polymerase II was examined in an *in vitro* system. Transcription was carried out using HeLa nuclear extract and a template containing the CMV promoter. The incorporation of a radiolabeled nucleotide into a 600-nucleotide run-off transcript was used to detect the level of transcription. As expected, inhibition of P-TEFb by the inclusion of 50 µmol/L DRB (16, 26) reduced the amount of run-off compared with the control reaction (Fig. 2A). Addition of increasing amounts of MLN944 (0.01–10 µmol/L) caused a decrease in the amount of run-off transcripts starting at 0.3 µmol/L, with a complete loss of run-off transcripts by 3 µmol/L (Fig. 2A). The amount of run-off was quantitated (Fig. 2B) and an IC₅₀ of 0.6 µmol/L




Figure 1. MLN944 inhibits RNA synthesis. RNA, DNA, and protein synthesis was examined in H460 (A) and HCT116 (B) cells treated with the indicated amounts of MLN944 for 1 h followed by pulse labeling with [5-6, ³H] uridine, [methyl-³H] thymidine, or L-[U-¹⁴C] leucine (for RNA, DNA, and protein synthesis, respectively) for 1 h. Trichloroacetic acid precipitation was done and the incorporation of label was quantitated with a scintillation counter.




Figure 2. MLN944 inhibits RNA polymerase II transcription *in vitro*. **A**, RNA polymerase II transcription using HNE and a template containing the CMV promoter was measured by generation of 600-nucleotide run-off transcripts in an *in vitro* transcription assay. Reactions contained a constant amount of DMSO to normalize for any slight effects it may have on transcription and were carried out with DMSO alone, 50 μmol/L DRB, or the indicated amounts of MLN944. The RNA was isolated, resolved on a 6% urea gel, quantitated with an InstantImager, and subjected to autoradiography. **B**, the data from three independent experiments was normalized to the run-off transcripts produced in the DMSO control and plotted.

was calculated. These results indicate that MLN944 inhibits transcription by RNA polymerase II but do not differentiate between inhibition of initiation and an effect on elongation.

MLN944 Does Not Inhibit P-TEFb

To determine if the effect of MLN944 on transcription was due to the inhibition of the elongation control factor P-TEFb, an *in vitro* kinase assay was done. The incorporation of labeled phosphate from γ -³²P-ATP into the largest subunit of DSIF (27) was examined in the presence of increasing amounts of MLN944 (0.01–10 μmol/L). No inhibition was seen at any concentration tested (Fig. 3). We conclude that the effects of MLN944 in the *in vitro* transcription assay (Fig. 2) are not due to inhibition of P-TEFb.

MLN944 Inhibits RNA Polymerase II Elongation

To determine if MLN944 affects elongation by inhibiting RNA polymerase II directly, an *in vitro* transcription assay using an immobilized template was done. Transcription was initiated using HeLa nuclear extract and a CMV template conjugated to paramagnetic beads. Early elongation complexes containing RNA polymerase II and short transcripts were isolated by washing with 1 mol/L KCl and 1% Sarkosyl to strip away all factors. Increasing amounts of MLN944 (0.3–10 μmol/L) were added to the early elongation complexes before the complexes were chased for 5 minutes in the presence of 0.5 mmol/L nucleotide triphosphates. The first effects on elongation were seen at 1 μmol/L as indicated by the generation of slightly shorter transcripts compared with the control reaction (Fig. 4). Elongation was further hindered, as the amount of

MLN944 was increased (Fig. 4). Clearly, MLN944 inhibited transcription elongation by RNA polymerase II directly; however, the concentration of MLN944 needed to inhibit elongation was significantly greater than that needed to inhibit the generation of run-off transcripts (Fig. 2A). This suggests that the effects of MLN944 on elongation may be secondary to its effects on initiation.

MLN944 Inhibits Initiation

To examine the ability of MLN944 to inhibit initiation by RNA polymerase II, short pulse labeling reactions were used. MLN944 has been shown to block the binding of a transcription factor, c-Jun (6). Because MLN944 might be more effective at blocking the binding of the basal transcription factors than inhibiting initiation of preformed preinitiation complexes, the compound was added to reactions before, at the same time, or after formation of preinitiation complexes. Transcription was initiated in the presence of a limiting radiolabeled nucleotide during a 45-second pulse that predominately produced a 15-nucleotide transcript (Fig. 5). As expected the P-TEFb inhibitor, DRB, had no effect on initiation compared with the control under all three conditions tested (Fig. 5). Increasing amounts of MLN944 (0.03–10 μmol/L) caused a decrease in the amount transcripts produced regardless of when the compound was added (Fig. 5). The results were quantitated (Fig. 5E) and transcription initiation from the CMV promoter under all three conditions was inhibited IC₅₀s of ~0.4 μmol/L. This is similar to 0.6 μmol/L calculated earlier for inhibition of the generation of run-off transcripts from the CMV promoter (Fig. 2). Inhibition of initiation was seen at concentrations of MLN944 that had no effect on elongation in the immobilized template transcription assay (Fig. 4), indicating that MLN944 inhibits RNA polymerase II transcription initiation from the CMV promoter before elongation is affected.




Figure 3. MLN944 is not a P-TEFb inhibitor. **A**, a P-TEFb *in vitro* kinase assay with DSIF as the substrate was done in the presence of the indicated amounts of MLN944. The reactions were resolved on a 7.5% SDS-PAGE gel and subjected to autoradiography. **B**, incorporation of γ -³²P-ATP into the largest subunit of DSIF, in three independent experiments, was quantitated with an InstantImager, normalized to the DMSO control, and plotted.




Figure 4. MLN944 inhibits elongation by RNA polymerase II. Isolated early elongation complexes (EEC) obtained from a pulse (P) labeling were incubated with the indicated amounts of MLN944 for 10 min and allowed to extend transcripts for 5 min. Markers (M), 25-bp ladder.

To determine if the effect of MLN944 on initiation was promoter specific, transcription from another template containing the SV40 promoter was analyzed. MLN944 was added after the formation of preinitiation complexes and initiation was allowed to occur during a 45-second pulse. Initiation was strongly inhibited between 1 and 3 $\mu\text{mol/L}$ MLN944 (Fig. 5D). Addition of MLN944 before formation of preinitiation complexes on the SV40 promoter exhibited an identical dose response (data not shown). Transcription initiation from the SV40 promoter was quantitated and an IC_{50} of $\sim 0.9 \mu\text{mol/L}$ was calculated. The reaction carried out in the presence of 0.1 $\mu\text{mol/L}$ MLN944 was excluded from this calculation, due to the poor recovery of the transcripts. The IC_{50} of $\sim 0.9 \mu\text{mol/L}$ is about 2-fold higher than that found when a CMV template was used. For this experiment, the SV40 template was a restriction digested plasmid rather than a PCR product as was used when the CMV promoter was tested. To compensate for the lower SV40 promoter concentration, more template DNA was added into SV40 transcription reactions and because of this more MLN944 was needed to inhibit transcription to the same level. Supporting this idea, we found that twice as much MLN944 was required to achieve the same level of inhibition of initiation from the CMV promoter, when the total amount of DNA was doubled by addition of plasmid DNA lacking an RNA polymerase II promoter (data not shown). These data strongly suggest that inhibition of initiation by MLN944 is not promoter specific.

Mechanism of Action of MLN944 Is Novel

The effects of MLN944 on transcription initiation and elongation were compared with actinomycin D, a mono-intercalating agent that inhibits RNA polymerase II (8). The generation of short transcripts was examined with increasing amounts of actinomycin D (0.3–10 $\mu\text{mol/L}$ actinomycin D) added to the template before or after formation of preinitiation complexes. When the compound was added before the formation of preinitiation complexes, inhibition of initiation was first seen at 3 $\mu\text{mol/L}$ and was complete by 10 $\mu\text{mol/L}$ (Fig. 6A). When it was added after the formation of preinitiation complexes similar, but not identical results were obtained (Fig. 6B). Both experiments were quantitated (Fig. 6C), and IC_{50} s were calculated for addition of actinomycin D before (1.6 $\mu\text{mol/L}$) or after (2.5 $\mu\text{mol/L}$) preinitiation complex formation. Actinomycin D was more inhibitory when added to the template before formation of preinitiation complexes, in contrast to MLN944 which inhibited equally well whether it was added before, at the same time, or after preinitiation




Figure 5. MLN944 inhibits RNA polymerase II transcription initiation. RNA polymerase II transcription initiation was examined by measuring the generation of short transcripts from the CMV promoter. Reactions were carried out in the presence of 50 $\mu\text{mol/L}$ DRB or the indicated amounts of MLN944 added either before (A), during (B), or after (C), formation of preinitiation complexes. D, transcription initiation from the SV40 promoter was also examined in the presence of the indicated amounts of MLN944. The RNA was resolved on a 6% urea gel. E, short transcripts were quantitated and plotted.

complex formation. The effect of actinomycin D on RNA polymerase II elongation was examined using an immobilized template assay. Isolated early elongation complexes were incubated for 10 minutes with increasing amounts of actinomycin D (0.01–30 $\mu\text{mol/L}$) and then chased for 5 minutes. The first effects of actinomycin D on transcript length were seen at 0.3 $\mu\text{mol/L}$ and became more dramatic as the concentration was increased (Fig. 6D). Actinomycin D inhibited initiation with an IC_{50} of 1.6 to 2.5 $\mu\text{mol/L}$ and began to inhibit elongation at 0.3 $\mu\text{mol/L}$ compared with MLN944 that inhibited initiation with an IC_{50} of 0.4 $\mu\text{mol/L}$ and began to inhibit elongation at 1 $\mu\text{mol/L}$. Overall, the data indicate that MLN944 preferentially inhibits initiation and that actinomycin D preferentially inhibits elongation.

MLN944 Inhibits Transcription *In vivo*

The ability of MLN944 to inhibit transcription *in vivo* was tested with nuclear run-on assays. Nuclei were isolated from HeLa S3 cells after undergoing a 1-hour treatment of 0, 30, 100, or 300 nmol/L MLN944. Nuclear run-ons were carried out and the amount of radiolabeled UTP incorporated after 10 or 20 minutes was quantitated. Total nuclear transcription was inhibited to below 50% after treatment with 30 nmol/L MLN944 ($\text{IC}_{50} \sim 20$ nmol/L; Fig. 7A). Nuclear run-ons were also carried out in the presence of 2 $\mu\text{g/mL}$ α -amanitin to inhibit RNA polymerase II allowing for examination of transcription by RNA polymerases I and III. Again, over half of the RNA polymerase I and III transcription was inhibited by 30 nmol/L ($\text{IC}_{50} \sim 10$ nmol/L; Fig. 7B). Transcription by RNA polymerase II was calculated by subtracting the α -amanitin insensitive counts from the total counts. A dose-dependent decrease of RNA polymerase II transcription was seen with increasing concentrations of MLN944, with approximately half of transcription inhibited at 30 nmol/L ($\text{IC}_{50} \sim 30$ nmol/L; Fig. 7C). A comparison of the 20-minute time points from run-ons looking at amanitin-resistant and amanitin-sensitive counts suggests that RNA

polymerases I and/or III are inhibited at slightly lower concentrations than RNA polymerase II (Fig. 7D). This slight difference may not be significant, but it is clear that MLN944 inhibits cellular transcription at very low concentrations, even after only a short time of treatment.

Discussion

The data presented here clearly show that the bis-intercalator MLN944 inhibits transcription and suggest that this is its primary mechanism of action. Incorporation of radiolabeled precursors into RNA was inhibited by MLN944 in H460 and HCT116 cells with IC_{50} s of 0.3 and 3 nmol/L, respectively. MLN944 inhibited both initiation and elongation by RNA polymerase II during *in vitro* transcription reactions using the CMV promoter and HeLa nuclear extract. Nuclear run-ons determined that all RNA polymerases are inhibited by a short treatment with MLN944 *in vivo*. Although MLN944 and actinomycin D are both intercalators and both inhibit transcription, differences were found in the stage of transcription preferentially affected by the two compounds.

In vitro transcription assays were used to examine the effect of MLN944 on initiation and elongation by RNA polymerase II. MLN944 inhibited initiation by RNA polymerase II with an IC_{50} of 0.4 $\mu\text{mol/L}$ and at higher concentrations directly inhibited elongation of polymerases initiated before drug addition. It is impossible to calculate an IC_{50} for an effect on elongation, but it is clear that concentrations of MLN944 that completely blocked initiation of transcription did not stop elongation. The patterns of transcripts resulting from elongation in the presence of MLN944 suggested that the drug was preferentially binding to specific sequences of the DNA because the pattern was different to that caused by intrinsic pause sites normally found in the template. The mechanism of MLN944 inhibition of initiation was not revealed by our data; however, because the drug had the same effect if added either before or after




Figure 6. Actinomycin D inhibits transcription elongation. The generation of short transcripts was examined in a transcription initiation assay. **A**, template DNA was incubated with the indicated amounts of actinomycin D for 10 min before HeLa nuclear extract was added for an additional 10-min incubation. **B**, template DNA was incubated with HeLa nuclear extract for 10 min before the indicated amounts of actinomycin D were added and incubated for another 10 min. **C**, quantitation of short transcripts. **D**, isolated elongation complexes obtained from a pulse (P) labeling were incubated with the indicated amounts of actinomycin D for 10 min and allowed to extend transcripts for 5 min. **A**, **B**, and **D**, RNA was isolated, resolved on a 6% urea gel, and subjected to autoradiography. **D**, transcript sizes are indicated.




Figure 7. MLN944 inhibits transcription *in vivo*. Nuclei were isolated from HeLa S3 cells that had been treated with increasing amounts of MLN944 (0, 30, 100, or 300 nmol/L) for 1 h. Nuclear run-ons were done, with or without 2 μ g/mL α -amanitin, and incorporation of 32 P-UTP was quantitated. Inhibition of total (A), α -amanitin insensitive (RNA polymerase I and III; B), and α -amanitin sensitive (RNA polymerases II; C) was examined. D, comparison of the effect of MLN944 on transcription by RNA polymerase II with RNA polymerases I and/or III.

preinitiation complex formation, it is not likely that formation of the preinitiation complex is affected. Perhaps the sequences or structures used by RNA polymerase II during abortive initiation, promoter escape, or very early stages of elongation are preferential binding sites for the compound. It is possible that other promoters might have different sensitivities to MLN944, but the SV40 promoter was inhibited in the same manner.

IC₅₀s determined for MLN944 inhibition of transcription *in vivo* and *in vitro* varied widely, but this can be explained by the compound's ability to intercalate DNA. As the ratio of DNA to MLN944 present in each experiment increased, the IC₅₀ also increased. The concentration of DNA in the *in vitro* transcription reactions is much higher than in the *in vivo* experiments, and higher concentrations of MLN944 are required for inhibition of transcription *in vitro*. At the IC₅₀ for inhibition of transcription *in vitro*, there is one molecule of MLN944 for every 80 bp of DNA, and at the IC₅₀ for inhibition of cellular transcription, there is one molecule of MLN944 present for every 50 bp of DNA while the cells are being treated in culture. The MLN944/DNA ratio would suggest that MLN944 is slightly more effective *in vitro*. Although MLN944 is taken up by the cells and concentrated in nuclei, as indicated by their yellow color, it is likely that not all of the compound is removed from the growth medium. Taken together, these results suggest that caution should be taken when interpreting results from different types of experiments. Because the compound is concentrated by cells during long-term treatment, it would take less MLN944 to achieve similar inhibitory effects, especially if the medium containing fresh compound was periodically replaced. This would explain why LD₅₀s were very low (0.04–0.4 nmol/L) in initial experiments with Chinese hamster ovary and Jurkat cells that received long-term treatments (2).

The effects of actinomycin D on initiation and elongation by RNA polymerase II were compared with those of MLN944. Actinomycin D had its first effect on elongation at 0.3 μ mol/L, whereas MLN944 did not have an effect until 1 μ mol/L. Comparison of the pattern of transcripts obtained with both compounds indicated that the induced pause sites were different from each other and different from the intrinsic sites on the template (compare Fig. 4 with Fig. 6D). Therefore, the effect of either compound on elongation will likely be dependent on the sequence of the template. Using the CMV promoter MLN944 was about five times more effective at blocking initiation than actinomycin D (IC₅₀s of 0.4 and 2 μ mol/L, respectively). At other promoters, the difference between the effect of MLN944 and actinomycin D might be more or less pronounced.

Our results support a model in which MLN944 could generally inhibit transcription; however, it is likely that at low concentrations the compound might alter expression of specific genes. This could occur through sequence specific binding of MLN944 leading to decreased initiation, an effect on chromatin structure, or to an effect on activation or repression of transcription by blocking interaction of regulatory molecules with their target sequences. This latter possibility is supported by the finding that MLN944 can block c-Jun binding to activator protein, but not nuclear factor- κ B binding to its target sequence *in vitro* (6). Supporting the idea that MLN944 might alter expression of a subset of genes, Sappal et al. found that in mice treated with the compound, xenographic colon tumors expressed a different pattern of genes than those in untreated mice or mice given the topoisomerase inhibitor, CPT-11 (5). In this way, MLN944 may selectively inhibit transcription of rapidly dividing tumor cells before the general effects on transcription are seen in noncancerous cells.

References

1. Gamage SA, Spicer JA, Finlay GJ, et al. Dicationic bis(9-methylphenazine-1-carboxamides): relationships between biological activity and linker chain structure for a series of potent topoisomerase targeted anticancer drugs. *J Med Chem* 2001;44:1407-15.
2. Stewart AJ, Mistry P, Dangerfield W, et al. Antitumor activity of XR5944, a novel and potent topoisomerase poison. *Anticancer Drugs* 2001;12:359-67.
3. Di Nicolantonio F, Knight LA, Whitehouse PA, et al. The *ex vivo* characterization of XR5944 (MLN944) against a panel of human clinical tumor samples. *Mol Cancer Ther* 2004;3:1631-7.
4. Harris SM, Mistry P, Freathy C, Brown JL, Charlton PA. Antitumor activity of XR5944 *in vitro* and *in vivo* in combination with 5-fluorouracil and irinotecan in colon cancer cell lines. *Br J Cancer* 2005;92:722-8.
5. Sappal DS, McClendon AK, Fleming JA, et al. Biological characterization of MLN944: a potent DNA binding agent. *Mol Cancer Ther* 2004;3:47-58.
6. Dai J, PUNCHIHEWA C, Mistry P, Ooi AT, Yang D. Novel DNA Bis-intercalation by MLN944, a potent clinical bisphenazine anticancer drug. *J Biol Chem* 2004;279:46096-103. Epub 2004 Aug 17.
7. Hurley LH. DNA and its associated processes as targets for cancer therapy. *Nat Rev Cancer* 2002;2:188-200.
8. Perry RP, Kelley DE. Inhibition of RNA synthesis by actinomycin D: characteristic dose-response of different RNA species. *J Cell Physiol* 1970;76:127-39.
9. Kimura H, Sugaya K, Cook PR. The transcription cycle of RNA polymerase II in living cells. *J Cell Biol* 2002;159:777-82.
10. Martin B, Vaquero A, Priebe W, Portugal J. Bisanthracycline WP631 inhibits basal and Sp1-activated transcription initiation *in vitro*. *Nucleic Acids Res* 1999;27:3402-9.
11. Jordan P, Carmo-Fonseca M. Molecular mechanisms involved in cisplatin cytotoxicity. *Cell Mol Life Sci* 2000;57:1229-35.
12. Mello JA, Lippard SJ, Essigmann JM. DNA adducts of *cis*-diamminedichloroplatinum(II) and its trans isomer inhibit RNA polymerase II differentially *in vivo*. *Biochemistry* 1995;34:14783-91.
13. Zlatanova J, Yaneva J, Leuba SH. Proteins that specifically recognize cisplatin-damaged DNA: a clue to anticancer activity of cisplatin. *FASEB J* 1998;12:791-9.
14. Chao SH, Fujinaga K, Marion JE, et al. Flavopiridol inhibits P-TEFb and blocks HIV-1 replication. *J Biol Chem* 2000;275:28345-8.
15. Chao SH, Price DH. Flavopiridol inactivates P-TEFb and blocks most RNA polymerase II transcription *in vivo*. *J Biol Chem* 2001;276:31793-9.
16. Marshall NF, Price DH. Purification of P-TEFb, a transcription factor required for the transition into productive elongation. *J Biol Chem* 1995;270:12335-8.
17. Marshall NF, Peng J, Xie Z, Price DH. Control of RNA polymerase II elongation potential by a novel carboxyl-terminal domain kinase. *J Biol Chem* 1996;271:27176-83.
18. Senderowicz AM. Small-molecule cyclin-dependent kinase modulators. *Oncogene* 2003;22:6609-20.
19. Marshall NF, Price DH. Control of formation of two distinct classes of RNA polymerase II elongation complexes. *Mol Cell Biol* 1992;12:2078-90.
20. Price DH, Sluder AE, Greenleaf AL. Fractionation of transcription factors for RNA polymerase II from *Drosophila* Kc cell nuclear extracts. *J Biol Chem* 1987;262:3244-55.
21. Renner DB, Yamaguchi Y, Wada T, Handa H, Price DH. A highly purified RNA polymerase II elongation control system. *J Biol Chem* 2001;276:42601-9.
22. Orozco IJ, Kim SJ, Martinson HG. The poly(A) signal, without the assistance of any downstream element, directs RNA polymerase II to pause *in vivo* and then to release stochastically from the template. *J Biol Chem* 2002;277:42899-911.
23. Peng J, Zhu Y, Milton JT, Price DH. Identification of multiple cyclin subunits of human P-TEFb. *Genes Dev* 1998;12:755-62.
24. Adamson TE, Shore SM, Price DH. Analysis of RNA polymerase II elongation *in vitro*. *Methods Enzymol* 2003;371:264-75.
25. Marzluff WF. Preparation of active nuclei. *Methods Enzymol* 1990;181:30-6.
26. Chodosh LA, Fire A, Samuels M, Sharp PA. 5,6-Dichloro-1- β -D-ribofuranosylbenzimidazole inhibits transcription elongation by RNA polymerase II *in vitro*. *J Biol Chem* 1989;264:2250-7.
27. Ping YH, Rana TM. DSIF and NELF interact with RNA polymerase II elongation complex and HIV-1 Tat stimulates P-TEFb-mediated phosphorylation of RNA polymerase II and DSIF during transcription elongation. *J Biol Chem* 2001;276:12951-8.

Molecular Cancer Therapeutics

The antiproliferative agent MLN944 preferentially inhibits transcription

Sarah A. Byers, Blanca Schafer, Darshan S. Sappal, et al.

Mol Cancer Ther 2005;4:1260-1267.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/4/8/1260>

Cited articles This article cites 27 articles, 15 of which you can access for free at:
<http://mct.aacrjournals.org/content/4/8/1260.full#ref-list-1>

Citing articles This article has been cited by 1 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/4/8/1260.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/4/8/1260>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.