

Down-regulation of Notch-1 contributes to cell growth inhibition and apoptosis in pancreatic cancer cells

Zhiwei Wang,¹ Yuxiang Zhang,² Yiwei Li,¹
Sanjeev Banerjee,¹ Joshua Liao,¹
and Fazlul H. Sarkar¹

¹Department of Pathology, Karmanos Cancer Institute, Wayne State University School of Medicine, Detroit, Michigan and
²Department of Biochemistry and Molecular Biology, Capital University of Medical Science, Beijing, China

Abstract

Pancreatic cancer remains the fourth most common cause of cancer-related death in the United States. Notch signaling plays a critical role in maintaining the balance among cell proliferation, differentiation, and apoptosis, and thereby may contribute to the development of pancreatic cancer. To characterize Notch pathway function in pancreatic cancer cells, we explored the consequences of down-regulation of Notch-1 in BxPC-3, HPAC, and PANC-1 pancreatic cancer cells. Using multiple cellular and molecular approaches such as 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay, apoptosis assay, flow cytometry, gene transfection, real-time reverse transcription-PCR (RT-PCR), Western blotting, and electrophoretic mobility shift assay for measuring DNA binding activity of nuclear factor κ B (NF- κ B), we found that down-regulation of Notch-1 inhibited cell growth and induced apoptosis in pancreatic cancer cells. Notch-1 down-regulation also increased cell population in the G₀-G₁ phase. Compared with control, small interfering RNA-transfected cells decreased expression of cyclin A, cyclin D1, and cyclin-dependent kinase 2. We found up-regulation of p21 and p27, which was correlated with the cell cycle changes. In addition, Notch-1 down-regulation also induced apoptosis, which could be due to decreased Bcl-2 and Bcl-X_L protein expression in pancreatic cancer cells. Because Notch-1 is known to cross-talk with another major cell growth and apoptotic regulatory pathway (i.e., NF- κ B),

we found that NF- κ B is a downstream target of Notch because down-regulation of Notch reduced NF- κ B activity. We also found that genistein, a prominent isoflavone, could be an active agent for the down-regulation of the Notch pathway. These findings suggest that Notch-1 down-regulation, especially by genistein, could be a novel therapeutic approach for the treatment of pancreatic cancer. [Mol Cancer Ther 2006;5(3):483–93]

Introduction

Pancreatic cancer has the worst prognosis among all major cancers and remains the fourth most common cause of cancer related death in the United States and throughout the world (1). This could be due to the fact that no effective methods of early diagnosis are currently available as well as the lack of effective therapies, resulting in high mortality of patients diagnosed with pancreatic cancer. This disappointing outcome strongly suggests that innovative research is needed to control this deadly disease.

Notch signaling is involved in cell proliferation and apoptosis, which affect the development and function of many organs (2, 3). *Notch* genes encode proteins which can be activated by interacting with a family of its ligands (4, 5). On activation, Notch is cleaved, releasing intracellular Notch which translocates into the nucleus. The intracellular Notch associates with transcriptional factors, which regulate the expression of target genes, and thus plays important roles in development and cell growth (6, 7). To date, four vertebrate *Notch* genes have been identified: *Notch-1*, *Notch-2*, *Notch-3*, and *Notch-4*. In addition, five ligands, Dll-1, Dll-3, Dll-4, Jagged-1, and Jagged-2, have been found in mammals (8, 9).

Because Notch signaling plays important roles in the cellular developmental pathway including proliferation and apoptosis (10), alterations in Notch signaling are associated with tumorigenesis. These observations suggest that dysfunction of intracellular Notch prevents differentiation, ultimately guiding undifferentiated cells toward malignant transformation (11). It has been reported that the Notch signaling network is frequently deregulated in human malignancies with up-regulated expression of Notch receptors and their ligands in cervical, lung, colon, head and neck, renal carcinoma, acute myeloid, Hodgkin and large-cell lymphomas, and pancreatic cancer (12–14). It has also been reported that Notch-1 expression inhibits apoptosis (15–17), suggesting a possible role of *Notch* as an oncogene in many cancers.

Notch-1 has been reported to cross-talk with another major cell growth and apoptotic regulatory pathway [i.e., nuclear factor κ B (NF- κ B)]. Specifically, Notch-1 has been reported to strongly induce NF- κ B promoter activity in reporter assays (18) and expression of several NF- κ B subunits (19). Notch ligands activate NF- κ B in human

Received 8/3/05; revised 10/31/05; accepted 1/5/06.

Grant support: National Cancer Institute, NIH grant 5R01CA101870-02 and The University of Texas M.D. Anderson Cancer Center subcontract award (F.H. Sarkar) through Specialized Program of Research Excellence in Pancreatic Cancer grant 1P20-CA010193-01 (J. Abbruzzese).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Fazlul H. Sarkar, Department of Pathology, Karmanos Cancer Institute, Wayne State University School of Medicine, 9374 Scott Hall, 540 East Canfield, Detroit, MI 48201. Phone: 313-576-8327; Fax: 313-576-8389. E-mail: sarkarf@karmanos.org

Copyright © 2006 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-05-0299

keratinocytes, and down-regulation of Notch-1 results in lower NF- κ B activity. Levels of basal and stimulation-induced NF- κ B activity were significantly decreased in mice with reduced Notch levels (20, 21). Constitutive levels of Notch activity are essential in maintaining NF- κ B activity in various cell types. Notch and NF- κ B pathways are key regulators of numerous cellular events such as proliferation, differentiation, and apoptosis. Therefore, inactivation of Notch-1-mediated cell growth inhibition and induction of apoptosis could be partly mediated via inactivation of NF- κ B activity.

However, the mechanisms by which Notch-1 inhibits apoptosis in pancreatic cancer cells are still unclear. Because Notch-1 down-regulation showed antineoplastic effects *in vivo* and *in vitro* (15–17, 22), the potential for treating certain cancers could be achievable by inhibiting Notch signal transduction. Genistein, a natural isoflavonoid found in soybean products, consumed in a diet, has been associated with lower incidences of endometrial, breast, prostate, and pancreatic cancers and is believed to be a chemopreventive agent (23). Studies from our laboratory and others have shown that genistein can inhibit cell growth and induce apoptosis in various cancer cell lines (24–26). However, little is known about the *Notch-1* gene alteration in pancreatic cancer cells after genistein treatment. Therefore, we hypothesized that genistein may inhibit Notch-1 activation in pancreatic cancer cells leading to apoptotic cell death. Therefore, in this report, we tested our hypothesis on whether down-regulation of *Notch-1* gene expression, either by small interfering RNA (siRNA) or by genistein, could inhibit cell growth and induce apoptosis, which are mechanistically associated with the

down-regulation of NF- κ B in pancreatic cancer cells. Our data show that down-regulation of Notch-1 inhibits cell growth with concomitant induction of apoptosis. Our data also show that genistein down-regulated the expression of Notch-1 and its downstream molecules, suggesting that Notch-1 down-regulation, especially by genistein, could be a novel therapeutic approach for the treatment of pancreatic cancer.

Materials and Methods

Cell Culture and Experimental Reagents

Human pancreatic cancer cell lines BxPC-3, HPAC, and PANC-1 (American Type Culture Collection, Manassas, VA) were cultured in RPMI 1640 (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum and 1% penicillin and streptomycin in a 5% CO₂ atmosphere at 37°C. Cell Death Detection ELISA Kit was obtained from Roche (Indianapolis, IN). Primary antibodies for Notch-1, Hes-1, cyclin D1, cyclin A, p21, p27, Bcl-2, and Bcl-X_L were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). All secondary antibodies were obtained from Pierce (Rockford, IL). Notch-1 siRNA and siRNA control were obtained from Santa Cruz Biotechnology. Lipofectamine 2000 was purchased from Invitrogen. Chemiluminescence detection of proteins was done with the use of a kit from Amersham Biosciences (Amersham Pharmacia Biotech, Piscataway, NJ). Genistein was obtained from Toronto Research Chemicals (North York, Ontario, Canada). Protease inhibitor cocktail, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), and all other chemicals were obtained from Sigma (St. Louis, MO).




Figure 1. Constitutive expression and down-regulation of Notch-1 by siRNA in pancreatic cancer cell lines BxPC-3, HPAC, and PANC-1. CS, control siRNA; NS, Notch-1 siRNA; CP, control plasmid; NP, Notch-1 plasmid. **A** and **B**, Notch-1 mRNA and protein levels were measured by real-time RT-PCR and Western blotting. **C**, Notch-1 mRNA levels were down-regulated by Notch-1 siRNA. **D**, *top*, Notch-1 protein levels were down-regulated by siRNA in all three pancreatic cancer cells. *Bottom*, Notch-1 protein levels were overexpressed by cDNA transfection in different cell lines.




Figure 2. Effect of down-regulation of Notch-1 by siRNA on pancreatic cancer cell growth and apoptosis. **A to C**, inhibition of cancer cell growth tested by MTT assay. **D to F**, cell death assay for measuring apoptosis tested by ELISA. *, $P < 0.05$; **, $P < 0.01$, relative to control siRNA.

Plasmids and Transfections

The Notch-1 cDNA plasmid encoding the Notch-1 intracellular domain and control plasmids were kind gifts from L. Miele (Department of Biopharmaceutical Sciences and Cancer Center, University of Illinois at Chicago, Chicago, IL; ref. 11). Cells were transfected with Notch-1 siRNA and siRNA control, respectively, using Lipofectamine 2000. Cells were stably transfected with human intracellular Notch or vector alone (pcDNA3) and maintained under neomycin selection.

Cell Growth Inhibition Studies by MTT Assay

The transfected cells (5×10^3) were seeded in a 96-well culture plate and subsequently incubated with MTT reagent (0.5 mg/mL) at 37°C for 2 hours and MTT assay was done as described earlier (27). Results were plotted as mean \pm SD of three separate experiments having six determinations per experiment for each experimental condition.

Histone/DNA ELISA for Detection of Apoptosis

The Cell Death Detection ELISA Kit was used for assessing apoptosis in transfected cells according to the protocol of the manufacturer. Briefly, the cells were lysed and the cell lysates were overlaid and incubated in microtiter plate modules coated with antihistone antibody. Samples were then incubated with anti-DNA peroxidase

followed by color development with ABTS substrate. The absorbances of the samples were determined with the Ultra Multifunctional Microplate Reader (Tecan, Durham, NC) at 405 nm.

Flow Cytometry and Cell Cycle Analysis

The cells were synchronized in G_0 by serum starvation for 24 hours in phenol red-free RPMI with 0.1% serum. Subsequently, cells were released into complete medium containing 10% fetal bovine serum. The cell cycle was analyzed by flow cytometry. Briefly, 1×10^6 cells were harvested and washed in PBS, then fixed in 70% alcohol for 30 minutes at 4°C . After washing in cold PBS thrice, cells were resuspended in 1 mL of PBS solution with 40 μg of propidium iodide and 100 μg of RNase A for 30 minutes at 37°C . Samples were then analyzed for their DNA content by FACSCalibur (Becton Dickinson, Mountain View, CA).

Western Blot Analysis

Cells were lysed in lysis buffer [50 mmol/L Tris (pH 7.5), 100 mmol/L NaCl, 1 mmol/L EDTA, 0.5% NP40, 0.5% Triton X-100, 2.5 mmol/L sodium orthovanadate, 10 $\mu\text{L}/\text{mL}$ protease inhibitor cocktail, 1 mmol/L phenylmethylsulfonyl fluoride] by incubating for 20 minutes at 4°C . The protein concentration was determined with the Bio-Rad assay system (Hercules, CA). Total proteins were fractionated by SDS-PAGE and transferred onto nitrocellulose membrane

for Western blotting as described earlier (27). Quantification of blotting was done using laser densitometry and the results are presented as the mean of three independent experiments with error bars representing SD.

Real-time Reverse Transcription-PCR Analysis for Gene Expression Studies

The total RNA from transfected cells was isolated by Trizol (Invitrogen) and purified by RNeasy Mini Kit and RNase-free DNase Set (Qiagen, Valencia, CA) according to the protocols of the manufacturer. One microgram of total RNA from each sample was subjected to first-strand cDNA synthesis using TaqMan Reverse Transcription Reagents (Applied Biosystems, Foster City, CA) in a total volume of 50 μ L, including 6.25 units of MultiScribe reverse transcriptase and 25 pmol of random hexamers. Reverse transcription reaction was done at 25°C for 10 minutes, followed by 48°C for 30 minutes and 95°C for 5 minutes. The primers used in the PCR reaction are Notch-1 forward primer (5'-CACTGTGGGCGGGTCC-3') and reverse primer (5'-GTTGTATTGGTTCGGCACCAT-3') and β -actin forward primer (5'-CCACACTGTGCCCATC-TACG-3') and reverse primer (5'-AGGATCTTCATGAGG-TAGTCAGTCAG-3'). Real-time PCR amplifications were done as described earlier (28).

Electrophoretic Mobility Shift Assay for Measuring NF- κ B Activity

The transfected cells were washed with cold PBS and suspended in 0.15 mL of lysis buffer [10 mmol/L HEPES (pH 7.9), 10 mmol/L KCl, 0.1 mmol/L EDTA, 0.1 mmol/L EGTA, 1 mmol/L DTT, 1 mmol/L phenylmethylsulfonyl fluoride, 2 μ g/mL leupeptin, 2 μ g/mL aprotinin, 0.5 mg/mL benzamide]. The nuclear protein was prepared and subjected to DNA binding activity of NF- κ B by electrophoretic mobility shift assay (EMSA) as described earlier (27).

Genistein Treatment of BxPC-3 Cells for Different Periods of Time

The transfected BxPC-3 cells were seeded in 100-mm dishes and allowed to attach for 24 hours, followed by the addition of different concentrations of genistein or 0.5 mmol/L Na₂CO₃ (solvent control) for different periods of time. The proteins were extracted and measured by Western blotting. In addition, the cell growth and apoptotic cell death in transfected cells with treatments were detected using MTT assay and Cell Death Detection ELISA Kit, respectively, following the procedure described earlier. NF- κ B activity was measured by EMSA as discussed above.





Figure 3. Effect of up-regulation of Notch-1 by cDNA plasmid on pancreatic cancer cell growth and apoptosis. **A to C**, promotion of cancer cell growth tested by MTT assay. **D to F**, cell death assay for measuring apoptosis tested by ELISA. *, $P < 0.05$; **, $P < 0.01$, relative to control plasmid.

Figure 4. Effect of the down-regulation of Notch-1 on cell cycle distribution. HPAC cells were harvested for cell cycle analysis using propidium iodide staining. X axis, DNA content; Y axis, number of nuclei. Compared with the control, down-regulation of Notch-1 caused G₀-G₁ cell cycle arrest.



Densitometric and Statistical Analysis

The bidimensional absorbances of Notch-1 and β -actin proteins on the films were quantified and analyzed with Molecular Analyst software (Bio-Rad). The ratios of Notch-1 against β -actin were calculated. The cell growth inhibition by transfection or genistein treatment was statistically evaluated with GraphPad StatMate software (GraphPad Software, Inc., San Diego, CA). Comparisons were made between control and transfection or genistein treatment. $P < 0.05$ was used to indicate statistical significance.

Results

Down-Regulation of Notch-1 Expression by siRNA Inhibited Cell Growth and Induced Apoptosis

Initial studies were done to examine the relative levels of Notch-1 in three pancreatic cancer cells, BxPC-3, HPAC, and PANC-1, by real-time RT-PCR and Western blot analysis. All three cell lines expressed high levels of Notch-1 at both mRNA and protein levels (Fig. 1A and B). To determine whether Notch-1 could be an effective therapeutic target for pancreatic cancer, the effect of Notch-1 siRNA on cell growth of the pancreatic cancer cells was examined. The efficacy of Notch-1 siRNA for knockdown of Notch-1 mRNA and protein was confirmed through real-time RT-PCR and Western blotting. We observed that both Notch-1 mRNA and protein levels were barely detectable in Notch-1 siRNA-transfected cells compared with siRNA control-transfected cells (Fig. 1C and D). The cell viability was determined by MTT and the effect of Notch-1 siRNA on the growth of cancer cells was shown in Fig. 2A to C. We found that down-regulation of Notch-1 expression caused cell growth inhibition in all three pancreatic cancer cell lines.

To investigate whether the growth inhibitory effects of Notch-1 siRNA are partially related to the induction of apoptosis, the effect of Notch-1 siRNA on apoptotic cell death was examined using an ELISA-based assay. These results provided convincing data that down-regulation of

Notch-1 induces apoptosis in all three pancreatic cancer cell lines (Fig. 2D-F). These data suggest that the growth inhibitory activity of Notch-1 down-regulation is partly attributed to an increase in cell death.

Overexpression of Notch-1 by cDNA Transfection Promoted Cell Growth and Inhibited Apoptosis

Pancreatic cancer cells were stably transfected with human intracellular Notch or empty vector alone (pcDNA3) and maintained under neomycin selection. The proteins were measured by Western blotting. The results showed that Notch-1 protein level was increased by intracellular Notch transfection (Fig. 1D). Intracellular Notch-transfected cells showed significant promotion of cell growth compared with empty vector-transfected control cells (Fig. 3A-C). We also found that overexpression of Notch-1 protected cells from apoptosis to a certain degree (Fig. 3D-F).

Down-Regulation of Notch-1 Induced Cell Cycle Arrest in G₀-G₁ Phase

To further investigate the growth inhibitory effect of Notch-1 knockdown in pancreatic cancer cells, we did cell




Figure 5. The level of expression of several known G₀-G₁ cell cycle regulatory factors as detected by Western blotting in all three pancreatic cancer cells.

Table 1. Flow cytometric analysis of Notch-1-transfected cells

	G ₀ -G ₁ (%)	S (%)	G ₂ -M (%)
BxPC-3			
CS	55.8 ± 0.7	30.6 ± 1.1	13.6 ± 0.9
NS	72.4 ± 1.4	22.1 ± 1.5	5.5 ± 0.6
CP	52.3 ± 3.5	29.2 ± 3.1	18.5 ± 0.4
NP	37.8 ± 1.2	42.4 ± 2.0	19.8 ± 3.1
HPAC			
CS	46.9 ± 2.4	36.3 ± 1.5	16.8 ± 1.2
NS	66.2 ± 1.8	24.7 ± 1.5	9.1 ± 1.1
CP	53.8 ± 2.0	34.4 ± 1.3	11.8 ± 2.1
NP	40.1 ± 1.9	36.8 ± 1.5	23.1 ± 1.6
PANC-1			
CS	40.5 ± 2.4	33.7 ± 1.8	25.8 ± 1.6
NS	67.4 ± 3.5	15.3 ± 1.2	17.3 ± 2.8
CP	46.1 ± 2.1	35.5 ± 1.5	18.4 ± 0.9
NP	29.8 ± 2.6	54.3 ± 1.2	15.9 ± 1.7

NOTE: Flow cytometry was done as described in Fig. 4. The mean values (± SE) represent the percentage of cells in the indicated phase of the cell cycle from three independent experiments.

Abbreviations: CS, control siRNA; NS, Notch-1 siRNA; CP, control plasmid; NP, Notch-1 plasmid.

cycle analysis using propidium iodide staining and flow cytometry. These results showed a typical G₀-G₁ arrest pattern in Notch-1 siRNA-transfected cells (Fig. 4). In contrast, synchronized intracellular Notch-transfected cells caused a greater drop in the fraction of cells at G₀-G₁ phase (Table 1).

To further characterize the G₀-G₁ arrest, we examined the level of expression of several known G₀-G₁ cell cycle regulatory factors. Consistent with cell cycle arrest, expression of cyclin A1, cyclin D1, and cyclin-dependent kinase (Cdk)-2 was found to be decreased whereas p21 and p27 expression was increased (Fig. 5), suggesting

mechanistic roles of these molecules during Notch-1-induced cell cycle progression and cell cycle arrest by Notch-1 siRNA. We have also detected the expression of other proteins, such as cyclin B, Cdk4, and Cdk6, and did not find any change in the expression of those proteins (data not shown).

We also analyzed the expression of the apoptosis-related proteins Bcl-2 and Bcl-X_L. Our data showed that Bcl-2 and Bcl-X_L expression was down-regulated in siRNA-transfected cells (Fig. 5).

Down-regulation of *Notch-1* gene expression by Notch-1 siRNA inhibited NF-κB DNA-binding activity. We investigated whether the downstream effect of Notch-1 down-regulation was mechanistically associated with the NF-κB pathway. Nuclear proteins from transfected cells were subjected to analysis for NF-κB DNA-binding activity as measured by EMSA. The specificity of NF-κB DNA binding to the DNA consensus sequence was confirmed by supershift assay. The results showed that down-regulation of Notch-1 significantly inhibited NF-κB DNA-binding activity compared with control whereas Notch-1 cDNA transfection caused activation of NF-κB DNA-binding activity in all three cell lines tested (Fig. 6). These results provide evidence for a mechanistic cross-talk between Notch-1 and NF-κB in pancreatic cancer.

Down-Regulation of Notch-1 by Genistein Inhibited Cell Growth and Induced Apoptosis

Thus far, our results clearly show that Notch-1 down-regulation contributes to cell growth inhibition and apoptosis. However, the usefulness of siRNA approach for the inactivation of Notch-1 in a therapeutic setting is not available at the present time. Therefore, we have tested whether natural products could down-regulate the Notch signaling, and if so, then our approach could be useful therapeutically. Using various known chemopreventive




Figure 6. Down-Regulation of *Notch-1* gene expression by Notch-1 siRNA inhibited NF-κB DNA-binding activity. Nuclear proteins from siRNA- and cDNA-transfected cells were subjected to analysis for NF-κB DNA-binding activity as measured by EMSA. **A to C**, down-regulation of Notch-1 inhibited NF-κB DNA-binding activity compared with control whereas Notch-1 cDNA transfection caused activation of NF-κB DNA-binding activity in all three cell lines tested. **D**, NF-κB supershift analyses. EMSA experiments were done by additional 30-min incubations with polyclonal supershift antibodies against p65 before the addition of labeled probe. 1, nonspecific antibody (anti-cyclin D1); 2, p65 antibody.




Figure 7. Down-regulation of Notch-1 by genistein inhibited cell growth and induced apoptosis. **A**, inhibition of Notch-1 mRNA after 6, 12, 18, 24, 48, and 72 h of treatment with 25 $\mu\text{mol/L}$ genistein in BxPC-3 pancreatic cancer cells. The mRNA level in BxPC-3 pancreatic cancer cells treated with genistein was assessed by real-time RT-PCR. The expression of Notch-1 at the mRNA level was down-regulated after genistein treatment. *, $P < 0.05$; **, $P < 0.01$, relative to solvent control. **B**, inhibition of Notch-1, Hes-1, cyclin D1, and Bcl-X_L protein expression by 25 $\mu\text{mol/L}$ genistein in BxPC-3 pancreatic cancer cells. Cells were treated with 25 $\mu\text{mol/L}$ genistein for 24, 48, and 72 h. Western blot analysis showed that the protein levels of Notch-1, Hes-1, Bcl-X_L, and cyclin D1 were down-regulated in genistein-treated BxPC-3 pancreatic cancer cells in a time-dependent manner. **C**, inhibitory effect of genistein on the growth of BxPC-3 pancreatic cancer cells tested by MTT assay. BxPC-3 cells treated with 15, 25, and 50 $\mu\text{mol/L}$ genistein for 3 d. The treatment of BxPC-3 pancreatic cancer cells with genistein resulted in cell growth inhibition. The inhibition of cell growth was dose and time dependent. *, $P < 0.01$, relative to solvent control. **D**, genistein-induced apoptosis in BxPC-3 pancreatic cancer cells measured by the histone/DNA fragment analysis using ELISA. Cells were treated with 25 $\mu\text{mol/L}$ genistein for 24, 48, or 72 h. The induction of apoptosis was time dependent and was found to be more pronounced after 48 to 72 h of treatment. *, $P < 0.01$, relative to solvent control. **E**, NF- κ B DNA binding activity was measured by EMSA. Genistein inhibits NF- κ B DNA binding activity in BxPC-3 pancreatic cancer cells. Cells were treated with 10, 25, and 50 $\mu\text{mol/L}$ genistein for 48 h. Nuclear extracts were prepared from control and genistein-treated cells and subjected to analysis for NF- κ B DNA binding activity as measured by EMSA. Genistein significantly inhibited NF- κ B DNA-binding activity whereas there was no change in Rb (used as a protein loading control).

agents such as curcumin, indole-3-carbinol, 3,3'-diindolylmethane, epigallocatechin gallate, genistein, and resveratrol, we found that genistein is the best agent tested thus far to down-regulate the expression of Notch-1 (data not shown). We used real-time RT-PCR and Western blotting to detect the Notch-1 level in BxPC-3 cells treated with genistein. The expression of *Notch-1* gene at the mRNA level was down-regulated after genistein treatment. The altered expression of *Notch-1* gene was observed as early as 6 hours after genistein treatment and was significantly more pronounced with longer treatment (Fig. 7A), suggesting transcriptional inactivation of *Notch-1* gene expression.

To verify whether the alternation of *Notch-1* gene at the level of transcription ultimately results in alternations at the level of translation, we conducted Western blotting for detection of Notch-1. Western blot analysis showed that the protein level of Notch-1 was down-regulated in genistein-

treated BxPC-3 cells in a time-dependent manner (Fig. 7B). These results are in direct agreement with the RT-PCR data showing that genistein regulates the transcription and translation of *Notch-1* gene. In addition, we found that the expression of Notch-1 downstream target genes, including *Hes-1*, *Bcl-X_L*, and *cyclin D1*, were also down-regulated in genistein-treated cells (Fig. 7B).

We further investigated whether down-regulation of Notch-1 by genistein resulted in the inhibition of cell growth and induction of apoptosis in BxPC-3 cells. The treatment of BxPC-3 pancreatic cancer cells for 1 to 3 days with 10, 25, and 50 $\mu\text{mol/L}$ of genistein resulted in cell growth inhibition in a dose- and time-dependent manner (Fig. 7C). BxPC-3 cells were treated with 25 $\mu\text{mol/L}$ genistein for 24, 48, and 72 hours, respectively, to measure the degree of apoptosis. The induction of apoptosis was time dependent (Fig. 7D) and was found to be more pronounced after 48 to 72 hours of treatment. Our results

also showed that genistein significantly inhibited NF- κ B DNA-binding activity (Fig. 7E), which was in direct agreement with results previously published by our laboratory (29).

Down-Regulation of Notch-1 Expression by siRNA Promotes Genistein-Induced Cell Growth Inhibition and Apoptosis

Down-regulation of Notch-1 by siRNA transfection showed less expression of Notch-1 protein as confirmed by Western blotting (Fig. 8A). We have also found that the down-regulation of Notch-1 expression significantly inhibited cell growth induced by genistein (Fig. 8B). Notch-1 siRNA-transfected BxPC-3 cells were significantly more sensitive to spontaneous and genistein-induced apoptosis (Fig. 8C). Nuclear extracts from Notch-1 siRNA-transfected BxPC-3 cells with different treatments were subjected to analysis for NF- κ B DNA-binding activity as measured by EMSA. The results showed that Notch-1 siRNA also inhibited NF- κ B DNA-binding activity; however, genistein plus Notch-1 siRNA inhibited NF- κ B activity to a greater degree compared with genistein alone (Fig. 8D).

Overexpression of Notch-1 by cDNA Transfection Reduced Genistein-Induced Cell Growth Inhibition and Apoptosis

Overexpression of Notch-1 by cDNA transfection showed overexpression of Notch-1 protein as confirmed by Western blot analysis (Fig. 9A), and this overexpression in Notch-1 rescued genistein-induced cell growth inhibition and

abrogated genistein-induced apoptosis to a certain degree (Fig. 9B and C). Overexpression of Notch-1 by cDNA transfection partly abrogated inactivation of NF- κ B DNA-binding activity by genistein (Fig. 9D). These results provide evidence for a potential cross-talk between Notch-1 and NF- κ B signaling pathways during genistein-induced cell growth inhibition and apoptosis in BxPC-3 cells.

Discussion

Notch signaling plays important roles in maintaining the balance among cell proliferation, differentiation, and apoptosis (10). The *Notch* gene is abnormally activated in many human malignancies. It has been reported that the function of Notch signaling in tumorigenesis can be either oncogenic or antiproliferative, and the function is context dependent (15). In a limited number of tumor types, including human hepatocellular carcinoma and small lung cancer, Notch signaling is antiproliferative rather than oncogenic (30–32). However, most of the studies showed opposite function of Notch in many human cancers including pancreatic cancer (17, 33). Notch family members, some Notch ligands, and downstream molecules, such as Hes-1, have been found to be up-regulated in pancreatic cancer tissues (34). In the present study, we investigated the role of Notch-1 in cell proliferation and apoptosis in pancreatic cancer cell lines. In our study, down-regulation of Notch-1 elicited a dramatic effect on growth inhibition and induction of





Figure 8. BxPC-3 pancreatic cancer cell growth inhibition and cell death induced by Notch-1 siRNA and genistein. **A**, Notch-1 expression was down-regulated by genistein and Notch-1 siRNA. Western blot analysis was used to detect the protein level of Notch-1. 1, control; 2, 25 μ mol/L genistein; 3, Notch-1 siRNA; and 4, Notch-1 siRNA plus 25 μ mol/L genistein. **B**, down-regulation of Notch-1 expression significantly inhibited cell growth. Genistein plus siRNA Notch-1 inhibited cell growth to a greater degree compared with genistein alone. **C**, BxPC-3 pancreatic cancer cell death induced by Notch-1 siRNA and genistein. Down-regulation of Notch-1 expression significantly increased apoptosis induced by genistein. Notch-1 siRNA-transfected BxPC-3 cells were significantly more sensitive to spontaneous and genistein-induced apoptosis. **D**, NF- κ B DNA binding activity was measured by EMSA. Genistein and Notch-1 siRNA significantly inhibited NF- κ B DNA-binding activity. Genistein plus Notch-1 siRNA inhibited NF- κ B DNA binding activity to a greater degree compared with genistein alone. Rb level served as nuclear protein loading control. Control, cells treated with DMSO; 1, control; 2, 25 μ mol/L genistein; 3, Notch-1 siRNA; and 4, Notch-1 siRNA plus 25 μ mol/L genistein.

Figure 9. Overexpression of Notch-1 by cDNA transfection reduced genistein-induced cell growth inhibition and apoptosis. **A**, the efficacy of Notch-1 cDNA for overexpression of Notch-1 protein was tested by Western blot analysis. **B** and **C**, overexpression of Notch-1 by cDNA transfection rescued genistein-induced cell growth inhibition and abrogated genistein-induced apoptosis to a certain degree. **D**, overexpression of Notch-1 by cDNA transfection partly abrogated inactivation of NF- κ B DNA-binding activity by genistein in BxPC-3 cells whereas there was no change in the levels of Rb (used as protein loading control). Control, cells treated with DMSO; 1, control plasmid; 2, control plasmid plus 25 μ mol/L genistein; 3, Notch-1 cDNA; and 4, Notch-1 cDNA plus 25 μ mol/L genistein.



apoptotic processes in pancreatic cancer cells, as shown by MTT assay and DNA/histone fragmentation analysis, respectively. In contrast, up-regulation of Notch-1 expression caused cell growth promotion and protected cells from apoptosis to a certain degree. Thus, our results provide *in vitro* evidence in support of the role of *Notch-1* as an oncogene rather than a tumor suppressor gene in pancreatic cancer cells.

Because down-regulation of Notch-1 by siRNA reduced cell growth, we wondered if cell cycle arrest was related to the cell growth inhibition. Indeed, we found that Notch-1 down-regulation increased cell population in G₀-G₁ phase. In contrast, up-regulation by overexpression of Notch-1 cDNA reduced the cell number in G₀-G₁ phase. Cell proliferation is tightly regulated by expression and activation of cell cycle-dependent cyclins, Cdks, and Cdk inhibitors. Cdk inhibitors have negative effects on cell cycle machinery by binding to various cyclin-Cdk complexes and inhibiting their activities. There are two classes of Cdk inhibitors, the INK4 family and the KIP/CIP family. The KIP/CIP family, including p21^{CIP}, p27^{KIP1}, and p57^{KIP2}, interact with cyclin A-Cdk2, cyclin E-Cdk2, cyclin D-Cdk4, and cyclin D-Cdk6 complexes and inhibit their activities (35–38). Progression of a cell through the cell cycle is promoted by a number of cyclin-dependent kinases (Cdk) that, when complexed with specific regulatory proteins called cyclins, drive the cells forward through the cell cycle.

Cyclin D interacts with Cdk2, Cdk4, and Cdk6, resulting in cell cycle progression through G₁ phase. To explore the mechanism involved in down-regulation of Notch-1-induced cell growth arrest, the expression of cell cycle proteins was examined. We observed a marked reduction in cyclin A, cyclin D1, and Cdk2 expression and a dramatic increase in p21^{CIP} and p27^{KIP1} expression in Notch-1 siRNA-transfected cells. In our study, the decrease in cyclin D1, cyclin A, and Cdk2 and the increase in Cdk inhibitor proteins, including p21^{CIP} and p27^{KIP1}, were strongly correlated with the altered cell cycle distribution phenotype and growth suppression. These results suggest that Notch-1 affects pancreatic cancer cell cycle by regulating the expression levels of some cyclins (cyclin D1 and cyclin A) and Cdk inhibitors (p21^{CIP} and p27^{KIP1}).

Recent reports have shown that Notch-1 expression regulates cell death through both apoptosis and cell cycle pathways in erythroleukemia cells with regulation of c-Jun NH₂-terminal kinase, Bcl-X_L, p21^{cip1}, p27^{kip1}, NF- κ B, and the retinoblastoma protein Rb (19). Besides the role of Notch on proliferation, Notch may also play a role in apoptosis. In the present study, we clearly showed that down-regulation of Notch-1 induced apoptosis in pancreatic cancer cells, as assessed by the cell death ELISA assay. To explore the molecular mechanism by which down-regulation of Notch-1 results in the induction of apoptosis in pancreatic cancer cells, we examined the expression of

antiapoptotic proteins Bcl-2 and Bcl-X_L in siRNA-transfected cells. We observed that down-regulation of Notch-1 reduced Bcl-2 and Bcl-X_L protein expression level. Because Bcl-2 and Bcl-X_L protect cells from apoptosis, our findings suggest that decreased Bcl-2 and Bcl-X_L expression may participate in apoptosis induced by down-regulation of Notch-1 in human pancreatic cancer cells. Thus, the inhibition of cell growth observed in pancreatic cancer cells treated with siRNA may be partly due to the increase in apoptosis.

Because NF- κ B plays important roles in many cellular processes including transcriptional regulation of Bcl-2 and Bcl-X_L, research on the interaction of NF- κ B activation with other cell signal transduction pathways, including the Notch pathway, has received increased attention in recent years. Notch-1 has been reported to cross-talk with NF- κ B pathway (18–20). Constitutive levels of Notch activity are essential in maintaining NF- κ B activity in various cell types. Levels of basal and stimulation-induced NF- κ B activity were significantly decreased in mice with reduced Notch levels (21). We observed that down-regulation of Notch-1 reduced NF- κ B activity. In contrast, overexpression of wild-type Notch-1 cDNA enhanced NF- κ B activity. Because NF- κ B pathways are key regulators of numerous cellular processes such as proliferation, differentiation, and apoptosis, our results clearly provide molecular evidence for a potential cross-talk between Notch and NF- κ B pathways and suggest that cell growth inhibition and apoptosis induced by the down-regulation of Notch-1 may be partly mediated by the NF- κ B pathway. Although we have shown that down-regulation of Notch-1 is feasible by Notch-1 siRNA, these approaches are not yet practically useful in the therapeutic arena. For that reason, we have used genistein in the current study, primarily because genistein was found to be a potent agent in the down-regulation of NF- κ B pathway and also because of the known cross-talk between Notch and NF- κ B pathways. We found that genistein down-regulated the transcription and translation of Notch-1 and its downstream genes, *Hes-1*, *cyclin D1*, *Bcl-X_L*, and *NF- κ B*. In addition, genistein elicited a dramatic effect on growth inhibition and induction of apoptotic processes in BxPC-3 cells. Overexpression of Notch-1 by Notch-1 cDNA transfection abrogated genistein-induced apoptosis to a certain degree. Therefore, we strongly believe that down-regulation of Notch-1 by genistein is mechanistically linked to cell proliferation and apoptotic processes. The molecular mechanism(s) by which genistein exerts its inhibitory effects on BxPC-3 pancreatic cells, as revealed in the present study, has opened up exciting avenues for devising novel therapeutic strategies. Therefore, the down-regulation of Notch-1 and NF- κ B by genistein could be a useful strategy in the schema of therapeutic approaches for the treatment of pancreatic cancer.

In summary, we found that Notch-1 plays a role in pancreatic cancer cell growth and apoptosis. Our data support the potential oncogenic role of Notch-1 in

pancreatic cancer. Down-regulation of Notch-1 induced G₀-G₁ phase cell cycle arrest, with reduced levels of cyclin D1 expression and increased p21^{CIP} and p27^{KIP1} expression. In addition, Notch-1 down-regulation also induced apoptosis, which was partly due to decreased Bcl-2 and Bcl-X_L protein expression in pancreatic cancer cells. It seems that NF- κ B is downstream of Notch-1 signaling because down-regulation of Notch-1 reduced NF- κ B activity. We also found that genistein could be an active agent for the down-regulation of Notch-1 and NF- κ B pathways. From these results, we conclude that Notch-1 down-regulation by genistein could be a novel therapeutic approach in pancreatic cancer.

Acknowledgments

We thank Carrie Koerner for her editorial assistance.

References

- Jemal A, Tiwari RC, Murray T, et al. Cancer statistics, 2004. *CA Cancer J Clin* 2004;54:8–29.
- Greenwald I. LIN-12/Notch signaling: lessons from worms and flies. *Genes Dev* 1998;12:1751–62.
- Artavanis-Tsakonas S, Rand MD, Lake RJ. Notch signaling: cell fate control and signal integration in development. *Science* 1999;284:770–6.
- Aster JC, Xu L, Karnell FG, Patriub V, Pui JC, Pear WS. Essential roles for ankyrin repeat and transactivation domains in induction of T-cell leukemia by notch1. *Mol Cell Biol* 2000;20:7505–15.
- Bresnick EH, Chu J, Christensen HM, Lin B, Norton J. Linking Notch signaling, chromatin remodeling, and T-cell leukemogenesis. *J Cell Biochem Suppl* 2000;Suppl 35:46–53.
- Osborne B, Miele L. Notch and the immune system. *Immunity* 1999;11:653–63.
- Yanagawa S, Lee JS, Kakimi K, Matsuda Y, Honjo T, Ishimoto A. Identification of Notch1 as a frequent target for provirus insertional mutagenesis in T-cell lymphomas induced by leukemogenic mutants of mouse mammary tumor virus. *J Virol* 2000;74:9786–91.
- Bigas A, Martin DI, Milner LA. Notch1 and Notch2 inhibit myeloid differentiation in response to different cytokines. *Mol Cell Biol* 1998;18:2324–33.
- Mumm JS, Kopan R. Notch signaling: from the outside in. *Dev Biol* 2000;228:151–65.
- Ohishi K, Katayama N, Shiku H, Varnum-Finney B, Bernstein ID. Notch signalling in hematopoiesis. *Semin Cell Dev Biol* 2003;14:143–50.
- Weijzen S, Rizzo P, Braid M, et al. Activation of Notch-1 signaling maintains the neoplastic phenotype in human Ras-transformed cells. *Nat Med* 2002;8:979–86.
- Leethanakul C, Patel V, Gillespie J, et al. Distinct pattern of expression of differentiation and growth-related genes in squamous cell carcinomas of the head and neck revealed by the use of laser capture microdissection and cDNA arrays. *Oncogene* 2000;19:3220–4.
- Rae FK, Stephenson SA, Nicol DL, Clements JA. Novel association of a diverse range of genes with renal cell carcinoma as identified by differential display. *Int J Cancer* 2000;88:726–32.
- Tohda S, Nara N. Expression of Notch1 and Jagged1 proteins in acute myeloid leukemia cells. *Leuk Lymphoma* 2001;42:467–72.
- Joutel A, Tournier-Lasserre E. Notch signalling pathway and human diseases. *Semin Cell Dev Biol* 1998;9:619–25.
- Jundt F, Anagnostopoulos I, Forster R, Mathas S, Stein H, Dorken B. Activated Notch1 signaling promotes tumor cell proliferation and survival in Hodgkin and anaplastic large cell lymphoma. *Blood* 2002;99:3398–403.
- Miele L, Osborne B. Arbiter of differentiation and death: Notch signaling meets apoptosis. *J Cell Physiol* 1999;181:393–409.
- Oswald F, Liptay S, Adler G, Schmid RM. NF- κ B2 is a putative target gene of activated Notch-1 via RBP-J κ . *Mol Cell Biol* 1998;18:2077–88.

19. Jang MS, Miao H, Carlesso N, et al. Notch-1 regulates cell death independently of differentiation in murine erythroleukemia cells through multiple apoptosis and cell cycle pathways. *J Cell Physiol* 2004;199:418–33.
20. Nickoloff BJ, Qin JZ, Chaturvedi V, Denning MF, Bonish B, Miele L. Jagged-1 mediated activation of notch signaling induces complete maturation of human keratinocytes through NF- κ B and PPAR γ . *Cell Death Differ* 2002;9:842–55.
21. Wang Y, Chan SL, Miele L, et al. Involvement of Notch signaling in hippocampal synaptic plasticity. *Proc Natl Acad Sci U S A* 2004;101:9458–62.
22. Nickoloff BJ, Osborne BA, Miele L. Notch signaling as a therapeutic target in cancer: a new approach to the development of cell fate modifying agents. *Oncogene* 2003;22:6598–608.
23. Goodman MT, Wilkens LR, Hankin JH, Lyu LC, Wu AH, Kolonel LN. Association of soy and fiber consumption with the risk of endometrial cancer. *Am J Epidemiol* 1997;146:294–306.
24. Li Y, Bhuiyan M, Sarkar FH. Induction of apoptosis and inhibition of c-erbB-2 in MDA-MB-435 cells by genistein. *Int J Oncol* 1999;15:525–33.
25. Li Y, Upadhyay S, Bhuiyan M, Sarkar FH. Induction of apoptosis in breast cancer cells MDA-MB-231 by genistein. *Oncogene* 1999;18:3166–72.
26. Li Y, Sarkar FH. Inhibition of nuclear factor κ B activation in PC3 cells by genistein is mediated via Akt signaling pathway. *Clin Cancer Res* 2002;8:2369–77.
27. Zhang Y, Banerjee S, Wang ZW, Marciniak DJ, Majumdar AP, Sarkar FH. Epidermal growth factor receptor-related protein inhibits cell growth and induces apoptosis of BxPC3 pancreatic cancer cells. *Cancer Res* 2005;65:3877–82.
28. Li Y, Hong X, Hussain M, Sarkar SH, Li R, Sarkar FH. Gene expression profiling revealed novel molecular targets of docetaxel and estramustine combination treatment in prostate cancer cells. *Mol Cancer Ther* 2005;4:389–98.
29. Li Y, Ellis KL, Ali S, et al. Apoptosis-inducing effect of chemotherapeutic agents is potentiated by soy isoflavone genistein, a natural inhibitor of NF- κ B in BxPC-3 pancreatic cancer cell line. *Pancreas* 2004;28:e90–5.
30. Shou J, Ross S, Koeppen H, de Sauvage FJ, Gao WQ. Dynamics of notch expression during murine prostate development and tumorigenesis. *Cancer Res* 2001;61:7291–7.
31. Sriuranpong V, Borges MW, Ravi RK, et al. Notch signaling induces cell cycle arrest in small cell lung cancer cells. *Cancer Res* 2001;61:3200–5.
32. Talora C, Sgroi DC, Crum CP, Dotto GP. Specific down-modulation of Notch1 signaling in cervical cancer cells is required for sustained HPV-E6/E7 expression and late steps of malignant transformation. *Genes Dev* 2002;16:2252–63.
33. Wang Z, Zhang Y, Banerjee S, Li Y, Sarkar FH. Inhibition of nuclear factor kappa B activity by genistein is mediated via Notch-1 signaling pathway in pancreatic cancer cells. *Int J Cancer* 2006;118:1930–6.
34. Miyamoto Y, Maitra A, Ghosh B, et al. Notch mediates TGF α -induced changes in epithelial differentiation during pancreatic tumorigenesis. *Cancer Cell* 2003;3:565–76.
35. Aprelikova O, Xiong Y, Liu ET. Both p16 and p21 families of cyclin-dependent kinase (CDK) inhibitors block the phosphorylation of cyclin-dependent kinases by the CDK-activating kinase. *J Biol Chem* 1995;270:18195–7.
36. Bartek J, Bartkova J, Lukas J. The retinoblastoma protein pathway in cell cycle control and cancer. *Exp Cell Res* 1997;237:1–6.
37. Pines J. Four-dimensional control of the cell cycle. *Nat Cell Biol* 1999;1:E73–9.
38. Sherr CJ. Mammalian G₁ cyclins. *Cell* 1993;73:1059–65.

Retraction: Downregulation of Notch-1 Contributes to Cell Growth Inhibition and Apoptosis in Pancreatic Cancer Cells



This article (1) has been retracted at the request of the editors. Following an institutional review by Wayne State University (Detroit, MI), the primary affiliation for several authors, it was determined that multiple Western blot bands in Fig. 5 were falsified and/or fabricated. As a result of these findings, the institution recommended retraction and, upon internal review, the editors agree with this recommendation.

A copy of this retraction notice was sent to the last known email addresses for five of the six authors. One author (J. Liao) agreed to the retraction; four authors (Z. Wang, Y. Li, S. Banerjee, and F.H. Sarkar) did not respond; the remaining author (Y. Zhang) could not be located.

Reference

1. Wang Z, Zhang Y, Li Y, Banerjee S, Liao J, Sarkar FH. Downregulation of Notch-1 contributes to cell growth inhibition and apoptosis in pancreatic cancer cells. *Mol Cancer Ther* 2006;5:483–93.

Published online October 1, 2018.

doi: 10.1158/1535-7163.MCT-18-0524

©2018 American Association for Cancer Research.

Molecular Cancer Therapeutics

Down-regulation of Notch-1 contributes to cell growth inhibition and apoptosis in pancreatic cancer cells

Zhiwei Wang, Yuxiang Zhang, Yiwei Li, et al.

Mol Cancer Ther 2006;5:483-493.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/5/3/483>

Cited articles This article cites 37 articles, 15 of which you can access for free at:
<http://mct.aacrjournals.org/content/5/3/483.full#ref-list-1>

Citing articles This article has been cited by 29 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/5/3/483.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/5/3/483>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.