

Targeting the Akt/mammalian target of rapamycin pathway for radiosensitization of breast cancer

Jeffrey M. Albert, Kwang Woon Kim,
Carolyn Cao, and Bo Lu

Department of Radiation Oncology, Vanderbilt Ingram
Cancer Center, Vanderbilt University School of Medicine,
Nashville, Tennessee

Abstract

The phosphatidylinositol 3-kinase (PI3K)/Akt pathway is known to be activated by radiation. The mammalian target of rapamycin (mTOR) is downstream of Akt, and we investigated the effects of radiation on Akt/mTOR signaling in breast cancer cell models. RAD001 (everolimus), a potent derivative of the mTOR inhibitor rapamycin, was used to study the effects of mTOR inhibition, as the role of mTOR inhibition in enhancing radiation remains unexplored. RAD001 decreased clonogenic cell survival in both breast cancer cell lines MDA-MB-231 and MCF-7, although the effect is greater in MDA-MB-231 cells. Irradiation induced Akt and mTOR signaling, and this signaling is attenuated by RAD001. The radiation-induced signaling activation is mediated by PI3K because inhibition of PI3K with LY294002 inhibited the increase in downstream mTOR signaling. Additionally, caspase-dependent apoptosis is an important mechanism of cell death when RAD001 is combined with 3 Gy radiation, as shown by induction of caspase-3 cleavage. An increase in G₂-M cell cycle arrest was seen in the combination treatment group when compared with controls, suggesting that cell cycle arrest may have been a contributing factor in the increased radiosensitization seen in this study. We conclude that RAD001 attenuates radiation-induced prosurvival Akt/mTOR signaling and enhances the cytotoxic effects of radiation in breast cancer cell models, showing promise as a method of radiosensitization of breast cancer. [Mol Cancer Ther 2006;5(5):1183–9]

Received 10/3/05; revised 1/23/06; accepted 3/2/06.

Grant support: Vanderbilt Discovery Grant, Vanderbilt Physician Scientist Grant, Mesothelioma Applied Research Foundation, and Department of Defense grants PC031161 and BC030542.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: J.M. Albert and K.W. Kim contributed equally to this work.

Requests for reprints: Bo Lu, Department of Radiation Oncology, Vanderbilt University, B-902 The Vanderbilt Clinic, 1301 22nd Avenue South, Nashville, TN 37232-5671. Phone: 615-343-9233; Fax: 615-343-3075. E-mail: bo.lu@vanderbilt.edu

Copyright © 2006 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-05-0400

Introduction

The phosphatidylinositol 3-kinase (PI3K)/Akt pathway is a cell survival pathway that is important for normal cell growth and proliferation (1). This pathway has also been implicated in tumorigenesis (2) and is becoming an important target for cancer treatment (3, 4). The PI3K/Akt pathway has also been specifically shown to promote breast cancer cell survival and resistance to tamoxifen (5) and many other chemotherapeutic agents (6). Clinically, increased Akt activity has been associated with decreased survival in tamoxifen-treated breast cancer patients, supporting the role of Akt in tamoxifen resistance (7). Additionally, patients who test positive for activated phospho-Akt are more likely to relapse with distant metastasis (8).

Mammalian target of rapamycin (mTOR) is a 289-kDa serine/threonine kinase that is a downstream target of Akt (9). It has been shown that mTOR is important for the oncogenic transformation induced specifically by PI3K and Akt, and that the mTOR inhibitor rapamycin inhibits cellular transformation (10). Recently, mTOR has been explored as a target for cancer therapy (3). mTOR presents an attractive target in the pathway because it is downstream of PI3K and Akt. Thus, mTOR inhibition could avoid possible side effects from inhibiting these broader function upstream proteins (3, 11). The normal activation of mTOR results in an increase in protein translation because mTOR phosphorylates and activates the translation regulators eukaryotic initiation factor 4E-binding protein 1 and ribosomal p70 S6 kinase (12, 13). Therefore, by inhibiting mTOR, rapamycin causes a decrease in phosphorylation of these effectors, and a decrease in protein synthesis, effectively blocking the pro-growth, pro-proliferative, and prosurvival actions of mTOR (14).

The rapamycin analogue CCI-779 has been shown to be an effective chemotherapeutic agent against breast cancer in phase II clinical trials (15), and the rapamycin derivative RAD001 is also currently in clinical trials. It has been recently suggested that combining RAD001 with DNA-damaging chemotherapeutic agents may be an effective means of enhancing cancer treatment (16). However, the role of mTOR inhibition in enhancing radiation-induced DNA damage remains unexplored. We aim to investigate the combination of the mTOR inhibitor RAD001 with radiation.

Breast cancer is the most common type of cancer in women, excluding nonmelanoma skin cancer, and is the second leading cause of cancer deaths in women. Based on current data, 13.2% of all women will be diagnosed with invasive breast cancer (17). Although early detection is leading to declining mortality rates, the high incidence of breast cancer requires continued use of different treatment modalities, including combinations of surgery,

radiotherapy, and chemotherapy. In particular, radiotherapy plays a crucial role in achieving local control following surgery. Accordingly, development of radioresistance in breast cancer cells presents a difficult problem in the course of treatment (18). Therefore, a breast cancer cell model is an important and useful model when studying radiosensitization and cancer cell survival pathways. The purpose of the present study was to determine the effects of radiation on mTOR signaling and determine whether mTOR inhibition with RAD001 enhances the cytotoxic effects of radiation in breast cancer cells. We found that radiation induces prosurvival Akt/mTOR signaling, and that RAD001 attenuates this induction and enhances the cytotoxic effects of radiation. Additionally, caspase-dependent apoptosis and cell cycle arrests seemed to contribute to the increased levels of cell death.

Materials and Methods

Cell Culture

MDA-MB-231 cells and MCF-7 cells (obtained from the American Type Culture Collection, Rockville, MD) were cultured in BMEM (DMEM, Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin. All cells were incubated at 37°C and humidified 5% CO₂.

Western Immunoblots

Cells were treated with various radiation doses and drugs and collected at different time points, according to the individual study. The cells were harvested and then washed with ice-cold PBS twice before the addition of lysis buffer. Protein concentration was quantified using the Bio-Rad method. Equal amounts of protein were loaded into each well and separated by 10% SDS-PAGE gel followed by transfer onto polyvinylidene difluoride membranes (Bio-Rad, Richmond, CA). Membranes were blocked using 5% nonfat dry milk in PBS-T. The blots were then incubated with total caspase-3, cleaved caspase-3, mTOR, phospho-mTOR (Ser²⁴⁸; Cell Signaling, Beverly, MA), Akt, phospho-Akt (Ser⁴⁷³; Cell Signaling), S6 ribosomal protein, or phospho-S6 ribosomal protein (Ser²⁴⁰/Ser²⁴⁴; Cell Signaling) antibodies overnight at 4°C. Goat anti-rabbit IgG secondary antibody (1:1,000; Santa Cruz Biotechnology, Santa Cruz, CA) was incubated for 1 hour at room temperature. Immunoblots were developed using the chemiluminescence detection system (Perkin-Elmer, Wellesley, MA) according to the manufacturer's protocol and autoradiography. Image analysis was used to quantify the intensity of the bands.

In vitro Clonogenic Assay

Cells were treated with RAD001 (10 nmol/L) or DMSO control for 1 hour. Various doses of radiation were given, and the medium was changed. After irradiation, cells were returned to the 37°C incubator and maintained for 10 days. Cells were fixed for 15 minutes with 3:1 methanol/acetic acid and stained for 15 minutes with 0.5% crystal violet (Sigma, St. Louis, MO) in methanol. After staining, colonies were counted with a cutoff of 50 viable cells. Surviving fraction was calculated as (mean colonies counted) / (cells

plated × plating efficiency), where plating efficiency was defined as (mean colonies counted) / (cells plated) for nonirradiated controls. Mean, SD, and *P* (for RAD001-treated cells versus DMSO control, using a Student's *t* test) were calculated.

Cell Cycle Analysis

Cells (10⁶) were seeded in 10-cm² dishes 24 hours before treatment with 5 Gy radiation and/or 10 nmol/L RAD001. The cells were collected at 8 and 24 hours by trypsinization. They were fixed with 70% ethanol and stored overnight at -20°C. After thawing, cells were collected by centrifugation, and the cell pellet was resuspended in 1 mL of PBS with propidium iodide (50 µg/mL). Cell number in each phase of the cell cycle was determined and calculated as a percentage of the total cell population. The analysis was repeated thrice, and the mean value and SD were calculated and graphed.

Results

Radiation Induces mTOR Signaling in Breast Cancer Cells

Western analysis was used to determine the effects of radiation and mTOR inhibition on mTOR signaling in breast cancer cells. The relative levels of the probed proteins were estimated by the densitometer and compared among the samples collected at different time points. After 15 minutes, phospho-mTOR was increased in both breast cancer cell lines used, and this increase persisted through 180 minutes (Fig. 1A and B). The phospho-mTOR was increased 1.4-fold from control in MDA-MB-231 cells and increased 2.0-fold in MCF-7 cells. Additionally, phospho-Akt was increased between 1.7- and 1.8-fold in both cell lines, and phospho-S6 ribosomal protein (a downstream marker of mTOR signaling) was increased 1.2- to 1.3-fold in both cell lines, showing increased signaling both upstream and downstream from mTOR. Notably, phospho-S6 levels began to decrease after 60 minutes in both cell lines despite phospho-mTOR levels remaining elevated and were reduced to control levels by 180 minutes (Fig. 1). To establish whether activation of mTOR signaling was mediated by PI3K, cells were treated with either the known PI3K inhibitor LY294002 (10 µmol/L) or RAD001 (20 nmol/L). A slightly higher concentration of RAD001 was used for Western immunoblots than for the other assays to emphasize cell signaling inhibition and protein level changes. After 1 hour of treatment, cells were treated with 3 Gy radiation. Cell lysates were then collected and analyzed. As shown in Fig. 2, MDA-MB-231 cells treated with 3 Gy showed dramatically increased levels of S6 phosphorylation, from negligible levels to strong induction. A moderate increase in S6 phosphorylation (~1.2-fold) was seen in MCF-7 cells (Fig. 2). Cells treated with LY294002 or RAD001 showed very little S6 phosphorylation with or without irradiation, similar to nonirradiated controls.

mTOR Inhibition Sensitizes Breast Cancer Cells to Radiation by Decreasing Cell Survival

Using clonogenic assays, the PI3K inhibitor LY294002 has been previously shown to sensitize cancer cells to




Figure 1. Radiation induces mTOR signaling in breast cancer cells. MDA-MB-231 (**A**) and MCF-7 (**B**) cells were incubated without serum for 7 h and irradiated with 3 Gy. Total cell lysates were extracted at indicated time points after irradiation. C indicates nonirradiated control in both **A** and **B**; 50 μ g total protein per lane were probed with phospho-mTOR (*p-mTOR*), phospho-Akt (*p-Akt*), and phospho-S6 (*P-S6*) ribosomal protein antibodies. mTOR, Akt, and S6 ribosomal protein were also probed to show equal loading.

radiation (19–21). In the present study, clonogenic assays were used to determine whether downstream mTOR inhibition with RAD001 radiosensitizes MDA-MB-231 and MCF-7 breast cancer cells. Cells were treated with RAD001 (10 nmol/L) or DMSO control for 1 hour before irradiation, based on previous studies (22, 23). One hour is long enough for mTOR inhibition to occur but not long enough for RAD001 to show a significant cytotoxic effect. Radiation doses from 0 to 6 Gy were then given, and the medium was changed. The colonies grew for 10 days and were then stained, scored, and graphed. A decrease in the survival curve of irradiated cells was seen for both breast cancer cell lines (Fig. 3). The effect is greater in MDA-MB-231 cells (**A**) than in MCF-7 cells (**B**).

Induction of Caspase-3 Cleavage in MDA-MB-231 Breast Cancer Cells

Western analysis was done to determine whether mTOR inhibition with RAD001 induces caspase-dependent apoptosis in MDA-MB-231 cells. This cell line was chosen for closer mechanism examination because the clonogenic assay shown a stronger radiosensitizing effect than in the MCF-7 cells (Fig. 3), and Western analysis showed that radiation induced much more S6 phosphorylation than in MCF-7 cells (Fig. 2). Some caspase-3 cleavage was induced by 3 Gy or RAD001 alone. However, significantly more caspase-3 was cleaved using the RAD001/3 Gy combination than with either treatment alone (Fig. 4A). To further analyze the effects of combination treatment on caspase-3 activity, pan-caspase-inhibitor z-VAD-fmk was combined with 3 Gy and RAD001. As expected, z-VAD-fmk notably reduced the level of cleaved caspase-3 seen with combination treatment. To quantify the caspase-3 activity seen in the different treatment groups, the intensity of the cleaved caspase-3 bands was then measured and calculated relative to actin control (Fig. 4B). The control cells showed insignificant levels of caspase-3 cleavage; 3 Gy alone had a relative intensity of about 0.33, RAD001 alone was 0.19, but combination 3 Gy + RAD001 was 0.7. Adding z-VAD-fmk to the 3 Gy/RAD001 combination reduced the band intensity to 0.27.

Cell Cycle Analysis of MDA-MB-231 Breast Cancer Cells

Changes in the cell cycle have been previously analyzed to explore radiosensitivity after mTOR inhibition (22). To determine whether changes in cell cycle distribution contributed to the increased radiosensitivity of MDA-MB-231 cells, propidium iodide staining and flow cytometry were used. The experiment was repeated thrice, and the mean cell cycle distribution for DMSO control, 5 Gy radiation, 10 nmol/L RAD001, and 5 Gy/RAD001 combination was graphed (Fig. 5). Eight hours after treatment, there was no notable change in cell cycle distribution seen after treatment with 5 Gy, RAD001, or combination 5 Gy/RAD001 (data not shown). At 24 hours, there was no significant change in cell cycle distribution after treatment with RAD001 alone. However, there was a moderate increase in G₂-M phase cells after treatment with 5 Gy




Figure 2. RAD001 inhibits radiation-induced mTOR signaling. MDA-MB-231 cells and MCF-7 cells were treated with DMSO, LY294002 (10 μ mol/L), or RAD001 (20 nmol/L) for 1 h before irradiation (3 Gy). After 30 min of irradiation, total cell lysates were prepared in lysis buffer; 50 μ g total protein per lane were immunoblotted for phospho-S6 (*P-S6*) ribosomal protein. S6 ribosomal protein was probed to show equal loading.




Figure 3. mTOR inhibition radiosensitizes MDA-MB-231 and MCF-7 breast cancer cells. **A**, MDA-MB-231 cells treated with radiation alone and radiation with RAD001 at indicated doses. **B**, MCF-7 cells that were similarly treated. Cells were treated with 10 nmol/L RAD001 or DMSO for 1 h. Cells were then irradiated with 0, 2, 4, or 6 Gy, and the medium was changed. After 2 wks, colonies were scored. Points, mean; bars, SD. $P < 0.003$, for RAD001 versus control in MDA-MB-231 cells; $P < 0.004$, for RAD001 versus control in MCF-7 cells.

alone, and a large increase in G₂-M phase cells was seen 24 hours after combination treatment (Fig. 5). At 24 hours, the percentage of cells in G₂-M phase was 15.0% (SD 1.2%) for control, 33.5% (SD 0.9%) for 5 Gy alone, 7.7% (SD 0.4%) for RAD001 alone, and 69.6% (SD 0.5%) for RAD001 + 5 Gy.

Discussion

We have found that mTOR signaling is increased by radiation in breast cancer cell models, and that inhibition of mTOR obstructs this signaling activation. The mTOR inhibitor RAD001 radiosensitizes the two models used in this study, although the effect is greater in MDA-MB-231 cells than in MCF-7 cells. The study of mTOR as a therapeutic target is becoming increasingly important in cancer research. The role of mTOR as a downstream kinase in the PI3K/Akt pathway, a pathway crucial to cell growth and survival, makes it a clear target.

Rapamycin is well characterized as an mTOR inhibitor (24) and has led to the development of rapamycin derivatives with improved pharmaceutical properties (i.e., CCI-779 and RAD001; ref. 25). Both of these mTOR inhibitors are in clinical trials, and have shown significant chemotherapeutic antitumor activity in various models (14, 15).

In the present study, we showed that mTOR signaling is induced by ionizing radiation in the two breast cancer cell lines MDA-MB-231 and MCF-7, which is consistent with the PI3K/Akt pathway activation seen in breast cancer cells in previous studies (26, 27). Because radiotherapy is important as adjuvant therapy for breast cancer, it is undesirable that radiation can contribute to cancer cell survival. Therefore, blocking radiation-induced PI3K/Akt/mTOR pathway activation presents a method of enhancing the cytotoxic effects of radiation. To show this, we showed that treating breast cancer cells with the mTOR inhibitor RAD001 blocked the increase in phosphorylation of the downstream marker S6 ribosomal protein. This strongly suggests that mTOR inhibition is an effective means of blocking the prosurvival response of breast cancer cells to radiation and suggests a mechanism for how RAD001 may enhance the efficacy of radiotherapy.

One notable unexpected result was that phospho-S6 levels decreased after 60 minutes in both cell lines, despite no corresponding decrease in phospho-mTOR levels. This finding suggests that in addition to mTOR, other pathways may be regulating the phosphorylation of S6 protein, perhaps through feedback inhibition. For example,




Figure 4. Induction of caspase-3 cleavage in treated MDA-MB-231 cells. MDA-MB-231 cells were treated with DMSO, 3 Gy, RAD001 (20 nmol/L), RAD001 + 3 Gy, z-VAD-fmk (50 μ mol/L), or z-VAD-fmk + 3 Gy + RAD001. **A**, Western blots probed for cleaved caspase-3, total caspase-3, and actin. **B**, the intensity of the bands was then quantified and calculated as intensity relative to actin control.




Figure 5. Cell cycle analysis of treated MDA-MB-231 cells. Cells were treated with DMSO control, 5 Gy radiation, 10 nmol/L RAD001, and 5 Gy/RAD001 combination. Cells were then fixed and resuspended in propidium iodide (50 μ g/mL). Cell number in each phase of the cell cycle was determined and calculated as a percentage of the total cell population and then graphed. Columns, mean from three repeated studies; bars, SD.

phosphoinositide-dependent protein kinase 1 has been shown to activate ribosomal p70 S6 kinase, which in turn phosphorylates S6. Phosphoinositide-dependent protein kinase 1 can bypass the Akt/mTOR pathway and directly activate ribosomal p70 S6 kinase (28–30). It has also been suggested that the c-Raf/mitogen-activated protein/extracellular signal-regulated kinase kinase/extracellular signal-regulated kinase pathway can control activation of ribosomal p70 S6 kinase (31), providing another possible mechanism for the late decrease in phospho-S6 seen in the present study.

A recent study used a combination of RAD001 and the DNA-damaging agent cisplatin to show that RAD001 sensitizes cancer cells to DNA damage-induced apoptosis (16). We hypothesized that combining radiation, a DNA-damaging agent, with RAD001 would also be effective at sensitizing cancer cells to apoptosis. We, therefore, investigated apoptosis in MDA-MB-231 cells treated with RAD001 and radiation. Western analysis probing for cleaved caspase-3 showed that apoptosis was greatly increased in these cells after combination treatment with RAD001 and 3 Gy versus treatment with RAD001 or 3 Gy alone (Fig. 4A), indicating that caspase-dependent apoptosis contributes to the mechanism of the observed increase in cell death. When considering the relative intensities of the cleaved caspase-3 bands in the Western blot, 3 Gy alone was 0.33, RAD001 alone was 0.19, and 3 Gy/RAD001 combination was 0.7, showing that the combination treatment induced more than an additive amount of apoptosis (Fig. 4B). However, this does not account for the total observed increase in cell death, suggesting that RAD001 functions via both apoptosis-dependent and apoptosis-independent pathways. Additionally, cell cycle regulation is important in mediating radiosensitivity. It is known that cell cycle arrested cells can be radioresistant (32), and cell cycle distribution has been previously

examined in glioma cells to assess radiosensitivity after treatment with rapamycin (22). It is also established that cells have varying radiosensitivity in different cell cycle phases. That is, cells are most sensitive to radiation during the G₂-M phase, less sensitive during G₁, and least sensitive near the end of S phase (33). We found a time-dependent effect on the cell cycle after treatment of MDA-MB-231 cells with radiation and/or RAD001. At 8 hours, there was a minimal change in cell cycle distribution compared with control for treatment with RAD001, 5 Gy, or combination. However, at 24 hours, there was a radiation-induced G₂-M arrest (33.5% G₂-M compared with 15.0% G₂-M in control), but no G₂-M arrest was seen with RAD001 alone.

Furthermore, a much larger increase in G₂-M arrest was seen after combination treatment (69.6% G₂-M; Fig. 5). Therefore, cell cycle arrest may be a contributing factor in the increased radiosensitivity seen in this study.

It is important to consider the differences between the two cell lines used in this study. Again, the MDA-MB-231 cells showed more radiation-induced mTOR signaling than MCF-7 cells and increased radiosensitivity after treatment with RAD001. There are several possible explanations for these differences. Our finding that mTOR signaling is less activated by radiation in MCF-7 cells implies that there is a smaller target for mTOR inhibition by RAD001. Because MDA-MB-231 cells have much more mTOR signaling activation after irradiation, there is a greater potential for mTOR inhibition. This suggests a possible mechanism for the observed difference in radiosensitization. It is also noteworthy that MCF-7 cells are deficient in caspase-3, an important effector in cellular apoptosis (34, 35). We also found that there were insignificant levels of caspase-3 present in MCF-7 cells (data not shown). Apoptosis resistance in MCF-7 cells treated with radiation has been shown to be caused by lack of caspase-3, and reexpression of caspase-3 in these cells caused an increase in apoptotic changes (36). Consistent with Zapata et al. (37), we found that MDA-MB-231 cells express high levels of caspase-3, and we also found that cleaved caspase-3 was increased after treatment with radiation and RAD001 (Fig. 4). This shows that the radiosensitization observed in MDA-MB-231 cells is at least partially attributable to caspase-dependent apoptotic cell death. Therefore, in the present study, the inability of MCF-7 cells to undergo caspase-dependent apoptosis may have been a contributing factor in their decreased radiosensitivity.

Our data provide additional evidence to the growing amount of research that indicates the effectiveness of mTOR inhibition in treating cancer. Specifically, we suggest that mTOR inhibition with rapamycin derivatives provides a means of radiosensitizing breast cancer cells. Of note, mTOR inhibitors are likely to be more effective at targeting tumor cells than normal cells. This is because transformed cells show increased activation of the Akt/mTOR pathway (10) and therefore present a better target with larger potential for the effects of mTOR inhibition. There are additional possible mechanisms of rapamycin derivatives

against tumorigenesis and tumor growth that were not investigated in the present study but propose interesting directions for future studies. For example, phosphate and tensin homologue on chromosome 10 (PTEN) normally opposes activation of the PI3K/Akt pathway (38). Although both cell lines used in this study are wild type for PTEN (39), mutations in at least one copy of the *PTEN* gene are found in ~50% of patients with breast cancer (40). Decreased PTEN expression in breast cancer cells has been shown to increase sensitivity to inhibition of the PI3K/Akt pathway (39, 41), emphasizing the expanded role that mTOR inhibitors can play in treating the large percentage of breast cancers with mutated PTEN.

It is possible that mTOR inhibition with rapamycin analogues would show a greater increase in radiosensitivity in breast cancer cells *in vivo*. Glioma models have previously shown no increased radiosensitivity in response to rapamycin derivatives *in vitro* (22, 23) but have shown significantly increased radiosensitivity *in vivo* (22). This suggests that rapamycins have an additional effect that is dependent on its action within whole tumors, as opposed to a tumor cell monolayer. Additionally, it has been shown that rapamycin inhibits angiogenesis (42), and tumor vasculature is significantly sensitized to radiotherapy by rapamycin and RAD001 both *in vitro* and *in vivo* (23). This suggests that combining mTOR inhibition with radiation may result in a dual mechanism of tumor inhibition, promoting both tumor cell cytotoxicity and inhibiting tumor angiogenesis. Therefore, further studies are indicated to examine the effects of mTOR inhibition combined with radiation in breast tumor models *in vivo*.

References

- Cantley LC. The phosphoinositide 3-kinase pathway. *Science* 2002; 296:1655–7.
- Nicholson KM, Anderson NG. The protein kinase B/Akt signalling pathway in human malignancy. *Cell Signal* 2002; 14:381–95.
- Lu Y, Wang H, Mills GB. Targeting PI3K-AKT pathway for cancer therapy. *Rev Clin Exp Hematol* 2003; 7:205–28.
- Morgensztern D, McLeod HL. PI3K/Akt/mTOR pathway as a target for cancer therapy. *Anticancer Drugs* 2005; 16:797–803.
- Campbell RA, Bhat-Nakshatri P, Patel NM, Constantinidou D, Ali S, Nakshatri H. Phosphatidylinositol 3-kinase/AKT-mediated activation of estrogen receptor alpha: a new model for anti-estrogen resistance. *J Biol Chem* 2001; 276:9817–24.
- Knuefermann C, Lu Y, Liu B, et al. HER2/PI-3K/Akt activation leads to a multidrug resistance in human breast adenocarcinoma cells. *Oncogene* 2003; 22:3205–12.
- Kirkegaard T, Witton CJ, McGlynn LM, et al. AKT activation predicts outcome in breast cancer patients treated with tamoxifen. *J Pathol* 2005; 207:139–46.
- Perez-Tenorio G, Stal O. Activation of AKT/PKB in breast cancer predicts a worse outcome among endocrine treated patients. *Br J Cancer* 2002; 86:540–5.
- Nave BT, Ouwens M, Withers DJ, Alessi DR, Shepherd PR. Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino-acid deficiency on protein translation. *Biochem J* 1999; 344 Pt 2: 427–31.
- Aoki M, Blazek E, Vogt PK. A role of the kinase mTOR in cellular transformation induced by the oncoproteins P3k and Akt. *Proc Natl Acad Sci U S A* 2001; 98:136–41.
- Chan S. Targeting the mammalian target of rapamycin (mTOR): a new approach to treating cancer. *Br J Cancer* 2004; 91:1420–4.
- Brunn GJ, Hudson CC, Sekulic A, et al. Phosphorylation of the translational repressor PHAS-1 by the mammalian target of rapamycin. *Science* 1997; 277:99–101.
- Burnett PE, Barrow RK, Cohen NA, Snyder SH, Sabatini DM. RAFT1 phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1. *Proc Natl Acad Sci U S A* 1998; 95:1432–7.
- Huang S, Houghton PJ. Targeting mTOR signaling for cancer therapy. *Curr Opin Pharmacol* 2003; 3:371–7.
- Chan S, Scheulen ME, Johnston S, et al. Phase II study of temsirolimus (CCI-779), a novel inhibitor of mTOR, in heavily pretreated patients with locally advanced or metastatic breast cancer. *J Clin Oncol* 2005; 23:5314–22.
- Beuvink I, Boulay A, Fumagalli S, et al. The mTOR inhibitor RAD001 sensitizes tumor cells to DNA-damaged induced apoptosis through inhibition of p21 translation. *Cell* 2005; 120:747–59.
- Ries LAG, Eisner MP, Kosary CL, et al. (eds). SEER cancer statistics review, 1975–2002. Bethesda, MD: National Cancer Institute; 2005.
- Jameel JK, Rao VS, Cawkwell L, Drew PJ. Radioresistance in carcinoma of the breast. *Breast* 2004; 13:452–60.
- Liang K, Lu Y, Jin W, Ang KK, Milas L, Fan Z. Sensitization of breast cancer cells to radiation by trastuzumab. *Mol Cancer Ther* 2003; 2: 1113–20.
- Nakamura JL, Karlsson A, Arvold ND, et al. PKB/Akt mediates radiosensitization by the signaling inhibitor LY294002 in human malignant gliomas. *J Neurooncol* 2005; 71:215–22.
- Gottschalk AR, Doan A, Nakamura JL, Stokoe D, Haas-Kogan DA. Inhibition of phosphatidylinositol-3-kinase causes increased sensitivity to radiation through a PKB-dependent mechanism. *Int J Radiat Oncol Biol Phys* 2005; 63:1221–7.
- Eshleman JS, Carlson BL, Mladek AC, Kastner BD, Shide KL, Sarkaria JN. Inhibition of the mammalian target of rapamycin sensitizes U87 xenografts to fractionated radiation therapy. *Cancer Res* 2002; 62: 7291–7.
- Shinohara ET, Cao C, Niermann K, et al. Enhanced radiation damage of tumor vasculature by mTOR inhibitors. *Oncogene* 2005; 24: 5414–22.
- Sabers CJ, Martin MM, Brunn GJ, et al. Isolation of a protein target of the FKBP12-rapamycin complex in mammalian cells. *J Biol Chem* 1995; 270:815–22.
- Huang S, Houghton PJ. Inhibitors of mammalian target of rapamycin as novel antitumor agents: from bench to clinic. *Curr Opin Investig Drugs* 2002; 3:295–304.
- Contessa JN, Hampton J, Lammering G, et al. Ionizing radiation activates Erb-B receptor dependent Akt and p70 S6 kinase signaling in carcinoma cells. *Oncogene* 2002; 21:4032–41.
- Liang K, Jin W, Knuefermann C, et al. Targeting the phosphatidylinositol 3-kinase/Akt pathway for enhancing breast cancer cells to radiotherapy. *Mol Cancer Ther* 2003; 2:353–60.
- Dufner A, Thomas G. Ribosomal S6 kinase signaling and the control of translation. *Exp Cell Res* 1999; 253:100–9.
- Shah OJ, Anthony JC, Kimball SR, Jefferson LS. 4E-BP1 and S6K1: translational integration sites for nutritional and hormonal information in muscle. *Am J Physiol Endocrinol Metab* 2000; 279: E715–29.
- Asnaghi L, Bruno P, Priulla M, Nicolin A. mTOR: a protein kinase switching between life and death. *Pharmacol Res* 2004; 50: 545–9.
- Iijima Y, Laser M, Shiraishi H, et al. c-Raf/MEK/ERK pathway controls protein kinase C-mediated p70S6K activation in adult cardiac muscle cells. *J Biol Chem* 2002; 277:23065–75.
- Hwang HS, Davis TW, Houghton JA, Kinsella TJ. Radiosensitivity of thymidylate synthase-deficient human tumor cells is affected by progression through the G₁ restriction point into S-phase: implications for fluoropyrimidine radiosensitization. *Cancer Res* 2000; 60:92–100.
- Pawlik TM, Keyomarsi K. Role of cell cycle in mediating sensitivity to radiotherapy. *Int J Radiat Oncol Biol Phys* 2004; 59:928–42.

34. Janicke RU, Sprengart ML, Wati MR, Porter AG. Caspase-3 is required for DNA fragmentation and morphological changes associated with apoptosis. *J Biol Chem* 1998;273:9357–60.
35. Gewirtz DA. Growth arrest and cell death in the breast tumor cell in response to ionizing radiation and chemotherapeutic agents which induce DNA damage. *Breast Cancer Res Treat* 2000;62:223–35.
36. Essmann F, Engels IH, Totzke G, Schulze-Osthoff K, Janicke RU. Apoptosis resistance of MCF-7 breast carcinoma cells to ionizing radiation is independent of p53 and cell cycle control but caused by the lack of caspase-3 and a caffeine-inhibitable event. *Cancer Res* 2004;64:7065–72.
37. Zapata JM, Krajewska M, Krajewski S, et al. Expression of multiple apoptosis-regulatory genes in human breast cancer cell lines and primary tumors. *Breast Cancer Res Treat* 1998;47:129–40.
38. Simpson L, Parsons R. PTEN: life as a tumor suppressor. *Exp Cell Res* 2001;264:29–41.
39. Yu K, Toral-Barza L, Discafani C, et al. mTOR, a novel target in breast cancer: the effect of CCI-779, an mTOR inhibitor, in preclinical models of breast cancer. *Endocr Relat Cancer* 2001;8:249–58.
40. Pandolfi PP. Breast cancer: loss of PTEN predicts resistance to treatment. *N Engl J Med* 2004;351:2337–8.
41. DeGraffenried LA, Fulcher L, Friedrichs WE, Grunwald V, Ray RB, Hidalgo M. Reduced PTEN expression in breast cancer cells confers susceptibility to inhibitors of the PI3 kinase/Akt pathway. *Ann Oncol* 2004;15:1510–6.
42. Guba M, von Breitenbuch P, Steinbauer M, et al. Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor. *Nat Med* 2002;8:128–35.

Molecular Cancer Therapeutics

Targeting the Akt/mammalian target of rapamycin pathway for radiosensitization of breast cancer

Jeffrey M. Albert, Kwang Woon Kim, Carolyn Cao, et al.

Mol Cancer Ther 2006;5:1183-1189.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/5/5/1183>

Cited articles This article cites 40 articles, 15 of which you can access for free at:
<http://mct.aacrjournals.org/content/5/5/1183.full#ref-list-1>

Citing articles This article has been cited by 19 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/5/5/1183.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/5/5/1183>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.