

Chemotherapeutic selectivity conferred by selenium: a role for p53-dependent DNA repair

Joshua L. Fischer,^{1,3} Elaine M. Mihelc,^{1,3}
Karen E. Pollok,² and Martin L. Smith^{1,3}

¹Department of Microbiology and Walther Oncology Center, Indiana University Cancer Center, and ²Herman B. Wells Center for Pediatric Research, Indiana University School of Medicine; and ³Walther Cancer Institute, Indianapolis, Indiana

Abstract

Selenium in various chemical forms has been the subject of cancer chemoprevention trials, but, more recently, selenium has been used in combination with DNA-damaging chemotherapeutics. Specifically, selenium protected tissues from dose-limiting toxicity and, in fact, allowed delivery of higher chemotherapeutic doses. At the same time, selenium did not protect cancer cells. Therefore, we seek to define the genetic basis for the observed selectivity of selenium in combination chemotherapeutics. The tumor suppressor p53 is mutated in the vast majority of cancers, but is by definition wild-type in nontarget tissues such as bone marrow and gut epithelium, tissues that are often dose-limiting due to DNA damage. We used primary, low-passage mouse embryonic fibroblasts that are wild-type or null for *p53* genes to test differential effects of selenium. Seleno-L-methionine, nontoxic by itself, was used to pretreat cell cultures before exposure to UV radiation or UV-mimetic cancer chemotherapy drugs. Seleno-L-methionine pretreatment caused a DNA repair response, which protected from subsequent challenge with DNA-damaging agents. The observed DNA repair response and subsequent DNA damage protection were p53 dependent as neither was observed in p53-null cells. The data suggest that (a) p53 may be an important genetic determinant that distinguishes normal cells from cancer cells, and (b) combinatorial chemotherapeutics that act by p53-dependent mechanisms may enhance chemotherapeutic efficacy by increasing the chemotherapeutic window distinguishing cancer cells from normal cells. [Mol Cancer Ther 2007;6(1):355 – 61]

Received 8/7/06; revised 11/1/06; accepted 11/28/06.

Grant support: U.S. Department of Defense, American Cancer Society, American Institute for Cancer Research, Indiana University Cancer Center.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Martin L. Smith, Indiana University Cancer Center, Indiana University School of Medicine, Room 155, 1044 West Walnut Street, Indianapolis, IN 46202. Phone: 317-278-0225; Fax: 317-274-7592. E-mail: marlsmith@iupui.edu

Copyright © 2007 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-06-0472

Introduction

The majority of investigation with selenium has emphasized cancer chemoprevention, and there are a number of large prevention clinical trials, many focusing on prostate cancer (1). The potential role for selenium in cancer chemotherapeutics is an area that has shown significant promise in preclinical and small clinical trials, but this potential has been overshadowed by the prevention studies. Compelling preclinical work has shown that nude mice bearing human tumor xenografts that received daily seleno-L-methionine (SeMet) supplementation before and during chemotherapy better tolerated increasing doses of irinotecan. Dose escalation allowed elimination of previously chemoresistant tumors (2). A clinical trial using selenium supplementation during chemotherapy has been initiated based on these results (3). Furthermore, phase I trials have shown that SeMet can be administered in very high doses without significant toxicity (4, 5).

There have been relatively few clinical trials investigating the effect of selenium supplementation during cancer chemotherapy; nevertheless, the results have been positive. Forty-one patients undergoing cisplatin chemotherapy were randomized into two groups, and the group that received selenium showed significantly higher WBC counts on day 14 after initiation of chemotherapy (6). Furthermore, consumption of granulocyte colony-stimulating factor and volumes of blood transfusion were significantly less in the selenium-supplemented group. An ovarian cancer study was done with 62 women undergoing cisplatin and cyclophosphamide combination chemotherapy and half of the patients received selenium supplementation (7). The group that received selenium showed significantly reduced neutropenia as well as increased WBCs from the second to third chemotherapy cycle. The authors also report that with selenium supplementation, there seemed to be a significant decrease in all cited side effects: nausea, vomiting, hair loss, etc. It was also noted that serum and tissue selenium levels in the control group decreased during the chemotherapy regimen whereas levels in the study group increased. Neither of these studies observed any loss of chemotherapeutic efficacy in association with selenium supplementation.

Of the major types of DNA repair, nucleotide excision repair (NER) is the repair pathway responsible for removing bulky lesions. For example, 6-4 photoproducts and cyclobutane pyrimidine dimers caused by UV radiation are repaired by NER. Similarly, platinum-DNA adducts formed by platinum-containing cancer chemotherapeutics are repaired by NER (8, 9). NER is divided into two distinct pathways: global genomic repair and transcription-coupled repair. Both pathways have three

basic steps: recognition of the damaged lesion, excision of the lesion, and resynthesis. The pathways differ in the initial recognition step but use the same proteins for the subsequent steps.

The damage recognition step of NER is rate limiting. For global genomic repair, regulation of this step is controlled by p53. Cells that have defective p53, such as those from patients with Li-Fraumeni syndrome, have defective global genomic repair but retain proficient transcription-coupled repair (10–13). p53 regulates the rate-limiting step in global genomic repair through transcriptional control of the DNA damage recognition proteins xeroderma pigmentosum complement groups C (XPC) and E (XPE). It has been shown that p53 transcriptionally regulates p48/XPE/DDB2, and forced overexpression of p48/XPE/DDB2 enhances global genomic repair (14–17). Likewise, XPC mRNA and protein expression is increased in a p53- and DNA damage-dependent manner (18). It has also been shown that within minutes of UV irradiation, p48 and XPC proteins localize to the damaged sites and that p48 enhances XPC binding (15). Several studies highlight the analogous repair of UV-damaged DNA and damage caused by platinum chemotherapeutics. XPC^{-/-} cells are defective in the repair of cisplatin damage, and it has been shown that XPC protein is required for cisplatin damage recognition (19, 20).

A role for selenium in DNA repair was first noticed when selenium treatment was shown to enhance host cell reactivation of a UV-damaged reporter plasmid template (21). It was later shown that selenium could only modulate DNA repair in cells with normal p53 (22). Selenium protection from DNA damage requires redox factor 1 (Ref1), which interacts with p53 and reduces key p53 cysteine residues (22, 23). The selenoprotein thioredoxin reductase is also required for p53 cysteine reduction (24). A dominant-negative Ref1 mutant blocked SeMet-induced transactivation by p53 (22). The reduced conformation of p53, promoted by SeMet, induces its transcription factor activity and the transcriptional activation of proteins responsible for recognition of DNA damage. Furthermore, the subsequent results show that SeMet elevates DNA repair and protects cells from DNA damage in the absence of cell cycle arrest or apoptosis. A potential rationale for this differential activity by p53 is likely due, in part, to posttranslational effects. It has been shown that different chemical forms of selenium have different effects on p53 phosphorylation, which alter the cellular response (25–27).

Selective modulation of NER has significant implication for patients being treated with DNA-damaging chemotherapeutic agents. The following results show that bone marrow and gut epithelium exhibited enhanced DNA repair following selenium treatment. The DNA repair activity of the cancer cells was unaffected. This effect may allow patients to receive more intense treatment without exacerbating unpleasant side effects. Experiments using matched isogenic cell lines, as well as tumors and

genetically normal tissues, show that a selenium-inducible DNA repair response protects from DNA damage and is p53 dependent. Selenium treatment did not protect or increase DNA repair in p53-deficient cells.

Materials and Methods

Chemotherapeutic Drugs

Cisplatin (purchased from Sigma, St. Louis, MO) was dissolved in DMSO as a 10 mmol/L stock solution. Carboplatin was used in some experiments instead of cisplatin, and results were identical. Oxaliplatin (purchased from HandiTech, Houston, TX) was dissolved in sterile water as a 10 mmol/L stock solution. All chemotherapeutics were frozen in small aliquots and stored at -20°C . Final concentrations in tissue culture medium were as indicated. Interleukin-6 and stem cell factor were purchased from PeproTech (Rocky Hill, NJ).

Cell Lines and Treatments

Mouse embryonic fibroblasts (MEF) of wild-type and p53^{-/-} genotypes were of low passage from our frozen stocks as previously described (28). MEF were from a C57/129 genetic background. Noncancer cells were IEC6 rat gut epithelial cells (American Type Culture Collection, Rockville, MD) and primary mouse bone marrow cells (C57/129). Bone marrow cells were stimulated with interleukin-6 (200 units/mL) and stem cell factor (100 ng/mL) for 24 h, then treated with SeMet (10 $\mu\text{mol/L}$) for 15 h, followed by DNA damage by cisplatin or oxaliplatin at the concentrations indicated. Cancer cell lines of human origin A253 and FaDu were from a previous study (2, 29). Both are squamous cell carcinoma of head and neck lines and carry mutant p53 genes. FaDu carries a R248L mutant p53 allele (30), whereas A253 carries deletions in both p53 alleles (31). Xeroderma pigmentosum XPA cells defective in DNA repair served as a negative control for some experiments, as previously described (28). Cell lines were likewise treated with SeMet (10 $\mu\text{mol/L}$, 15 h) and then with DNA-damaging chemotherapeutic drugs at concentrations and durations indicated. MEF were grown in DMEM (4.5 g/L glucose) plus 10% fetal bovine serum. Other cell lines were maintained in RPMI 1640 plus 10% fetal bovine serum, except for bone marrow, which was maintained in Iscove's modified Dulbecco's medium plus 20% fetal bovine serum, interleukin-6 (200 units/mL), and stem cell factor (100 ng/mL).

Cell Survival

Cell yield was determined by thiazolyl blue assay 7 days after DNA-damaging treatments. This assay can be applied to all cell lines irrespective of their colony-forming ability, a consideration for the MEF and other primary cells, which do not form colonies. Cells were plated at $\sim 1,000$ per well in 96-well culture plates, allowed to attach for 24 h, and treated with SeMet (10 $\mu\text{mol/L}$, 15 h) and then with DNA-damaging drugs for 2 h. Drugs were removed by washing the wells in culture medium with aspiration, then medium was replaced for the 7-day duration. On day 7, 50 μL of 2 mg/mL thiazolyl blue reagent were added to each well

and plates returned to incubator for 4 h to allow formation of a blue precipitate. The amount of blue precipitate was proportional to the number of viable cells by visual inspection. Precipitates were dissolved in DMSO and quantified by a Tecan plate reader at a wavelength of 592 nm. Data were normalized to control cells that did not receive DNA damage and expressed as percent cell yield relative to untreated controls. Data were averaged from three or more independent determinations, with wells in multiples of six in each experiment. Additionally, clonogenic cell survival was determined in some data sets. Clonogenic cell survival was conducted as described (32).

Unscheduled DNA Synthesis

DNA repair synthesis or unscheduled DNA synthesis was determined as previously described (28). Cells were treated with SeMet (15 h, 10 $\mu\text{mol/L}$), then with DNA-damaging agents to induce unscheduled DNA synthesis. The prototype DNA-damaging agent was UV radiation (20 J m^{-2} , 254 nm), which served as a positive control to induce unscheduled DNA synthesis (28). XPA cells served as a negative control because they are severely defective in nucleotide excision DNA repair (NER, <1% of normal) yet they are healthy cells unless exposed to DNA damage. After UV radiation, cellular DNA was labeled with tritiated thymidine (10 $\mu\text{Ci/mL}$) in tissue culture medium for 3 h. Cisplatin (100 $\mu\text{mol/L}$) or oxaliplatin (100 $\mu\text{mol/L}$) was delivered to cells for 5 h concurrent with tritiated thymidine uptake. Cells were fixed on glass slides in ethanol, then processed for autoradiography. S-phase nuclei were strongly labeled by the tritiated thymidine and were excluded from analysis. Non-S phase nuclei,




Figure 1. SeMet treatment (10 $\mu\text{mol/L}$, 15 h) caused elevated expression of p53-dependent DNA repair proteins XPC, XPE, and Gadd45a, which compose the "DNA repair branch" of the p53 pathway. Immunoblots were conducted with wild-type and p53^{-/-} MEF. DNA repair proteins were not detected in p53^{-/-} MEF. Proliferating cell nuclear antigen (PCNA) immunoblot and Ponceau S staining served as loading controls.




Figure 2. A, SeMet treatment (10 $\mu\text{mol/L}$, 15 h) increases the rate of repair of UV-induced DNA damage. MEF were treated with SeMet, then with UV radiation (20 J m^{-2} , 254 nm), and allowed indicated times for removal of UV lesions. An antibody to 6-4 photoproducts was used to assay 6-4 photoproduct removal from genomic DNA. The rate of 6-4 photoproduct removal was enhanced by selenium in wild-type MEF, but p53^{-/-} MEF were unaffected. Points, mean of three independent determinations; bars, SD. $P < 0.04$, Wilcoxon rank-sum test. Note the slow rate of lesion removal in p53^{-/-} MEF. B, *in vivo* evidence for a DNA repair response to SeMet. Feeding of mice with 200 $\mu\text{g/d}$ SeMet \times 5 wk leads to increased DNA repair. Removal of 6-4 photoproducts was determined as in A.

primarily in the G₁ phase of the cell cycle, exhibited DNA repair synthesis (unscheduled DNA synthesis). The number of DNA repair sites per nucleus was determined, and ≥ 200 nuclei were assayed for each data set.

Results

SeMet and Protein Expression

In cells treated with SeMet overnight, p53 is reduced to its transcriptionally active conformation and induces expression of NER damage recognition factors. XPC and p48XPE proteins are the main contributors to damage recognition in NER. Wild-type and p53^{-/-} MEF were treated overnight with SeMet. The selenium-induced expression of damage recognition proteins is p53 dependent. Wild-type cells treated with SeMet had increased

expression of several proteins known to be involved in NER DNA damage recognition whereas p53^{-/-} cells showed no change in expression of these factors (Fig. 1). Proliferating cell nuclear antigen and Ponceau S staining served as loading controls.

SeMet and Repair Rate

SeMet induces expression of damage recognition factors and has been shown to protect from DNA damage. Using an antibody to 6-4 photoproducts, a prototypical UV-inducible lesion, and cells exposed to UV radiation, the rate of repair can be assayed by monitoring the persistence of damaged lesions. Following overnight SeMet treatment, cells with wild-type p53 have fewer lesions at the indicated times (Fig. 2A). Furthermore, persistence of lesions in p53^{-/-} cells is not affected. Untreated cells serve as controls. Repair rates following SeMet treatment are expressed relative to untreated controls.

To ascertain if a DNA repair response to SeMet occurs *in vivo*, mice were given 200 µg/d SeMet orally for 5 weeks. Total bone marrow cells were UV irradiated and then incubated in tissue culture for 4 h to repair. Removal of 6-4 photoproducts was determined. Repair rates by SeMet feeding are shown relative to control mice (Fig. 2B).

SeMet and Chemotherapy

Selenomethionine protects wild-type MEF from UV radiation or cisplatin (Fig. 3). p53^{-/-} MEF were not protected. Cells were pretreated with 10 µmol/L selenomethionine for 15 h before DNA-damaging treatments. Cell survival was determined after 7 days by thiazolyl blue assay. Data of cell yield after 7 days are expressed relative to controls not treated with selenomethionine and controls not treated with DNA-damaging agents. The results shown for UV radiation are similar to those previously published (22). The similar results for UV and cisplatin treatment reiterate the requirement for p53-mediated NER for both types of damage. Cisplatin concentrations were as indicated. The implication is that p53 status is a molecular determinant that mediates DNA repair by selenium. The following experiments address this possibility.

SeMet and DNA Repair

The above findings show that SeMet treatment induces expression of NER damage recognition factors and elevates the rate of repair. Furthermore, SeMet protected wild-type, but not p53^{-/-}, cells from DNA damage. The unscheduled DNA synthesis assay was used to assay DNA repair *in vitro*. The method is illustrated in Fig. 4A. Isogenic wild-type and p53-deficient MEF (Fig. 4B) were treated with selenomethionine and various DNA-damaging agents and then unscheduled DNA synthesis was evaluated. Additionally, the effect of selenomethionine on DNA repair was evaluated in primary rat gut epithelial cells (IEC6), primary murine bone marrow, and two of the human squamous cell carcinoma of the head and neck (A253 and FaDu) cell lines used for xenografts in Cao et al.'s (2) study (Fig. 4C). Cells were treated with a variety of DNA-damaging agents: UV, cisplatin, or oxaliplatin. Wild-type MEF, rat gut epithelial cells, and murine bone marrow with genetically normal p53 show a significant increase in unscheduled DNA

synthesis when treated with selenomethionine before DNA damage $P < 0.02$ (*t* test). Cells lacking functional p53 [A253 (p53 mut), FaDu (p53^{-/-}), and p53^{-/-} MEF] were unresponsive to selenomethionine and showed no increase in unscheduled DNA synthesis.

SeMet Metabolites

Besides being used as seleno-amino acids for selenoprotein synthesis, low molecular weight metabolites of selenium compounds can mediate some biological responses. We used methyl selenenic acid as a representative SeMet metabolite. Although methyl selenenic acid showed some evidence for a DNA repair response at <1 µmol/L concentration (27), apoptosis predominated at methyl selenenic acid concentrations >1 µmol/L (Fig. 5). The DNA repair and protective effect of SeMet are therefore not likely due to low molecular weight metabolites.

Discussion

Clinical trials have shown that selenium supplementation during chemotherapy may partially alleviate the




Figure 3. SeMet treatment (10 µmol/L, 15 h) promotes cell survival in wild-type, but not p53^{-/-}, MEF. MEF were treated with SeMet, then with 254-nm UV radiation (A) or cisplatin (B). Cell survival was determined after 7 d by thiazolyl blue assay. Columns, mean of three independent determinations; bars, SD. $P < 0.04$, *t* test.




Figure 4. SeMet enhanced global genomic DNA repair as unscheduled DNA synthesis in wild-type, but not $p53^{-/-}$, MEF. **A**, illustration of methods and controls. Controls (normal human fibroblasts and DNA repair-defective XPA fibroblasts) were UV irradiated (20 J m^{-2} , 254 nm) and incubated in the presence of tritiated thymidine for 3 h, during which time the tritium label was incorporated into NER repair patches. Slides were processed for autoradiography. S-phase nuclei were excluded from analysis. By definition, unscheduled DNA synthesis (UDS; or repair synthesis) is confined to G_1 and G_2 nuclei. The number of tritium grains per nucleus is a direct measure of sites of repair synthesis. Cells not treated with DNA-damaging agents showed little or no unscheduled DNA synthesis (28). **B**, MEF treated with UV radiation (20 J m^{-2} , 254 nm), cisplatin ($50 \mu\text{mol/L}$), or oxaliplatin ($1 \mu\text{mol/L}$) for 4 h concurrent with tritiated thymidine labeling. SeMet was added to the medium 15 h before DNA-damaging treatments. Shown is relative repair synthesis (SeMet treated divided by SeMet untreated for each respective sample) in G_1 nuclei; bars, SD. At least 200 nuclei were determined per data point. SeMet induced NER in wild-type MEF ($P < 0.01$, t test). SeMet did not significantly induce NER in $p53^{-/-}$ MEF. **C**, SeMet induced NER in normal mouse bone marrow and in primary rat gut epithelial cells ($P < 0.01$, t test) but did not significantly induce NER in $p53$ -mutant cancer cell lines A253 and FaDu.

dose-limiting and poor quality-of-life side effects. In two studies, selenium supplementation significantly reduced myelotoxicity, and in one study, selenium reduced other side effects attributed to the toxicity to rapidly proliferating nontarget tissues (6, 7, 33). The findings herein show that selenium supplementation elevates expression of proteins responsible for recognition of DNA damage. The increased expression of recognition factors is concomitant with an increase in the rate of DNA repair and overall DNA repair synthesis. However, all of these selenium-inducible observations are absent in a $p53$ -null background. That is, selenium did not induce expression of key NER recognition factors or alter the rate or overall level of DNA repair in the $p53$ -null cells or tumor cell lines tested. The conclusion is that selenium selectively protects genetically normal cells from DNA-damaging chemotherapeutics, while simultaneously offering no detectable protection to cells either completely lacking $p53$ or possessing only mutant $p53$. This is important considering that $p53$ is the most widespread genetic alteration in human cancer,

with as many as 70% of tumors having a mutant $p53$ phenotype. One caveat is that some cancers with wild-type $p53$ may not be ideally suited for selenium therapies.

The results suggest a potential mechanism for selenium-inducible protection from chemotherapy in the clinical trials highlighted above and in the context of chemoprevention. In the nontarget tissues, an increase in the basal levels of NER damage recognition factors following selenium supplementation promotes an increase in the basal rate of NER, which can better tolerate the additional damage from chemotherapy. The elevated DNA repair synthesis in cells from nontarget tissues in this report, combined with the data from an earlier study showing selenium enhancing cure rates of xenograft tumors in nude mice (2), supports the proposed mechanism of selectivity.

The notion that $p53$ is an important marker for differentiating tumor cells from normal cells is not new. It is important to note, however, that a safe, reliable therapy that takes advantage of this widely known fact remains to be identified. The widespread $p53$ mutations in




Figure 5. The selenium metabolite methyl selenenic acid did not induce a protective response in MEF. Rather, p53-mediated apoptosis predominated at methyl selenenic acid (MSA) concentrations $>1 \mu\text{mol/L}$. p53^{+/+} and p53^{-/-} MEF were treated with indicated concentrations of methyl selenenic acid for 4 h. Cell survival was determined after 5 d in culture. p53^{+/+} MEF were preferentially sensitive to methyl selenenic acid ($P < 0.02$, Wilcoxon rank-sum test).

human cancer should be a benchmark for developing novel therapies. However, the inherent heterogeneity of tumors and their unpredictable responses to therapeutic strategies require extensive testing of tumor tissue. Whereas cells with altered p53 should be more sensitive to agents whose damage is repaired by the p53-regulated NER pathway, tumor cells have acquired other growth advantages, which may abrogate this potential weakness (34). A typical proposal for improving chemotherapeutic efficacy attempts to sensitize tumor cells by targeting their greatest defenses (e.g., apoptotic, cell cycle, and DNA repair targets). A strategy that protects normal cells instead is perhaps more reliable. Selenium supplementation has recently been shown in clinical trials to be nontoxic at very high doses (4). In fact, ongoing trials are attempting to reach levels of at least $15 \mu\text{mol/L}$, which shows that the concentrations used in the present study are physiologically relevant (5). The results of this study present a safe potential method of improving chemotherapeutic selectivity that focuses on the genetically normal, nontarget tissues, which may be a more promising foundation for novel therapeutic strategies.

In the United States, serum selenium concentrations of $1 \mu\text{mol/L}$ are fairly common (1). At $1 \mu\text{mol/L}$ concentration, both seleno-amino acids exemplified by SeMet and metabolic by-products of SeMet exemplified by methyl selenenic acid may contribute to DNA repair (27). At concentrations exceeding $1 \mu\text{mol/L}$, such as in this study, methyl selenenic acid induced apoptosis, which would mask any DNA repair response (Fig. 5). Therefore, it is likely that DNA repair and DNA damage protection observed *in vitro* (refs. 21, 22, and this study) and *in vivo* (Fig. 2B) at selenium concentrations in the $15 \mu\text{mol/L}$ range are due to selenoproteins (e.g., thioredoxin reductase).

Note, however, that the apoptotic response evoked by methyl selenenic acid also involves p53, as p53-wild-type MEF were preferentially sensitive to methyl selenenic acid (Fig. 5). DNA repair or apoptotic responses would each be important in chemotherapy, albeit mediated by different selenium chemical forms.

References

1. Meuillet E, Stratton S, Prasad Cherukuri D, et al. Chemoprevention of prostate cancer with selenium: an update on current clinical trials and preclinical findings. *J Cell Biochem* 2004;91:443–58.
2. Cao S, Durrani FA, Rustum YM. Selective modulation of the therapeutic efficacy of anticancer drugs by selenium containing compounds against human tumor xenografts. *Clin Cancer Res* 2004;10:2561–9.
3. Fakhri MG, Pendyala L, Smith PF, et al. A phase I and pharmacokinetic study of fixed-dose selenomethionine and irinotecan in solid tumors. *Clin Cancer Res* 2006;12:1237–44.
4. Reid ME, Stratton MS, Lillico AJ, et al. A report of high-dose selenium supplementation: response and toxicities. *J Trace Elem Med Biol* 2004;18:69–74.
5. Fakhri MG, Pendyala L, Creaven PJ, Smith P, Ross ME, Rustum Y. A Phase I dose escalation study of selenomethionine (SLM) in combination with fixed dose irinotecan (CPT-11) in patients with advanced solid tumors. *Proc Am Assoc Cancer Res* 2006;47:686.
6. Hu YJ, Chen Y, Zhang YQ, et al. The protective role of selenium on the toxicity of cisplatin-contained chemotherapy regimen in cancer patients. *Biol Trace Elem Res* 1997;56:331–41.
7. Sieja K, Talerczyk M. Selenium as an element in the treatment of ovarian cancer in women receiving chemotherapy. *Gynecol Oncol* 2004;93:320–7.
8. Reed E. Platinum-DNA adduct, nucleotide excision repair and platinum based anti-cancer chemotherapy. *Cancer Treat Rev* 1998;24:331–44.
9. Li Q, Yu JJ, Mu C, et al. Association between the level of ERCC-1 expression and the repair of cisplatin-induced DNA damage in human ovarian cancer cells. *Anticancer Res* 2000;20:645–52.
10. Ford JM, Hanawalt PC. Li-Fraumeni syndrome fibroblasts homozygous for p53 mutations are deficient in global DNA repair but exhibit normal transcription-coupled repair and enhanced UV resistance. *Proc Natl Acad Sci U S A* 1995;92:8876–80.
11. Ford JM, Hanawalt PC. Expression of wild-type p53 is required for efficient global genomic nucleotide excision repair in UV-irradiated human fibroblasts. *J Biol Chem* 1997;272:28073–80.
12. Ford JM, Baron EL, Hanawalt PC. Human fibroblasts expressing the human papillomavirus E6 gene are deficient in global genomic nucleotide excision repair and sensitive to ultraviolet irradiation. *Cancer Res* 1998;58:599–603.
13. Bowman KK, Sicard DM, Ford JM, Hanawalt PC. Reduced global genomic repair of ultraviolet light-induced cyclobutane pyrimidine dimers in simian virus 40-transformed human cells. *Mol Carcinog* 2000;29:17–24.
14. Hwang BJ, Ford JM, Hanawalt PC, Chu G. Expression of the p48 xeroderma pigmentosum gene is p53-dependent and is involved in global genomic repair. *Proc Natl Acad Sci U S A* 1999;96:424–8.
15. Wakasugi M, Kawashima A, Morioka H, et al. DDB accumulates at DNA damage sites immediately after UV irradiation and directly stimulates nucleotide excision repair. *J Biol Chem* 2002;277:1637–40.
16. Fitch ME, Cross IV, Ford JM. p53 responsive nucleotide excision repair gene products p48 and XPC, but not p53, localize to sites of UV-irradiation-induced DNA damage, *in vivo*. *Carcinogenesis* 2003;24:843–50.
17. Fitch ME, Cross IV, Turner SJ, et al. The DDB2 nucleotide excision repair gene product p48 enhances global genomic repair in p53 deficient human fibroblasts. *DNA Repair (Amst)* 2003;2:819–26.
18. Adimoolam S, Ford JM. p53 and DNA damage-inducible expression of the xeroderma pigmentosum group C gene. *Proc Natl Acad Sci U S A* 2002;99:12985–90.
19. Chen Z, Xu XS, Yang J, Wang G. Defining the function of XPC protein

- in psoralen and cisplatin-mediated DNA repair and mutagenesis. *Carcinogenesis* 2003;24:1111–21.
20. You JS, Wang M, Lee SH. Biochemical analysis of the damage recognition process in nucleotide excision repair. *J Biol Chem* 2003;278:7476–85.
21. Seo YR, Sweeney C, Smith ML. Selenomethionine induction of DNA repair response in human fibroblasts. *Oncogene* 2002;21:3663–9.
22. Seo YR, Kelley MR, Smith ML. Selenomethionine regulation of p53 by a ref1-dependent redox mechanism. *Proc Natl Acad Sci U S A* 2002;99:14548–53.
23. Jayaraman L, Murthy KG, Zhu C, Curran T, Xanthoudakis S, Prives C. Identification of redox/repair protein Ref-1 as a potent activator of p53. *Genes Dev* 1997;11:558–70.
24. Seemann S, Hainaut P. Roles of thioredoxin reductase 1 and APE/Ref-1 in the control of basal p53 stability and activity. *Oncogene* 2005;24:3853–63.
25. Goel A, Fuerst F, Hotchkiss E, Boland CR. Selenomethionine induces p53 mediated cell cycle arrest and apoptosis in human colon cancer cells. *Cancer Biol Ther* 2006;5:529–35.
26. Chung HJ, Yoon SI, Shin SH, et al. p53-mediated enhancement of radiosensitivity by selenophosphate synthetase 1 overexpression. *J Cell Physiol* 2006;209:131–41.
27. Smith ML, Lancia JK, Mercer TI, Ip C. Selenium compounds regulate p53 by common and distinctive mechanisms. *Anticancer Res* 2004;24:1401–8.
28. Smith ML, Ford JM, Hollander MC, et al. P53-mediated DNA repair responses to UV-radiation: studies of mouse cells lacking p53, p21 and/or Gadd45a genes. *Mol Cell Biol* 2000;20:3705–14.
29. Fischer JL, Cao S, Durrani FA, Fakhri M, Rustum YM, Smith ML. Chemotherapeutic selectivity conferred by selenium: a role for p53-dependent DNA repair. *Proc Am Assoc Cancer Res* 2006;47:127.
30. Kim MS, Li SL, Bertolami CN, Cherrick HM, Park NH. State of p53, Rb and DCC tumor suppressor genes in human oral cancer cell lines. *Anticancer Res* 1993;13:1405–13.
31. Courtois SJ, Woodworth CD, Degreef H, Garmyn M. Early ultraviolet B-induced G₁ arrest and suppression of the malignant phenotype by wild-type p53 in human squamous cell carcinoma cells. *Exp Cell Res* 1997;233:135–44.
32. Smith ML, Chen IT, Zhan Q, O'Connor PM, Fornace AJ, Jr. Involvement of the p53 tumor suppressor gene in repair of UV-type DNA damage. *Oncogene* 1995;10:1053–9.
33. Sieja K. Selenium (Se) deficiency in women with ovarian cancer undergoing chemotherapy and the influence of supplementation with this micro-element on biochemical parameters. *Pharmazie* 1998;53:473–6.
34. Brown JM, Wouters BG. Apoptosis, p53, and tumor cell sensitivity to anticancer agents. *Cancer Res* 1999;59:1391–9.

Molecular Cancer Therapeutics

Chemotherapeutic selectivity conferred by selenium: a role for p53-dependent DNA repair

Joshua L. Fischer, Elaine M. Mihelc, Karen E. Pollok, et al.

Mol Cancer Ther 2007;6:355-361.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/6/1/355>

Cited articles This article cites 33 articles, 13 of which you can access for free at:
<http://mct.aacrjournals.org/content/6/1/355.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/6/1/355.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/6/1/355>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.