

D,L-Sulforaphane causes transcriptional repression of androgen receptor in human prostate cancer cells

Su-Hyeong Kim and Shivendra V. Singh

Department of Pharmacology and Chemical Biology, and University of Pittsburgh Cancer Institute, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania

Abstract

D,L-Sulforaphane (SFN), a synthetic analogue of cruciferous vegetable-derived L-isomer, inhibits the growth of human prostate cancer cells in culture and *in vivo* and retards cancer development in a transgenic mouse model of prostate cancer. We now show that SFN treatment causes transcriptional repression of androgen receptor (AR) in LNCaP and C4-2 human prostate cancer cells at pharmacologic concentrations. Exposure of LNCaP and C4-2 cells to SFN resulted in a concentration-dependent and time-dependent decrease in protein levels of total AR as well as Ser^{210/213}-phosphorylated AR. The SFN-mediated decline in AR protein level was accompanied by a decrease in intracellular as well as secreted levels of prostate-specific antigen, an AR-regulated gene product. The decrease in AR protein level resulting from SFN exposure was not reversed in the presence of the protein synthesis inhibitor cycloheximide. Reverse transcription-PCR analysis revealed a dose-dependent decrease in AR mRNA levels, indicating transcriptional repression of this ligand-activated transcription factor. The SFN treatment inhibited AR promoter activity as revealed by luciferase reporter assay. Synthetic androgen (R1881)-stimulated nuclear translocation of AR was markedly suppressed in the presence of SFN in both cell lines. The SFN treatment also inhibited R1881-stimulated proliferation of LNCaP cells. Naturally occurring thio analogues (iberverin, erucin, and berteroin), but not the sulfonyl analogues (cheirolin, erysolin, and alyssin sulfone), of SFN were also effective in reducing protein levels of AR in LNCaP cells. In conclusion, the present study shows for the first time that SFN treatment causes transcription-

al repression of AR and inhibition of its nuclear localization in human prostate cancer cells. [Mol Cancer Ther 2009;8(7):1946–54]

Introduction

Observational studies suggest that dietary intake of cruciferous vegetables may be inversely associated with the risk of different malignancies, including cancer of the prostate (1–4). For example, Kolonel et al. (2) observed an inverse association between intake of yellow-orange and cruciferous vegetables and the risk of prostate cancer in a multicenter case-control study. The anticarcinogenic effect of cruciferous vegetables is ascribed to organic isothiocyanates (5, 6). Broccoli is a rather rich source of the isothiocyanate compound (–)-1-isothiocyanato-(4R)-(methylsulfinyl)-butane (L-SFN). L-SFN and its synthetic analogue D,L-sulforaphane (SFN) have sparked a great deal of research interest because of their anticancer effects. For example, L-SFN and SFN were equipotent as inducers of quinone reductase activity in Hepa 1c1c7 hepatoma cells (7). L-SFN was shown to cause transcriptional induction of phase 2 enzymes in prostate cancer cells (8). L-SFN or synthetic SFN offered protection against 9,10-dimethyl-1,2-benzanthracene-induced mammary cancer in rats, azoxymethane-induced colonic aberrant crypt foci in rats, and benzo(a)pyrene-induced forestomach cancer in mice (9–11). Dietary feeding of SFN and its N-acetylcysteine conjugate inhibited the malignant progression of lung adenomas induced by tobacco carcinogens in A/J mice (12). The induction of phase 2 carcinogen-inactivating enzymes is an important mechanism in the prevention of chemically induced cancers by SFN (5, 6).

Recent studies, including those from our laboratory, have documented novel cellular responses to SFN exposure in cultured human cancer cells (13–27). The known cellular responses to SFN exposure in cultured cancer cells include cell cycle arrest, induction of apoptosis and autophagy, inhibition of histone deacetylase, protein binding, and sensitization of cells to tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis (13–27). The mechanisms by which SFN causes growth arrest and apoptosis induction have been studied extensively in human prostate cancer cells (15–17, 19–22, 26, 27). For example, we have reported a novel mechanism for SFN-induced G₂-M phase cell cycle arrest in cultured prostate cancer cells involving checkpoint kinase 2-mediated phosphorylation of cell division cycle 25C, leading to its sequestration in the cytosol (15). We also found that the SFN-induced apoptosis is selective toward prostate cancer cells and is intimately linked to the generation of reactive oxygen species (18, 19). More recently, we have shown that the SFN-induced generation of reactive oxygen species is caused by inhibition of mitochondrial respiratory chain enzymes (20). Cellular effects

Received 2/6/09; revised 4/17/09; accepted 4/29/09; published OnlineFirst 7/7/09.

Grant support: USPHS grant CA115498 awarded by the National Cancer Institute.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Shivendra V. Singh, University of Pittsburgh School of Medicine, 2.32A Hillman Cancer Center Research Pavilion, 5117 Centre Avenue, Pittsburgh, PA 15213. Phone: 412-623-3263; Fax: 412-623-7828. E-mail: singhs@upmc.edu

Copyright © 2009 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-09-0104

downstream of reactive oxygen species generation in the execution of SFN-induced apoptosis in human prostate cancer cells include activation of Bax and caspases (19, 26). Oral gavage of SFN to male nude mice significantly retarded the growth of PC-3 human prostate cancer xenografts *in vivo* (27). In addition, oral administration of SFN inhibited prostate cancer progression and pulmonary metastasis in a transgenic mouse model of prostate cancer without causing weight loss or any other side effects (28). The SFN-mediated inhibition of prostate cancer development and metastasis in transgenic mice correlated with suppression of cellular proliferation and augmentation of natural killer cell lytic activity (28).

Androgen receptor (AR), a member of the steroid receptor superfamily of ligand-activated transcription factors, is critically involved in the initiation and progression of prostate cancer (29, 30). Accordingly, novel strategies for blockade of AR signaling are desirable. Here, we report that SFN treatment decreases protein levels of AR by causing transcriptional repression. We also show that SFN treatment blocks synthetic androgen-stimulated nuclear translation of AR. Together, our preclinical observations (refs. 27, 28 and the present study) merit clinical investigation of SFN for its efficacy against human prostate cancer.

Materials and Methods

Reagents

SFN and its naturally occurring thio analogues (iberiverin, erucin, and berteroin), sulfinyl analogues (iberin and alyssin), and sulfonyl analogues (cheirolin, erysolin, and alyssin sulfone) were purchased from LKT Laboratories. Reagents for cell culture, including fetal bovine serum (FBS), were from Invitrogen; phenol red-free RPMI 1640 was from Cellgro; charcoal-dextran-stripped FBS (CSS) was from Hyclone; synthetic androgen R1881 was from Perkin-Elmer; and protein synthesis inhibitor cycloheximide and anti-actin and anti-tubulin antibodies were from Sigma. Antibody against prostate-specific antigen (PSA) was from DakoCytomation; antibody specific for detection of Ser^{210/213}-phosphorylated AR was from Imgenex; a kit for measurement of PSA levels was from R&D Systems; and a kit for quantification of cytoplasmic histone-associated apoptotic DNA fragmentation was from Roche Applied Science. Antibody against AR and poly(ADP)ribose polymerase (PARP) were from Santa Cruz Biotechnology. The Dual-Luciferase Reporter Assay kit was from Promega.

Cell Culture and Treatments

The LNCaP and C4-2 cell lines were maintained in RPMI 1640 supplemented with 10 mmol/L HEPES, 1 mmol/L sodium pyruvate, 0.2% glucose, 10% (volume for volume) FBS, and antibiotics. Before each experiment, the cells were trypsinized, washed twice with PBS, plated in phenol red-free medium containing 10% (volume for volume) CSS, and allowed to attach by incubation for 24 h at 37°C before drug treatment. This condition was followed in each experiment, except in the experiments designed to determine the effect of SFN treatment on AR protein level and cell

viability/apoptosis in LNCaP cells cultured under regular 10% FBS. A stock solution of SFN and its analogues were prepared in DMSO, and an equal volume of DMSO (final concentration, 0.1%) was added to the controls.

Immunoblotting

After treatment with DMSO (control) or the desired concentration of SFN for specified time intervals, floating and attached cells were collected and lysed as described by us previously (31). Cell lysates were cleared by centrifugation at 14,000 rpm for 20 min. The nuclear fractions from control cells and cells exposed to SFN in the absence or presence of R1881 were prepared with the use of a nuclear extraction kit from Pierce according to the manufacturer's instructions. Lysate proteins were resolved by SDS-PAGE and transferred onto a polyvinylidene fluoride membrane. The membrane was incubated with TBS containing 0.05% Tween 20 and 5% (weight for volume) nonfat dry milk, and then exposed to the desired primary antibody. After treatment with an appropriate secondary antibody, the immunoreactive bands were visualized with the use of the enhanced chemiluminescence method.

Measurement of PSA Secretion

The LNCaP and C4-2 cells were treated with DMSO (control) and 10, 20, or 40 μ mol/L SFN for 24 h. The medium was collected and used for measurement of secreted PSA levels with the use of a kit from R&D Systems as suggested by the supplier.

Reverse Transcription-PCR

Total RNA from LNCaP and C4-2 cells treated for 24 h with DMSO (control) or SFN was extracted with the use of the RNeasy mini kit (Qiagen) according to the manufacturer's protocol. The cDNA was synthesized with the use of 1 μ g of total RNA through SuperScript reverse transcriptase (Invitrogen) with Oligo dT primer. PCR was done with a specific primer (AR sense 5'-ATGGTGAGCAGAGTGCCCTA-3'; antisense 5'-GTGGTGCTGGAAGCCTCTCCT-3') with the use of the following amplification conditions: 94°C for 2 min, followed by 28 cycles of 94°C for 1 min, 57°C for 1 min, and 68°C for 1 min. Human glyceraldehyde-3-phosphate dehydrogenase primer was used as a loading control (Biomol).

Luciferase Reporter Assay

The LNCaP and C4-2 cells were transiently cotransfected with 2 μ g pARLUC plasmid and 0.2 μ g pRL-CMV plasmid with the use of FuGENE 6 transfection reagent. Twenty-four hours after transfection, the cells were treated with SFN for the desired period and harvested in reporter lysis buffer. The samples were centrifuged, and a 20- μ L aliquot was used for measurement of dual luciferase activity with the use of a luminometer. The luciferase activity was normalized against protein concentration and expressed as a ratio of firefly luciferase to *Renilla* luciferase units.

Measurement of Cell Viability and Apoptosis

The effect of SFN treatment on cell viability was determined by trypan blue dye exclusion assay essentially as described by us previously (32). Briefly, cells in 1 mL of phenol red-free medium containing 10% CSS or under regular 10% FBS conditions were plated in 12-well plates and allowed to




Figure 1. SFN treatment decreased levels of total AR protein, Ser^{210/213}-phosphorylated AR, and PSA protein in LNCaP and C4-2 cells. Immunoblotting for AR, p-AR, and PSA with the use of lysates from LNCaP and C4-2 cells treated with the indicated concentrations of SFN for 24 h (A) and 20 μmol/L SFN for the indicated time periods (B). Numbers above bands, densitometric quantitation relative to corresponding control. Immunoblotting for each protein was done at least twice with the use of independently prepared lysates, and the results were consistent.

attach by incubation for 24 h. The medium was replaced with medium containing 20 μmol/L SFN or 1 nmol/L synthetic androgen R1881, and the plates were incubated for an additional 24 h. At the end of the incubation, both floating and adherent cells were collected and used for trypan blue dye exclusion assay or cytoplasmic histone-associated DNA fragmentation assay (32).

Results

SFN Treatment Decreased Protein Levels of AR and PSA in LNCaP and C4-2 Cells

We determined the effect of SFN treatment on AR protein level by immunoblotting with the use of the androgen-

responsive human prostate cancer cell line LNCaP and its androgen-independent variant (C4-2). The C4-2 cell line, which was generated through the coculture of parental LNCaP cells with human bone fibroblasts *in vivo* in castrated male athymic mice, displays elevated PSA expression and increased anchorage-independent growth in soft agar (33, 34). As can be seen in Fig. 1A, a 24-hour exposure to SFN in LNCaP and C4-2 cells, which were cultured for 24 hours in phenol red-free medium supplemented with 10% CSS before drug exposure, resulted in a concentration-dependent decrease in AR protein levels. The SFN-mediated decline in total AR protein level was accompanied by suppression of phosphorylated AR and PSA protein levels in both cell lines (Fig. 1A). In time course kinetic studies with the use of 20 μmol/L SFN, the decline in protein levels of total AR, phospho-AR, and PSA was evident as early as 6 hours of treatment (Fig. 1B).

To determine the lowest effective dose of SFN, we carried out immunoblotting experiments with the use of lysates from LNCaP cells cultured under the conditions described for Fig. 1, and treated for 24 and 48 hours with DMSO (control) and 1, 2.5, or 5 μmol/L SFN. As shown in Fig. 2A, the SFN-mediated decline in protein levels of AR, phospho-AR, and PSA was observed even at 1 to 2.5 μmol/L concentration, especially at the 48-hour time point (Fig. 2A). Consistent with these results, SFN treatment resulted in a dose-dependent and statistically significant decrease in secreted levels of PSA into the culture medium of both cell lines (Fig. 2B). These results indicated that SFN treatment decreased the protein level of AR and suppressed its transcriptional activity as evidenced by a decline in intracellular as well as secreted levels of the AR-regulated gene product PSA.

SFN Treatment Suppressed Transcription of AR in LNCaP and C4-2 Cells

Next, we designed experiments to test whether the SFN-mediated decrease in AR protein level was caused by inhibition of protein synthesis. The SFN-mediated reduction in AR protein expression was maintained in the presence of the protein synthesis inhibitor cycloheximide in both LNCaP and C4-2 cells (Fig. 3). These results indicated that the decrease in AR protein level resulting from SFN exposure was not due to inhibition of protein synthesis.

Figure 4A depicts the effect of SFN treatment on AR message as determined by quantitative reverse transcription-PCR. Twenty-four hour exposure of LNCaP and C4-2 cells to SFN resulted in a marked and concentration-dependent decrease in AR mRNA levels in both cell lines. We determined the effect of SFN treatment on AR promoter activity with the use of luciferase reporter assay, and the results are shown in Fig. 4B. The pARLUC plasmid used in the present study (ref. 35; a generous gift from Dr. William H. Walker, University of Pittsburgh, Pittsburgh, PA) is a modification of the pAR1.1LUC plasmid containing the proximal 1,047-bp rat AR promoter fragment inserted upstream of the luciferase gene in the pGL2-enhancer vector (35). Exposure of LNCaP and C4-2 cells to SFN for 24 hours resulted in ~40% to 80% decrease in AR promoter activity. The SFN-mediated




Figure 2. SFN treatment suppressed secretion of PSA into the culture medium of LNCaP and C4-2 cells. **A**, immunoblotting for AR, p-AR, and PSA with the use of lysates from LNCaP cells treated with DMSO (control) or the indicated concentrations of SFN for 24 or 48 h. *Numbers above bands*, densitometric quantitation relative to corresponding control. **B**, levels of PSA secreted into the medium of LNCaP and C4-2 cells after 24-h treatment of the cells with the indicated concentrations of SFN. *Columns*, mean ($n = 3$); *bars*, SE. *, $P < 0.05$, significantly different compared with control by one-way ANOVA followed by Dunnett's test. Each experiment was repeated at least twice, and the results were consistent.

decrease in AR promoter activity was evident at each concentration of SFN tested in the preset study ($P < 0.05$ by one-way ANOVA followed by Dunnett's test). These results indicated that the SFN-mediated decline in AR protein level correlated with repression of its promoter activity.

SFN Treatment Inhibited Androgen-Stimulated Nuclear Translocation of AR

Because AR is a ligand-activated transcription factor, the nuclear translocation of AR is critical for its transcriptional activity (29, 30). As expected, the nuclear level of AR was markedly increased in LNCaP and C4-2 cells in the presence of 1 nmol/L R1881 (2-hour stimulation with R1881) as revealed by immunoblotting (Fig. 5A). The R1881-stimulated nuclear translocation of AR was markedly suppressed in the presence of 20 μmol/L SFN (1-hour pretreatment with SFN followed by 2-hour treatment with R1881 in the presence of SFN) in both cell lines (Fig. 5A). These results indicated that SFN was effective in blocking R1881-stimulated nuclear localization of AR.

SFN Treatment Inhibited Androgen-Stimulated Growth of LNCaP Cells

We have shown previously that SFN treatment inhibits the viability of LNCaP cells cultured under regular FBS conditions by causing apoptosis induction (26). Initially, we determined growth-inhibitory and proapoptotic response to SFN exposure (24-hour treatment) in LNCaP cells cultured under regular 10% FBS conditions or cultured for 24 hours in phenol red-free medium supplemented with 10% CSS before drug exposure. As can be seen in Supplementary Fig. S1,¹ SFN treatment resulted in growth inhibition and apoptotic DNA fragmentation in a dose-dependent manner in LNCaP cells cultured in medium supplemented with regular FBS or CSS. Next, we determined the effect of SFN treatment (24-hour exposure) on cell viability and apoptosis induction in the presence or absence of synthetic androgen R1881 with the use of LNCaP cells cultured for 24 hours in phenol red-free medium supplemented with 10% CSS. The R1881 treatment caused a nearly 2-fold increase in the number of LNCaP cells as judged by trypan blue dye exclusion assay (Fig. 5B). The viability of LNCaP cells was decreased by ~45% by 24-hour exposure to 20 μmol/L SFN. The SFN-mediated suppression of LNCaP cell viability was maintained even in the presence of R1881. Consistent with these results, 24-hour exposure of LNCaP cells to 20 μmol/L SFN resulted in increased cytoplasmic histone-associated apoptotic DNA fragmentation compared with DMSO-treated control in the presence of R1881 (Fig. 5C). These results indicated that SFN treatment inhibited R1881-stimulated growth of LNCaP cells by causing apoptosis.

We designed experiments to gain insights into the kinetics of SFN-mediated decline in AR and PSA protein levels and apoptosis induction with the use of LNCaP cells (cultured in medium supplemented with 10% FBS) by immunoblotting for PARP. As can be seen in Fig. 5D, the SFN-mediated

¹ Supplementary material for this article is available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

cleavage of PARP (an indicator of apoptosis) was not readily evident until 16 hours of treatment, whereas the decline in AR protein level resulting from SFN exposure was obvious at the 6-hour time point.

Structure-Activity Relationship for Effect of SFN Analogues on AR Protein Level

We used naturally occurring thio, sulfinyl, and sulfonyl analogues of SFN to determine the possible impact of the oxidation state of sulfur and alkyl chain length on thioalkyl-isothiocyanate-mediated decline in AR protein level (please refer to Supplementary Table S1 for the chemical structures of the SFN analogues). As can be seen in Fig. 6, the thio derivatives of SFN (iberverin, erucin, and berteroin) were highly effective in suppressing AR protein level regardless of the alkyl chain length. For example, the propyl, butyl, and pentyl-thio analogues were more or less equally effective in reducing the AR protein level (Fig. 6). On the other hand, the sulfonyl derivatives of SFN (cheirolin, erysolin, and alysin sulfone) were either inactive or comparatively less effective than thio or sulfinyl derivatives in suppressing the AR protein level. A decline in the AR protein level was also observed in cells exposed to other sulfinyl analogues (iberin and alyssin). These results suggested that the oxidation state of the sulfur, but not the alkyl chain length, might be a critical structural determinant in thioalkyl-isothiocyanate-mediated suppression of AR protein level.

Discussion

Prostate cancer is a leading cause of cancer-related deaths among men in the United States (36). The mechanism




Figure 3. SFN-mediated decrease in AR protein level was not rescued by protein synthesis inhibitor cycloheximide (CHX). Immunoblotting for AR with the use of lysates from LNCaP and C4-2 cells treated with 20 µmol/L SFN for the indicated time periods in the absence or presence of 10 µg/ml cycloheximide (30-min pretreatment). *Numbers above bands*, densitometric quantitation relative to corresponding control. Similar results were observed in two independent experiments.




Figure 4. SFN treatment decreased AR mRNA level in LNCaP and C4-2 cells. **A**, AR and GAPDH (loading control) mRNA levels in LNCaP and C4-2 cells treated for 24 h with DMSO (control) or the indicated concentrations of SFN. *Numbers above bands*, quantitation relative to control. **B**, promoter activity of AR, determined by luciferase reporter assay, in LNCaP and C4-2 cells treated for 24 h with DMSO (control) or the indicated concentrations of SFN. *Columns*, mean ($n = 3$); *bar*, SE. *, $P < 0.05$, significantly different compared with DMSO-treated control by one-way ANOVA followed by Dunnett's test. Each experiment was repeated twice with similar results. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

underlying the pathogenesis of prostate cancer is not fully understood, but age, race, dietary habits, and androgen secretion and metabolism are some of the risk factors associated with this malignancy (37). AR, a ligand-activated transcription factor belonging to the steroid receptor superfamily, is critically involved not only in the development and maintenance of the male reproductive organ but also in prostate cancer progression (29, 30, 38, 39). Moreover, AR is believed to be a major player in the transition from hormone-sensitive to androgen-independent prostate cancer (29, 40, 41). Hormone ablation therapy is the main treatment for early-stage prostate cancer. However, this treatment modality is palliative and often leads to incurable and highly aggressive hormone refractory disease. Consequently, novel strategies to effectively eliminate AR signaling from prostate cancer are highly desirable for the clinical management of this deadly disease. We have shown previously that SFN inhibits the growth of human prostate cancer cells in culture by causing irreversible cell cycle arrest and apoptosis induction regardless of the androgen responsiveness or the p53 status (15, 19, 26, 27). The present study extends these findings and now shows that SFN treatment decreases the protein level of AR in both androgen-responsive and androgen-independent prostate cancer cells. The SFN-mediated decline in AR protein level is not blunted in the presence of cycloheximide. Instead, the SFN-mediated down-modulation of AR protein level




Figure 5. SFN treatment inhibited R1881-stimulated nuclear translocation of AR in LNCaP and C4-2 cells. **A**, immunoblotting for AR with the use of nuclear and cytosolic fractions prepared from LNCaP and C4-2 cells after treatment with 20 μmol/L SFN (1-h pretreatment) in the absence or presence of 1 nmol/L R1881 (2-h treatment with R1881 in the presence of SFN). The blots were stripped and reprobed with anti-PARP and anti-tubulin antibodies as loading controls for nuclear and cytosolic fractions, respectively. Numbers above bands, densitometric quantitation relative to DMSO-treated control. **B**, effect of SFN treatment (24-h exposure) on viability of LNCaP cells (cultured for 24 h in medium supplemented with 10% CSS before drug treatment) in the absence or presence of 1 nmol/L synthetic androgen R1881. **C**, cytoplasmic histone-associated DNA fragmentation in LNCaP cells (cultured for 24 h in medium supplemented with 10% CSS before drug treatment) treated for 24 h with 20 μmol/L SFN in the absence or presence of 1 nmol/L R1881. Columns, mean ($n = 3$); bars, SE (**B** and **C**). Significantly different ($P < 0.05$) compared with (a) DMSO-treated control and (b) R1881-treated cells by one-way ANOVA followed by Bonferroni's multiple comparison test. **D**, immunoblotting for AR, PSA, and PARP with the use of lysates from LNCaP cells (cultured under 10% FBS conditions) treated for 6 or 16 h with DMSO (control) or the indicated concentrations of SFN. Each experiment was repeated twice, and the results were comparable.

correlates with a reduction in AR message as judged by reverse transcription-PCR and inhibition of AR promoter activity as revealed by the luciferase reporter assay. Because the AR protein down-regulating effect is conserved for thio derivatives of SFN but lost for sulfonyl analogues regardless of the alkyl chain length, it is reasonable to conclude that the oxidation state of the sulfur atom influences the anti-AR effect of SFN.

Ligand-free AR predominantly resides in the cytoplasm complexed with chaperone proteins, including Hsp90, but in a conformational state receptive to ligand binding (29). Ligand-activated regulation of gene expression by AR is achieved by its nuclear translocation, dimerization, and

binding to androgen response elements in the DNA of target genes (29, 38). The present study indicates that SFN treatment inhibits the transcriptional activity of AR. This conclusion is supported by the following observations: (a) R1881-mediated nuclear translocation of AR is markedly suppressed in the presence of SFN in both LNCaP and C4-2 cell lines; (b) SFN treatment results in a significant decrease in intracellular as well as secreted levels of AR-regulated gene product PSA; and (c) SFN treatment inhibits R1881-stimulated growth of LNCaP cells in association with apoptosis induction.

An issue highly relevant to clinical application of SFN as an effective AR modulator relates to the plasma concentration of

the agent. Although pharmacokinetic variables for SFN in humans are yet to be determined, this information is available in rodents. SFN was detectable in the plasma after 1 hour of oral gavage of rats with 50 μmol SFN (42). The maximal plasma concentration of SFN was found to be ~ 20 $\mu\text{mol}/\text{L}$ after 4 hours of oral gavage and declined with a half-life of ~ 2.2 hours (42). Hanlon et al. (43) have determined absolute bioavailability of SFN in rats treated with either a single i.v. dose of 2.8 $\mu\text{mol}/\text{kg}$ or single oral doses of 2.8, 5.6, and 28 $\mu\text{mol}/\text{kg}$ (43). The plasma profile of SFN after i.v. administration was best characterized by a two-compartment pharmacokinetic model (43). These investigators also showed that SFN was very well and rapidly absorbed with an absolute bioavailability of $\sim 82\%$, which decreased at the higher doses (43). Our own work has revealed that the maximal plasma concentration of SFN in mice orally gavaged with 1 mg SFN is ~ 19 $\mu\text{mol}/\text{L}$ (28). The maximal plasma concentration of SFN and its cysteine conjugate varied between 0.65 and 0.82 $\mu\text{mol}/\text{L}$ in healthy volunteers who consumed a test meal of broccoli soup (150 mL) containing 100 g of broccoli (44).

Lower plasma level of SFN in the human volunteer study could be attributed to differences in absorption and metabolism between humans and rodents or the presence of ingredients in the broccoli soup affecting the pharmacokinetics of SFN. Nonetheless, it is plausible that low micromolar concentrations of SFN required for AR protein suppression are achievable in humans.

Elegant work by Dou et al. has established that calpain-mediated AR degradation is intrinsic to the induction of apoptosis in prostate cancer cells in response to treatment with a variety of cancer chemotherapeutic agents, including proteasomal inhibitor, topoisomerase inhibitor, DNA-damaging agents, and docetaxel (45). We have previously determined the proapoptotic response to SFN exposure in PC-3 and LNCaP cells (26). We found that the LNCaP cell line, which is androgen-responsive and expresses wild-type p53, is relatively more sensitive to apoptosis induction by SFN compared with PC-3 cells (an androgen-independent human prostate cancer cell line lacking functional p53; ref. 26). We showed further that the differential sensitivity of LNCaP and PC-3 cells to SFN-induced apoptosis was independent of p53 status (i.e., small interfering RNA-based knockdown of p53 protein in LNCaP cells failed to significantly alter proapoptotic response to SFN) but was explained by the difference in kinetics of Bax activation resulting from SFN exposure (26). The SFN-mediated activation of Bax was much more robust in the LNCaP cell line than in PC-3 cells (26). Moreover, the SFN-mediated suppression of AR expression in the LNCaP cell line occurs earlier (~ 6 hours; present study) than the onset of apoptosis (~ 16 hours; ref. 26; Fig. 5D). Based on these observations, we conclude that the proapoptotic response to SFN is not dependent on AR signaling. However, inhibition of AR signaling as well as apoptosis induction are likely to contribute to the overall anticancer effect of SFN.

SFN is known to trigger diverse anticancer responses in cultured cancer cells and therefore qualifies as a "promiscuous" agent (13–26). Promiscuity is not unique to SFN because many known successful pharmaceutical agents (e.g., aspirin) as well as a number of promising dietary cancer chemopreventive agents (e.g., garlic constituent diallyl trisulfide) are promiscuous (46). Because the pathogenesis of cancer is complex, involving abnormalities in multiple cellular checkpoints and signal transduction pathways, promiscuity may be an advantageous attribute especially for cancer chemopreventive agents. We are reluctant in assigning preference on the relative contribution of varied cellular responses (e.g., induction of phase 2 enzymes, modulation of AR signaling, apoptosis induction, inhibition of histone deacetylase, and autophagy) to the overall cancer chemopreventive effect of SFN. However, dose-dependent distinction in some cellular responses to SFN exposure is discernible. For example, SFN-mediated induction of phase 2 enzymes (7, 8) as well as suppression of AR signaling (present study) are evident at 1 to 5 $\mu\text{mol}/\text{L}$ concentrations. On the other hand, higher concentrations of SFN (20–40 $\mu\text{mol}/\text{L}$) are required to elicit an autophagic response, which serves to protect against apoptosis (22).




Figure 6. Effects of naturally occurring analogues of SFN on protein levels of AR and/or PSA in LNCaP cell line. Immunoblotting for AR and/or PSA with the use of lysates from LNCaP cells treated for 24 h with DMSO (control) or the indicated concentrations of SFN analogues. Chemical structures of the SFN analogues are shown in Supplementary Table S1. Numbers above bands, densitometric quantitation relative to DMSO-treated control.

In conclusion, the present study reveals that, analogous to certain other diet-derived agents (47), SFN is highly effective in reducing the protein level of AR, the androgen-stimulated nuclear translocation of AR, and the transcriptional activity of AR in human prostate cancer cells. We also present evidence to indicate a critical role of the oxidation state of sulfur in the thioalkyl-isothiocyanate-mediated decline in AR protein level. Results of the present study, together with our previous preclinical observations (27, 28), merit clinical investigation to determine the efficacy of SFN against prostate cancer in humans.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Dr. William H. Walker (University of Pittsburgh, Pittsburgh, PA) for the generous gift of pARLUC plasmid, and Yan Zeng for technical assistance.

References

- Verhoeven DT, Goldbohm RA, van Poppel G, Verhagen H, van den Brandt PA. Epidemiological studies on *Brassica* vegetables and cancer risk. *Cancer Epidemiol Biomarkers Prev* 1996;5:733–48.
- Kolonel LN, Hankin JH, Whittemore AS, et al. Vegetables, fruits, legumes and prostate cancer: a multiethnic case-control study. *Cancer Epidemiol Biomarkers Prev* 2000;9:795–804.
- Zhang SM, Hunter DJ, Rosner BA, et al. Intakes of fruits, vegetables, and related nutrients and the risk of non-Hodgkin's lymphoma among women. *Cancer Epidemiol Biomarkers Prev* 2000;9:477–85.
- Ambrosone CB, McCann SE, Freudenheim JL, Marshall JR, Zhang Y, Shields PG. Breast cancer risk in premenopausal women is inversely associated with consumption of broccoli, a source of isothiocyanates, but is not modified by GST genotype. *J Nutr* 2004;134:1134–8.
- Hecht SS. Inhibition of carcinogenesis by isothiocyanates. *Drug Metab Rev* 2000;32:395–411.
- Conaway CC, Yang YM, Chung FL. Isothiocyanates as cancer chemopreventive agents: their biological activities and metabolism in rodents and humans. *Curr Drug Metab* 2002;3:233–55.
- Zhang Y, Talalay P, Cho CG, Posner GH. A major inducer of anticarcinogenic protective enzymes from broccoli: isolation and elucidation of structure. *Proc Natl Acad Sci U S A* 1992;89:2399–403.
- Brooks JD, Paton VG, Vidanes G. Potent induction of phase 2 enzymes in human prostate cells by sulforaphane. *Cancer Epidemiol Biomarkers Prev* 2001;10:949–54.
- Zhang Y, Kensler TW, Cho CG, Posner GH, Talalay P. Anticarcinogenic activities of sulforaphane and structurally related synthetic nonbornyl isothiocyanates. *Proc Natl Acad Sci U S A* 1994;91:3147–50.
- Chung FL, Conaway CC, Rao CV, Reddy BS. Chemoprevention of colonic aberrant crypt foci in Fischer rats by sulforaphane and phenethyl isothiocyanate. *Carcinogenesis* 2000;21:2287–91.
- Fahey JW, Haristoy X, Dolan PM, et al. Sulforaphane inhibits extracellular, intracellular, and antibiotic-resistant strains of *Helicobacter pylori* and prevents benzo[a]pyrene-induced stomach tumors. *Proc Natl Acad Sci U S A* 2002;99:7610–5.
- Conaway CC, Wang CX, Pittman B, et al. Phenethyl isothiocyanate and sulforaphane and their *N*-acetylcysteine conjugates inhibit malignant progression of lung adenomas induced by tobacco carcinogens in A/J mice. *Cancer Res* 2005;65:8548–57.
- Gamet-Payraastre L, Li P, Lumeau S, et al. Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res* 2000;60:1426–33.
- Jackson SJT, Singletary KW. Sulforaphane: a naturally occurring mammary carcinoma mitotic inhibitor, which disrupts tubulin polymerization. *Carcinogenesis* 2004;25:219–27.
- Singh SV, Herman-Antosiewicz A, Singh AV, et al. Sulforaphane-induced G₂/M phase cell cycle arrest involves checkpoint kinase 2 mediated phosphorylation of Cdc25C. *J Biol Chem* 2004;279:25813–22.
- Myzak MC, Hardin K, Wang R, Dashwood RH, Ho E. Sulforaphane inhibits histone deacetylase activity in BPH-1, LNCaP and PC-3 prostate epithelial cells. *Carcinogenesis* 2006;27:811–9.
- Cho SD, Li G, Hu H, et al. Involvement of c-Jun N-terminal kinase in G₂/M arrest and caspase-mediated apoptosis induced by sulforaphane in DU145 prostate cancer cells. *Nutr Cancer* 2005;52:213–24.
- Choi S, Singh SV. Bax and Bak are required for apoptosis induction by sulforaphane, a cruciferous vegetable derived cancer chemopreventive agent. *Cancer Res* 2005;65:2035–43.
- Singh SV, Srivastava SK, Choi S, et al. Sulforaphane-induced cell death in human prostate cancer cells is initiated by reactive oxygen species. *J Biol Chem* 2005;280:19911–24.
- Xiao D, Powolny AA, Antosiewicz J, et al. Cellular responses to dietary cancer chemopreventive agent D,L-sulforaphane in human prostate cancer cells are initiated by mitochondrial reactive oxygen species. *Pharm Res* 2009;26:1729–38.
- Xu C, Shen G, Chen C, Gelinis C, Kong AN. Suppression of NF- κ B and NF- κ B-regulated gene expression by sulforaphane and PEITC through I κ B α , IKK pathway in human prostate cancer PC-3 cells. *Oncogene* 2005;24:4486–95.
- Herman-Antosiewicz A, Johnson DE, Singh SV. Sulforaphane causes autophagy to inhibit release of cytochrome c and apoptosis in human prostate cancer cells. *Cancer Res* 2006;66:5828–35.
- Kim H, Kim EH, Eom YW, et al. Sulforaphane sensitizes tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-resistant hepatoma cells to TRAIL-induced apoptosis through reactive oxygen species-mediated up-regulation of DR5. *Cancer Res* 2006;66:1740–50.
- Pledgie-Tracy A, Sobolewski MD, Davidson NE. Sulforaphane induces cell type-specific apoptosis in human breast cancer cell lines. *Mol Cancer Ther* 2007;6:1013–21.
- Mi L, Wang X, Govind S, et al. The role of protein binding in induction of apoptosis by phenethyl isothiocyanate and sulforaphane in human non-small lung cancer cells. *Cancer Res* 2007;67:6409–16.
- Choi S, Lew KL, Xiao H, et al. D,L-Sulforaphane-induced cell death in human prostate cancer cells is regulated by inhibitor of apoptosis family proteins and Apaf-1. *Carcinogenesis* 2007;28:151–62.
- Singh AV, Xiao D, Lew KL, Dhir R, Singh SV. Sulforaphane induces caspase-mediated apoptosis in cultured PC-3 human prostate cancer cells and retards growth of PC-3 xenografts *in vivo*. *Carcinogenesis* 2004;25:83–90.
- Singh SV, Warin R, Xiao D, et al. Sulforaphane inhibits prostate carcinogenesis and pulmonary metastasis in TRAMP mice in association with increased cytotoxicity of natural killer cells. *Cancer Res* 2009;69:2117–25.
- Burnstein KL. Regulation of androgen receptor levels: implications for prostate cancer progression and therapy. *J Cell Biochem* 2005;95:657–69.
- Kaarbø M, Klock TI, Saatcioglu F. Androgen signaling and its interactions with other signaling pathways in prostate cancer. *BioEssays* 2007;29:1227–38.
- Xiao D, Srivastava SK, Lew KL, et al. Allyl isothiocyanate, a constituent of cruciferous vegetables, inhibits proliferation of human prostate cancer cells by causing G₂/M arrest and inducing apoptosis. *Carcinogenesis* 2003;24:891–7.
- Xiao D, Choi S, Johnson DE, et al. Diallyl trisulfide-induced apoptosis in human prostate cancer cells involves c-Jun N-terminal kinase and extracellular-signal regulated kinase-mediated phosphorylation of Bcl-2. *Oncogene* 2004;23:5594–606.
- Thalmann GN, Anezinis PE, Chang SM, et al. Androgen-independent cancer progression and bone metastasis in the LNCaP model of human prostate cancer. *Cancer Res* 1994;54:2577–81.
- Wu HC, Hsieh JT, Gleave ME, Brown NM, Pathak S, Chung LW. Derivation of androgen-independent human LNCaP prostatic cancer cell sublines: role of bone stromal cells. *Int J Cancer* 1994;57:406–12.
- Delfino FJ, Boustead JN, Fix C, Walker WH. NF- κ B and TNF- α stimulate androgen receptor expression in Sertoli cells. *Mol Cell Endocrinol* 2003;201:1–12.
- Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57:43–66.

1954 *Transcriptional Repression of Androgen Receptor by SFN*

37. Nelson WG, DeMarzo A, Isaacs WB. Prostate cancer. *N Engl J Med* 2003;349:366–81.
38. Richter E, Srivastava S, Dobi A. Androgen receptor and prostate cancer. *Prostate Cancer Prostatic Dis* 2007;10:114–8.
39. Chang C, Saltzman A, Yeh S, et al. Androgen receptor: an overview. *Crit Rev Eukaryot Gene Exp* 1995;5:97–125.
40. Feldman BJ, Feldman D. The development of androgen-independent prostate cancer. *Nat Rev Cancer* 2001;1:34–45.
41. Tamura K, Furihata M, Tsunoda T, et al. Molecular features of hormone-refractory prostate cancer cells by genome-wide gene expression profiles. *Cancer Res* 2007;67:5117–25.
42. Hu R, Hebbar V, Kim BR, et al. *In vivo* pharmacokinetics and regulation of gene expression profiles by isothiocyanate sulforaphane in the rat. *J Pharmacol Exp Ther* 2004;310:263–71.
43. Hanlon N, Coldham N, Gielbert A, et al. Absolute bioavailability and dose-dependent pharmacokinetic behavior of dietary doses of the chemopreventive isothiocyanate sulforaphane in rat. *Br J Nutr* 2008;99:559–64.
44. Al Janobi AA, Mithen RF, Gasper AV, et al. Quantitative measurement of sulforaphane, iberin and their mercapturic acid pathway metabolites in human plasma and urine using liquid chromatography-tandem electrospray ionisation mass spectrometry. *J Chromatogr B (Analyt Technol Life Sci)* 2006;844:223–34.
45. Yang H, Murthy S, Sarkar FH, Sheng S, Reddy GP, Dou QP. Calpain-mediated androgen receptor breakdown in apoptotic prostate cancer cells. *J Cell Physiol* 2008;217:569–76.
46. Stan SD, Kar S, Stoner GD, Singh SV. Bioactive food components and cancer risk reduction. *J Cell Biochem* 2008;104:339–56.
47. Basak S, Pookot D, Noonan EJ, Dahiya R. Genistein down-regulates androgen receptor by modulating HDAC6-90 chaperone function. *Mol Cancer Ther* 2008;7:3195–202.

Molecular Cancer Therapeutics

d,l-Sulforaphane causes transcriptional repression of androgen receptor in human prostate cancer cells

Su-Hyeong Kim and Shivendra V. Singh

Mol Cancer Ther 2009;8:1946-1954. Published OnlineFirst July 7, 2009.

Updated version Access the most recent version of this article at:
doi:[10.1158/1535-7163.MCT-09-0104](https://doi.org/10.1158/1535-7163.MCT-09-0104)

Supplementary Material Access the most recent supplemental material at:
<http://mct.aacrjournals.org/content/suppl/2021/03/11/1535-7163.MCT-09-0104.DC1>

Cited articles This article cites 47 articles, 22 of which you can access for free at:
<http://mct.aacrjournals.org/content/8/7/1946.full#ref-list-1>

Citing articles This article has been cited by 10 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/8/7/1946.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/8/7/1946>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.