

Review

The role of vascular endothelial growth factor SNPs as predictive and prognostic markers for major solid tumors

Lokesh Jain,^{1,2} Craig A. Vargo,¹ Romano Danesi,³ Tristan M. Sissung,⁴ Douglas K. Price,⁴ David Venzon,⁵ Jürgen Venitz,² and William D. Figg^{1,2,4}

¹Clinical Pharmacology Program, National Cancer Institute, Bethesda, Maryland; ²Department of Pharmaceutics, School of Pharmacy, Virginia Commonwealth University, Richmond, Virginia; ³Department of Internal Medicine, University of Pisa, Pisa, Italy; ⁴Molecular Pharmacology Section and ⁵Biostatistics and Data Management Section, National Cancer Institute, Bethesda, Maryland

Abstract

Angiogenesis is crucial for development and metastasis of tumors, and vascular endothelial growth factor (VEGF) is a key mediator of this process. The importance of VEGF in tumorigenesis and tumor progression makes it an attractive target for the development of anticancer therapies. Inhibition of angiogenesis has shown promising clinical efficacy; however, not all patients treated with antiangiogenic agents derive benefit from them. Some patients are predisposed to refractory disease, whereas others develop resistance after initial response. Patients may also have different severity of drug-related adverse events. Optimization of drug administration based on disease status and individual responsiveness is important in limiting the treatment failure and minimization of side-effects. Single nucleotide polymorphisms (SNP) in *VEGF* may alter VEGF protein concentrations, influence the process of angiogenesis, and may relate to interindividual variation in the risk and progression of selected tumors, and their resistance to treatments. This review examines the role of SNPs in the *VEGF* gene as predictive and prognostic markers for major solid tumors, including the breast, non-small cell lung, co-

lorectal, and prostate cancers. Selected *VEGF* SNPs seem to be associated with risk of these cancers; however, there is lack of unanimity in findings, in part influenced by differences in study design and analysis. [Mol Cancer Ther 2009;8(9):2496–508]

Introduction

Antiangiogenic therapies have improved the clinical outcome of selected tumors by inhibiting tumor growth via restriction of new blood vessel formation. As the tumor grows, the lack of sufficient blood supply creates a hypoxic environment, stimulating the release of factors like hypoxia inducible factor (HIF)-1 α , which induces angiogenesis by activating the transcription of vascular endothelial growth factor (VEGF; ref. 1). VEGF and its receptors are frequently overexpressed in human tumors, including the breast, non-small cell lung, colorectal, and prostate cancers (2–5). VEGF and associated signaling pathways have been the target for many newly developed anticancer drugs. These agents have shown promising efficacy; however, they are often very expensive, and some patients experience drug resistance, limited activity, and severe toxicities. There is a strong need for identification of markers enabling *a priori* selection of patients who are likely to benefit from these agents. These markers might also carry prognostic value and may help in decisions about the overall treatment approach to manage more or less aggressive tumors with varying degrees of angiogenic involvement. This review evaluates the role of *VEGF* single nucleotide polymorphisms (SNP) in predicting the cancer susceptibility, progression, and anti-VEGF therapeutic response in subjects with major solid tumors including breast, non-small cell lung, colorectal, and prostate cancers. For these tumors anti-VEGF or anti-VEGFR treatments are either approved or in advance stages of clinical development. We also did a meta-analysis, using the methods of DerSimonian and Laird (6) to test the homogeneity of odds ratios (OR) across studies, and for calculation of pooled ORs and their confidence intervals (CI). Standard error estimates for individual ORs were calculated using exact methods from published genotype frequencies or were derived from published CIs.

VEGF Gene

The *VEGF* gene is located on chromosome 6 at location 6p21.1 (7). Its coding region spans approximately 14 kilobases and consists of eight exons (8, 9). Numerous SNPs in the promoter, 5'-, and 3'-untranslated regions (UTR) are present in *VEGF*. Some of the more frequent SNPs are

Received 4/1/09; revised 7/10/09; accepted 7/21/09; published OnlineFirst 9/15/09.

Grant support: Intramural Research Program of the National Cancer Institute (NCI), National Institutes of Health (NIH), U.S. Department of Health and Human Services.

Note: Supplementary material for this article is available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

Requests for reprints: William D. Figg, Medical Oncology Branch, CCR, NCI/NIH, 9000 Rockville Pike, Building 10, Room 5A01, Bethesda, MD, 20892. Phone: 301-402-3622; Fax: 301-402-8606. E-mail: wdfigg@helix.nih.gov

Copyright © 2009 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-09-0302




Figure 1. Structure of *VEGF* gene and position of VEGF SNPs relative to translation start site. *Italic* positions indicate alternate names. *Dashed lines* indicate the region consisting of coding sequence (CDS) and seven introns (adapted from ref. 29).

shown in Fig. 1, and their frequencies in control population are summarized in Table 1. The 5'- and 3'-UTR contain key regulatory elements that are sensitive to hypoxia (10), and contribute to high variability in VEGF production among tissues (11). For example, 634G > C SNP in the 5'-UTR of *VEGF* affects the protein translation efficiency (12), and 936C > T SNP in the 3'-UTR influences the circulating plasma concentrations (13) and tumor tissue expression of VEGF (14). However, it is likely that only a small number of these polymorphisms and haplotypes (linearly linked SNPs) actually have a functional effect on VEGF translation, whereas others act as proxies (15).

VEGF SNPs in Breast Cancer

VEGF has shown to be important for development, invasiveness, and metastasis of breast cancer in both preclinical and clinical settings (16). Higher levels of VEGF have shown to be related with adverse prognosis, decreased overall survival, and resistance to hormonal therapy (17–19). High microvessel density (MVD), a marker of VEGF expression and local activity, increases the likelihood of metastatic disease (20) and acts as a prognostic indicator for relapse-free and overall survival in node-negative breast cancer patients (21). These data suggest that SNPs in *VEGF* may play a substantial role in development and progression of breast cancer; studies evaluating this hypothesis are summarized in Table 2.

A case-control study, with 500 subjects in each group, evaluating the role of the germline *VEGF* 936C > T SNP in Austrian individuals, found that subjects with variant 936T allele were at reduced risk of breast cancer (22). However, the 936C > T SNP was not related to prognostic factors such as tumor size, histological grades, lymph node status, estrogen and progesterone receptor status, or age at diagnosis (22). In a small subset of 21 healthy, nonsmoking postmenopausal women, the 936T allele was associated with significantly lower plasma concentrations of VEGF, which may partially explain the reduced risk of cancer associated with this allele (22). A smaller study from the UK, including 144 cases and 263 controls, evaluated the effect of SNPs in the promoter regions of various cytokines, including the *VEGF* -1154G > A transition, on risk and prognosis of breast cancer. The -1154G > A allele was not associated with risk of

breast cancer; however, carriers of heterozygous -1154AG genotype displayed a trend toward favorable prognosis (23). A larger case-control study including 571 patients with familial breast cancer from Germany and Poland, and 974 patients with sporadic breast cancer from Sweden failed to find a relation between *VEGF* SNPs (-2578C > A, -1154G > A, -634G > C, and 936C > T) or their haplotypes and risk of breast cancer (24). This finding is in contrast with the earlier report (22) suggesting the protective role of 936T allele. However, patients with the -634CC genotype and -2578C/-634 C haplotype had larger, more histologically advanced tumors, whereas -2578AA and -2578A/-634G had low histological grade tumors, suggesting that these SNPs might act as genetic marker for tumor aggressiveness (24).

Two different publications (25, 26) evaluated data from the Shanghai Breast Cancer Study, a large population-based case-control study to evaluate the impact of germline *VEGF* -1498C > T, -634G > C, and 936C > T SNPs on survival and risk of breast cancer in Chinese women. On the basis of data from 1,133 cases and 1,233 age-matched controls, the -1498 CC allele was associated with shorter overall and disease-free survival [age-adjusted hazard ratio (HR) of 1.5 for both; ref. 25], but it had no association with risk of breast cancer (26). The carriers of the -634GG allele had significantly shorter overall survival than noncarriers with an age-adjusted HR of 1.6, but the disease-free survival and risk were not different between these groups (25, 26); whereas previously the complementary allele -634CC was shown to be related with advanced tumors (24). The 936C > T SNP was not related to overall or disease-free survival (25), but the 936TT genotype carriers were at reduced risk of cancer (OR = 0.65; ref. 26), agreeing with Krippel and colleagues (22) despite differences in ethnicity. The subset analysis based on menopausal status explained that the protective role of the 936TT genotype was restricted to premenopausal women (OR = 0.45). Haplotype analysis showed that -1498T/-634C/936C was related with longer overall survival (HR = 0.57), and premenopausal women with -1498T/-634G/936T were at reduced risk of breast cancer (OR = 0.47) compared with individuals with the reference -1498C/-634G/936T haplotype (25, 26).

A study evaluating the impact of four previously studied (24) *VEGF* SNPs (-2578C > A, -1154G > A, -634G > C, and 936C > T) in a Caucasian population from a different

Table 1. Frequencies of selected VEGF polymorphisms in control subjects

Polymorphism	Ethnicity	Controls	Variants			Allele Frequency		Reference No.
			No. (Percent Genotype Frequency)			P	q	
936 C > T (rs3025039)			CC	CT	TT			
	Austrian Caucasian	500	353 (70.6)	137 (27.4)	10 (2.0)			(22)
	Polish	422	297 (70.4)	114 (27.0)	11 (2.6)			(24)
	German	163	128 (78.5)	31 (19.0)	4 (2.5)			(24)
	Swedish	934	720 (77.1)	203 (21.7)	11 (1.2)			(24)
	Chinese	1,195	793 (66.4)	351 (29.4)	51 (4.3)			(26)
	UK Caucasian	708	531 (75.0)	165 (23.3)	12 (1.7)			(28)
	98% Caucasian	479	353 (73.7)	118 (24.6)	8 (1.7)			(27)
	NR	60	57 (95.0)	3 (5.0)	0 (0.0)			(30)
	Polish Caucasian	290	202 (69.7)	81 (27.9)	7 (2.4)			(32)
	Caucasian	1,458	1,125 (77.2)	308 (21.1)	25 (1.7)			(39)
	Korean	229	169 (73.8)	57 (24.9)	3 (1.3)			(43)
	Korean	413	252 (61.0)	149 (36.1)	12 (2.9)			(45)
	Austrian	427	308 (72.1)	108 (25.3)	11 (2.6)			(48)
	Tunisian	100	72 (72.0)	27 (27.0)	1 (1.0)			(55)
	Austrian	804				0.85	0.15	(29)
	-1154G > A (rs1570360)			GG	GA	AA		
NR		263	120 (45.6)	109 (41.4)	34 (12.9)			(23)
Polish		423	199 (47.0)	178 (42.1)	46 (10.9)			(24)
German		163	76 (46.6)	69 (42.3)	18 (11.0)			(24)
98% Caucasian		492	208 (42.3)	227 (46.1)	57 (11.6)			(27)
UK Caucasian		263	120 (45.6)	109 (41.4)	34 (12.9)			(53)
Tunisian		100	36 (36.0)	50 (50.0)	14 (14.0)			(55)
-634 G > C (rs2010963)			GG	GC	CC			
	Swedish	941	492 (52.3)	367 (39.0)	82 (8.7)			(24)
	98% Caucasian	500	232 (46.4)	221 (44.2)	47 (9.4)			(27)
	Korean	413	106 (25.7)	223 (54.0)	84 (20.3)			(45)
	Austrian	430	192 (44.7)	195 (45.4)	43 (10.0)			(48)
	Tunisian	100	44 (44.0)	46 (46.0)	10 (10.0)			(55)
	Chinese	1,198	418 (34.9)	598 (49.9)	182 (15.2)			(26)
	UK Caucasian	498	209 (42)	225 (45.2)	64 (12.9)			(28)
	Caucasian	1,458	650 (44.6)	644 (44.2)	164 (11.3)			(39)
	Austrian	804				0.66	0.34	(29)
-2578 C > A (rs699947)			CC	CA	AA			
	Polish	423	106 (25.1)	207 (48.9)	110 (26.0)			(24)
	German	162	50 (30.9)	72 (44.4)	40 (24.7)			(24)
	Swedish	940	257 (27.3)	451 (48.0)	232 (24.7)			(24)
	98% Caucasian	495	129 (26.1)	236 (47.7)	130 (26.3)			(27)
	Korean	203	106 (52.2)	82 (40.1)	15 (7.4)			(44)
	Austrian	427	85 (19.9)	238 (55.7)	104 (24.4)			(48)
	Austrian	804				0.53	0.47	(29)
-1498 C > T (rs833061)			CC	CT	TT			
	Chinese	1,222	78 (6.4)	479 (39.2)	665 (54.4)			(26)
	UK Caucasian	498	106 (21.3)	251 (50.4)	141 (28.3)			(28)
	Caucasian	1,458	342 (23.5)	694 (47.6)	422 (28.9)			(39)
	Taiwanese	119	4 (3.4)	72 (60.5)	43 (36.1)			(54)
	Turkish	157	50 (31.9)	94 (59.9)	13 (8.3)			(56)
	Japanese	252	23 (9.1)	97 (38.5)	132 (52.4)			(51)
	Austrian	804				0.54	0.46	(29)
-7 C > T (rs25648)			CC	CT	TT			
	UK Caucasian	493	332 (67.3)	151 (30.7)	10 (2.0)			(28)
-2489C > T (rs1005230)	Austrian	804				0.84	0.16	(29)
	Austrian	804				0.53	0.47	(29)
1612G > A (rs10434)			GG	GA	AA			
	Austrian	804				0.54	0.46	(29)

Abbreviation: NR, not reported.

Table 2. Summary of case-control studies assessing the VEGF polymorphisms in relation to risk of breast cancer

Polymorphism	Ethnicity	Case-Control	Genotype	Association of Cancer Risk with Respect to VEGF Genotype	OR	OR Adjusted (Yes or No)	L95% CI	U95% CI	Reference No.
936 C > T*	Austrian Caucasian	500/500	CC vs. (CT or TT) [†]	Yes (decreases risk)	0.51	No	0.38	0.7	(22)
			TT vs. CC [†]	No	0.77	No	0.29	2.03	
	Polish	412/422	CT vs. CC	No	0.87	No	0.63	1.21	(24)
			TT vs. CC	No	1.27	No	0.53	3.05	
			CC vs. (CT or TT) [†]	No	0.91	No	0.67	1.23	
	German	153/163	CT vs. CC	No	1.07	No	0.59	1.93	(24)
			TT vs. CC	No	0.53	No	0.07	3.46	
			CC vs. (CT or TT) [†]	No	1.01	No	0.58	1.75	
	Swedish	924/934	CT vs. CC	No	1.02	No	0.81	1.28	(24)
			TT vs. CC	No	1.11	No	0.45	2.71	
			CC vs. (CT or TT) [†]	No	1.03	No	0.82	1.28	
	Chinese	1,109/1,195	CT vs. CC	No	1.01	No	0.85	1.21	(26)
			TT vs. CC	Yes (decreases risk)	0.65	No	0.41	1.02	
			CC vs. (CT or TT) [†]	No	0.97	No	0.81	1.15	
	Caucasian (98%)	488/479	CC vs. (CT or TT)	No	1.02	Yes	0.76	1.36	(27)
TT vs. CC [†]			No	1.42	No	0.67	3.07		
UK Caucasian	848/708	CC vs. (CT or TT) [†]	No	1.08	No	0.85	1.36	(28)	
		TT vs. CC [†]	No	0.82	No	0.26	2.6		
Polish Caucasian	319/290	CC vs. (CT or TT)	Yes (decreases risk)	0.69	No	0.48	0.99	(32)	
		TT vs. CC [†]	No	0.82	No	0.26	2.6		
NR	60/60	CC vs. (CT or TT) [†]	Yes (increases risk)	6.91	Yes	1.84	32	(30)	
		TT vs. CC [†]	No	0.82	No	0.26	2.6		
-2578 C > A [‡]	Polish	411/423	CA vs. CC	No	0.96	No	0.68	1.36	(24)
			AA vs. CC	No	1.04	No	0.7	1.54	
			AA vs. (CA or CC) [†]	No	1.07	No	0.78	1.46	
	German	153/162	CA vs. CC	No	1.18	No	0.68	2.06	(24)
			AA vs. CC	No	0.97	No	0.5	1.86	
			AA vs. (CA or CC) [†]	No	0.87	No	0.51	1.49	
	Swedish	939/940	CA vs. CC	No	0.99	No	0.79	1.24	(24)
			AA vs. CC	No	1	No	0.77	1.29	
			AA vs. (CA or CC) [†]	No	1	No	0.81	1.24	
	98% Caucasian 89% Caucasian + 7% AA	498/495 NR (total 656)	CA vs. AA	No	1.18	Yes	0.86	1.61	(27)
			AA vs. CC [§]	No	0.8	Yes	0.56	1.14	
	NR	134/263	AA vs. (CA or CC)	Yes (increases risk)	1.99	Yes	1.06	3.74	(31)
			AA vs. GG [†]	No	0.61	No	0.27	1.3	
	Polish	411/423	GA vs. GG	No	1.22	No	0.91	1.65	(24)
			AA vs. GG	No	1.23	No	0.76	1.97	
GA vs. GG			No	0.76	No	0.46	1.25		
German	159/163	AA vs. GG	No	0.85	No	0.39	1.89	(24)	
		GA vs. AA	No	1.01	Yes	0.66	1.55		
98% Caucasian	495/492	AA vs. GG [§]	No	0.78	Yes	0.51	1.19	(27)	
		AA vs. GG [§]	No	0.78	Yes	0.51	1.19		

(Continued on the following page)

Table 2. Summary of case-control studies assessing the VEGF polymorphisms in relation to risk of breast cancer (Cont'd)

Polymorphism	Ethnicity	Case-Control	Genotype	Association of Cancer Risk with Respect to VEGF Genotype	OR	OR Adjusted (Yes or No)	L95% CI	U95% CI	Reference No.
-634 G > C [¶]	Chinese	1,095/1,198	CG vs. CC	No	0.81	No	0.64	1.02	(26)
			CC vs. GG [§]	No	1.12	No	0.88	1.44	
	UK Caucasian	490/498	CG vs. CC [†]	No	1.03	No	0.68	1.56	(28)
			CC vs. GG [†]	No	0.82	No	0.54	1.26	
	Swedish	936/941	GC vs. GG [†]	No	0.85	No	0.65	1.12	
			GC vs. GG	No	1	No	0.82	1.21	(24)
			CC vs. GG	No	1.05	No	0.74	1.47	
	98% Caucasian	495/500	CG vs. CC [†]	No	0.95	No	0.68	1.34	
			GC vs. GG	No	1.05	Yes	0.8	1.37	(27)
			CC vs. GG	No	1.18	Yes	0.76	1.83	
-1498 C > T**	Chinese	1,123/1,222	CT vs. TT	No	0.94	No	0.79	1.12	(26)
			CC vs. TT	No	1.23	No	0.89	1.7	
			CC vs. (CT or TT) [†]	No	1.26	No	0.92	1.73	
	UK Caucasian	493/498	CT vs. TT [†]	No	1.04	No	0.77	1.4	(28)
			CC vs. TT [†]	No	1.1	No	0.77	1.59	
			CC vs. (CT or TT) [†]	No	1.07	No	0.8	1.45	
	89% Caucasian + 7% AA	NR (total 656)	CC vs. (CT or TT)	Yes (increases risk)	2.01	Yes	1.08	3.76	(31)
	Chinese		Ref: -1498T/-634C/936C	Yes (decreases risk)	0.67	No	0.43	1.04	(26)
-1498T/-634G/936T				No	0.97	No	0.84	1.11	
-1498T/-634G/936C				No	1.02	No	0.86	1.21	
-1498C/-634G/936C				No	0.96	No	0.78	1.18	
-1498C/-634G/936T				No	0.98	No	0.76	1.25	
-1498T/-634C/936T				No	0.98	No	0.76	1.25	
-2578A/-1154A/-634G	98% Caucasian		1 HC vs. 0	Yes (decreases risk)	0.69	Yes	0.51	0.92	(27)
			2 HC vs. 0	Yes (decreases risk)	0.62	Yes	0.38	1	
-2578A/-1154G/-634G			1 HC vs. 0	No	0.97	Yes	0.71	1.33	
			2 HC vs. 0	No	1.71	Yes	0.7	4.15	
-2578C/-1154G/-634G			1 HC vs. 0	No	1.27	Yes	0.94	1.72	
			2 HC vs. 0	No	1.68	Yes	0.78	3.61	
-2578C/-1154G/-634C			1 HC vs. 0	No	1.07	Yes	0.8	1.43	

Abbreviations: HC, haplotype copy numbers; NR, not reported; L, lower; U, upper; AA, African-American; Ref, reference haplotype.

*Test for homogeneity of OR: pooled OR (95% CI); CT versus CC, $P = 0.86, 0.99 (0.88-1.13)$; TT versus CC, $P = 0.61, 0.86 (0.64-1.16)$; CC versus (CT or TT), $P = 0.0010, 0.91 (0.76-1.10)$.

[†]Calculated OR.

[‡]Test for homogeneity of OR: pooled OR (95% CI); CA versus CC, $P = 0.82, 1.00 (0.84-1.20)$; AA versus CC, $P = 0.73, 0.95 (0.80-1.13)$; AA versus (CA or CC), $P = 0.20, 1.08 (0.86-1.35)$.

[§]OR inverted from original publication.

[¶]Test for homogeneity of OR: pooled OR (95% CI); AA versus GG, $P = 0.36, 0.89 (0.67-1.18)$; GA versus GG, $P = 0.11, 1.01 (0.64-1.59)$.

^{‡‡}Test for homogeneity of OR: pooled OR (95% CI); GC versus GG, $P = 0.52, 0.97 (0.85-1.11)$; GC versus CC, $P = 0.52, 0.88 (0.74-1.05)$; CC versus GG, $P = 0.62, 1.06 (0.89-1.25)$.

**Test for homogeneity of OR: pooled OR (95% CI); CT versus TT, $P = 0.57, 0.96 (0.83-1.12)$; CC versus TT, $P = 0.66, 1.17 (0.92-1.49)$; CC versus (CT or TT), $P = 0.21, 1.23 (1.00-1.51)$.

geographic location (United States), including 501 cases and 504 controls, also failed to find any association between these SNPs and the risk of breast cancer (27), disregarding the spread of disease. However, subgroup analysis in pa-

tients with invasive ($n = 380$) and *in situ* ($n = 107$) breast cancer showed higher risk of invasive, but not of *in situ*, cancer in carriers of -2578CC (OR = 1.46) and -1154GG (1.64) genotypes (27). Conversely, the 936C allele was not

associated with invasive cancer, but was related to reduced risk of *in situ* breast cancer (OR = 0.59; ref. 27). Notably, this finding is in contrast with the majority of previous studies in which patients with the 936T allele had lower risk of breast cancer (22, 26). The -634G > C SNP had no association with risk of either *in situ* or invasive breast cancer. These results show the possible differences in genetic markers based on stage of disease (i.e., localized or metastatic). Haplotype analysis showed that one or two copies of -2578A/-1154A/-634G resulted in lower risk of invasive breast cancer compared with zero-copy carriers (OR = 0.69 and 0.62, respectively). Assessment of promoter diplotypes suggested no significant association with risk of invasive breast cancer (27). Another case-control study in 500 Caucasian patients and 500 matched controls analyzed the effect of germline VEGF -1498C > T, -634G > C, -7C > T, and 936C > T SNPs on risk and severity of breast cancer (assessed by tumor size, tumor grade, nodal involvement, vascular invasion, and estrogen receptor status), survival of patients, circulating levels of VEGF, and VEGF expression in tumor (28). Singularly, none of these SNPs were related to risk, but patients carrying the -1498T/-634C/-7C/936C haplotype were at lower risk of breast cancer than noncarriers. Except for -7C > T transition, which predicted longer overall survival, no other SNPs were related with survival (28). The serum and plasma VEGF levels and tumor VEGF expression were not related with any of these four SNPs (28). A large case-control study including 804 breast cancer patients and

804 age-matched healthy volunteers from Austria found that germline VEGF -2578C > A, -2489C > T, -1498C > T, -634G > C, -7C > T, 936C > T, and 1612G > A SNPs or their haplotypes were not related with risk of breast cancer (29). However the -634C allele was related with small tumor size (29), contrasting an earlier report (24) showing -634CC allele's relation with high tumor aggressiveness. A small case-control study with 60 patients and 60 age-matched healthy volunteers found no carriers of the 936TT genotype in two groups, and the frequency of the 936CT genotype was significantly higher in patients compared with controls (30). The 936C > T SNP had no association with clinicopathological characteristics of the patients (30).

An epidemiological study with predominantly Caucasian subjects, including 715 controls and 520 breast cancer cases, evaluated the impact of alleles in seven genes, including the VEGF -2578C > A, -1498C > T, -634G > C, -1154G > A, and 936C > T SNPs on risk of breast cancer. The -2578AA and -1498CC genotypes had higher risk of breast cancer, independent of nongenetic risk factors estimated by the Gail model (respective adjusted OR = 1.99 and 2.01), whereas other VEGF SNPs had no association with disease risk (31). Finally, a case-control study in Polish Caucasian subjects, including 319 breast cancers, 146 ovarian cancers, and 290 unaffected controls, investigated the effect of the 936C > T SNP in modifying the hereditary risk of breast and ovarian cancer in women harboring a mutated BRCA1

Table 3. Summary of case-control studies assessing the VEGF polymorphisms in relation to risk of NSCLC

Polymorphism	Ethnicity	Case-Control	Genotype	Association of Cancer Risk with Respect to VEGF Genotype	Gender	OR	L95% CI	U95% CI	Reference No.
936 C > T	Caucasian	1,900/1,458	(CT or TT) vs. CC	No	All	1.05	0.87	1.27	(39)
				No	Male	1.13	0.87	1.5	
				No	Female	0.99	0.27	1.29	
-1498 C > T	Caucasian	1,900/1,458	(CT or CC) vs. TT	No	All	1.16	0.96	1.4	(39)
				No	Male	1.07	0.81	1.4	
				No	Female	1.27	0.98	1.65	
-634 G > C	Caucasian	1,900/1,458	(CG or CC) vs. GG	No	All	1.14	0.96	1.36	(39)
				Yes (increases risk, lung adenocarcinoma)	Male	1.27	0.99	1.63	
				No	Female	1.04	0.82	1.32	
-1498T/-634C/936C	Caucasian	1,900/1,458	Ref: -1498C/-634G/936C	No	All	1.02	0.88	1.17	(39)
				No	Male	1.15	0.94	1.42	
				No	Female	0.91	0.74	1.11	
-1498T/-634G/936C	Caucasian	1,900/1,458	Ref: -1498C/-634G/936C	No	All	0.88	0.75	1.04	(39)
				Yes (decreases risk, lung adenocarcinoma)	Male	0.81	0.63	1.02	
				No	Female	0.93	0.74	1.17	
-1498C/-634G/936T	Caucasian	1,900/1,458	Ref: -1498C/-634G/936C	No	All	0.98	0.74	1.31	(39)
				No	Male	0.9	0.59	1.36	
				No	Female	1.04	0.69	1.55	
-1498T/-634C/936T	Caucasian	1,900/1,458	Ref: -1498C/-634G/936C	No	All	1.14	0.84	1.55	(39)
				No	Male	1.15	0.72	1.85	
				No	Female	1.04	0.69	1.57	

Abbreviations: L, lower; U, upper; Ref, reference haplotype.

gene (32). Carriers of the *VEGF* 936 T allele and *BRCA1* mutation had reduced risk of breast cancer (OR = 0.63 for CT+TT genotype) but the risk of ovarian cancer was not affected (32). These findings suggest a protective role for the 936T allele in women at risk for breast cancer carrying *BRCA1* mutations.

Recently two studies (33, 34) evaluated the relationship of *VEGF* plasma levels or somatic *VEGF* and *VEGFR-2* SNPs with treatment efficacy or toxicity. One of them compared the effect of paclitaxel with paclitaxel and bevacizumab combination in metastatic breast cancer by enrolling 183 patients in each arm (33). The *VEGF* -2578AA had longer median overall survival (HR 0.58) compared with the -2578CA+CC genotype in the combination arm (33). Similarly -1154A allele had longer overall survival (HR 0.62) in the combination arm, which increased additively with increase in number of A alleles (33). *VEGF* -634CC and -1498TT had significantly less likelihood of developing grade 3 or 4 hypertension in the combination arm compared with -634 GC+GG and -1498 CT+CC, respectively (33). The *VEGF* 936C > T, *VEGFR-2* 889G > A, and *VEGFR-2* 1416A > T had no significant relationship with either the efficacy or toxicity. Another study enrolled 56 breast cancer patients to examine the role of plasma *VEGF* levels as a predictor of treatment outcome with bevacizumab and vinorelbine therapy (34). Lower levels of baseline *VEGF* predicted longer progression free survival (34).

In conclusion, *VEGF* -634G > C (24, 26–29, 31), -7C > T (26, 28, 29), and -2489C > A (26, 29) SNPs were not related with risk of breast cancer. *VEGF* -1154G > A had no relation with overall risk of breast cancer (23, 24, 27, 31), however the G allele was related with increased risk of invasive disease (23, 27, 31). *VEGF* -1498C > T was shown to have no relation (26, 28, 29) or predicted higher risk of breast cancer (31). Results for *VEGF* -2578C > A were inconsistent, i.e., no association with risk (23, 24, 27, 29) and higher risk of invasive disease with the C allele (23, 27, 31) or A allele (31). For the 936C > T SNP, three studies showed a protective role of the 936T allele (22, 26, 32), four failed to find any correlation with risk (24, 28, 29, 31), and one found 936 C as a protective allele for *in situ* breast cancer (27). Differences in geographic location of the studied population did not explain the differences in the role of the 936C > T SNP. Results of meta-analysis (Table 2) show that, except for 936 CC versus (CT or TT), ORs for all other genotype comparisons were homogenous to be combined across studies. Despite a small number of studies that disagree with the others, the pooled analyses for these genotypes seem not to have found any significant associations with breast cancer risk.

VEGF SNPs in Non-Small Cell Lung Cancer

High *VEGF* mRNA or protein expression in tumor (35) or in serum have shown to be associated with advanced tumor stage (IIIB and IV; ref. 36), shorter survival (37), and prognosis of non-small cell lung cancer (NSCLC; ref. 38).

VEGF mRNA and protein expression were found to be significantly higher in adenocarcinomas than in squamous-cell carcinomas (35). These observations suggest that SNPs in the *VEGF* gene may affect the prognosis or clinical outcome of NSCLC by altering the *VEGF* levels; studies evaluating this hypothesis are summarized in Table 3.

A small study on DNA samples extracted from tumor and normal lung tissues from 36 surgically resected patients, examined the effect of *VEGF* -2578C > A, -634G > C, -1154G > A, and 936C > T SNPs on tumor *VEGF* expression and vascular density (14).

The -2578 CC, -634 GG, and -1154 AA and GA genotypes were associated with low *VEGF* expression, whereas high *VEGF* levels were detected in samples carrying the -2578 CA, -634 GC, and -1154 GG genotypes (14). The MVD in tumors bearing the -2578 CC and -634 GG genotypes was significantly lower compared with the -2578 CA and -634 CC+GC genotypes, respectively. *VEGF* -1154G > A and 936C > T were not related with tumor *VEGF* expression and MVD (14). A large case-control study on 1,900 NSCLC Caucasian patients and 1,458 healthy controls failed to find a correlation between germline *VEGF* -1498C > T, -634C > G, and 936C > T SNPs taken singularly on risk of NSCLC after adjusting for age, gender, smoking status, pack-years of smoking, and years since smoking cessation (39). However, male patients bearing the -634CC+CG genotype and -1498T/-634G/936C haplotype, respectively, had higher and reduced risk of developing lung adenocarcinoma (39). The lack of association in female patients was unclear and was not explained by differences in smoking status between genders (39). Another study on 462 patients with early-stage NSCLC treated with surgical resection, examined the impact of germline 936C > T, -1498C > T, and -634G > C SNPs on survival (3). Carriers of -634C and 936T alleles had longer 5-year overall survival than noncarriers, with respective survival of 61% versus 51%, and 67% versus 54% (3). While considering the combined effect of 936C > T and -634G > C SNPs, an increased number of pooled variant alleles resulted in significantly longer survival, with adjusted HR for one, two, and three variant allele(s) being 0.76, 0.61, and 0.61, respectively, compared with the zero variant allele (3). The -1498C > T SNP had no association with overall survival in these patients (3).

In conclusion, male subjects carrying the -634C allele seem to be at increased risk of lung adenocarcinoma (39). Considering both males and females together, -634G > C, 936C > T, and -1498C > T had no significant association with risk of disease (39), but a variant allele for these SNPs predicted longer overall survival (3). The -1498T/-634G/936C haplotype may predict the risk of NSCLC. The -2578C > A, -1154G > A, and -634G > C SNPs were associated with *VEGF* expression and angiogenesis, but these observations were limited by small sample size (14). There were not enough studies (Table 3) to do a meta-analysis.

VEGF SNPs in Colorectal Cancer

Like other solid tumors, overexpression of *VEGF* mRNA and protein has shown to be associated with tumor progression and poor prognosis of colon carcinoma (40). *VEGF* has shown to be an important predictor of prognosis in cases of advanced CRC (41). High MVD and *VEGF* expression were shown to be related with poor relapse-free and overall survival (42). Therefore, *VEGF* SNPs may play an important role in determining the risk, prognosis, and survival in colorectal cancer patients; studies evaluating this hypothesis are summarized in Table 4.

A study that analyzed *VEGF* SNPs in peripheral blood and their impact on prognosis in 125 patients with locally advanced colon cancer found that the *VEGF* 936C genotype had a significantly shorter median time to relapse compared with CT or TT genotypes (2.6 versus 11.1 years; ref. 40). A case-control study on the *VEGF* 936C > T germline SNP in Korean colon cancer patients ($n = 262$) and healthy controls ($n = 229$) showed that the frequency of T allele-bearing genotypes (CT+TT) was higher in cases than in controls. When

stratified by gender and age, the frequency of CT+TT was significantly associated with increased risk for colon cancer in women and patients younger than 55 years old, thus suggesting that the *VEGF* 936C > T SNP might be a genetic determinant for colon cancer in Korean patients (43). Another case-control study on the *VEGF* -2578C > A germline SNP was carried out in a cohort of 246 patients with colon cancer and 203 healthy controls and found that *VEGF* -2578C > A SNP had no association with susceptibility to colon cancer, although gender stratification indicated that the -2578CA and AA genotypes had a marginal protective effect for colon cancer in women. Therefore, the *VEGF* -2578C > A SNP, at least in Koreans, seems to have a borderline role in colon cancer risk (44). The analysis of two *VEGF* SNPs (-634G > C and 936C > T) in colorectal tissue from 465 patients and peripheral blood lymphocytes from 413 healthy controls showed that the -634 GC and CC genotypes were associated with a decreased risk of cancer, indicating a dominant role of the C allele, whereas the 936TT genotype correlated with advanced stage, high serum levels of CA19-9, and higher

Table 4. Summary of case-control studies assessing the *VEGF* polymorphisms in relation to risk of colon cancer

Polymorphism	Ethnicity	Case-Control	Genotype	Association of Cancer Risk with Respect to <i>VEGF</i> Genotype	OR	L95%CI	U95%CI	Reference No.
936C > T*	Korean	262/229	CT vs. CC	No	1.45	0.97	2.16	(43)
			TT vs. CC	No	2.98	0.79	11.21	
	NR	465/413	(CT or TT) vs. CC	Yes (increases risk)	1.52	1.03	2.25	(45)
			CT vs. CC	No	0.89	0.67	1.18	
-2578C > A [†]	Austrian	427/427	TT vs. CC	No	1.19	0.5	2.35	(48)
			(CT or TT) vs. CC	No	0.92	0.7	1.21	
	Korean	246/203	T vs. C	No	0.96	0.76	1.21	(44)
			CA vs. CC	No	0.76	0.55	1.05	
-634G > C [§]	Austrian	433/427	TT vs. CC [†]	No	0.68	0.27	1.7	(48)
			(CT or TT) vs. CC [†]	No	0.75	0.55	1.02	
	NR	465/413	AA vs. CC	No	0.72	0.49	1.07	(45)
			(CA or AA) vs. CC	No	0.66	0.31	1.43	
-634C/936C	Austrian	433/427	CA vs. CC [†]	No	1	0.7	1.43	(48)
			AA vs. CC [†]	No	1.31	0.88	1.95	
	NR	465/413	(CA or AA) vs. CC [†]	No	1.1	0.78	1.54	(45)
			GC vs. GG	Yes (decreases risk)	0.55	0.41	0.75	
-634G/936T	Austrian	432/430	CC vs. GG	No	0.81	0.55	1.18	(48)
			(GC or CC) vs. GG	Yes (decreases risk)	0.62	0.47	0.83	
	NR	465/413	C vs. G	No	0.86	0.71	1.04	(45)
			GC vs. GG [†]	No	0.98	0.74	1.3	
NR	465/413	CC vs. GG [†]	No	1.09	0.69	1.72	(45)	
		(GC or CC) vs. GG [†]	No	1	0.76	1.31		
-634C/936C	NR		Ref: -634G/936C	Yes (decreases risk)	0.53	0.43	0.66	(45)
-634G/936T	NR		Ref: -634G/936C	Yes (decreases risk)	0.56	0.42	0.74	

Abbreviations: NR, not reported; L, lower; U, upper; Ref, reference haplotype.

*Test for homogeneity of OR: pooled OR (95% CI); CT versus CC, $P = 0.042$, 0.98 (0.70–1.37); TT versus CC, $P = 0.20$, 1.17 (0.57–2.39); (CT or TT) versus CC, $P = 0.019$, 1.00 (0.69–1.44).

[†]Calculated OR.

[‡]Test for homogeneity of OR: pooled OR (95% CI); CA versus CC, $P = 0.22$, 0.86 (0.62–1.19); AA versus CC, $P = 0.12$, 1.01 (0.53–1.93); (CA or AA) versus CC, $P = 0.095$, 0.89 (0.58–1.36).

[§]Test for homogeneity of OR: pooled OR (95% CI); GC versus GG, $P = 0.0070$, 0.74 (0.42–1.30); CC versus GG, $P = 0.33$, 0.91 (0.68–1.22); (GC or CC) versus GG, $P = 0.018$, 0.79 (0.49–1.26).

histologic grades (45). Haplotype analysis showed that -634C/936C and -634G/936T were associated with decreased susceptibility of colorectal cancer (45). The *VEGF* -1498C > T, -634G > C, and -7C > T SNPs were determined in 36 Japanese patients and showed that the -1498CC allele, but not the -634G > C and -7C > T SNPs, was predictive of poorly differentiated tumors and thereby poor prognosis (46). A larger study on 445 patients with colorectal cancer examined the *VEGF* -2578C > A, -634G > C, and 936C > T SNPs and showed that the survival for patients with the -634GC or CC genotypes was better than for those with the -634GG genotype (47). Furthermore, the 936CT or TT genotypes were associated with worse survival compared with the 936CC genotype, and -2578C > A was not related with survival (47). Finally, in haplotype analysis, the -2578A/-634G/936T exhibited a significantly less favorable survival when compared with the wild-type -2578C/-634G/936C haplotype (47). A case-control study examined *VEGF* 936C > T, -2578C > A, and -634G > C SNPs in 427 patients and 427 age- and sex-matched healthy control subjects and found no correlation between all these variants and tumor size, histological grading, positive regional lymph node metastases, or tumor stage (48). Patients treated with cetuximab were genotyped to assess the possible predictive role of SNPs of *cyclin D1*, *cyclooxygenase 2*, *EGF*, *EGFR*, *IL-8*, and *VEGF* genes in blood samples. Although a significant association was found between *cyclin D1* 870A > G and the *EGF* 61A > G SNPs and survival, *VEGF* 936C > T showed no association (49). Finally, a study on RNA from 25 colorectal cancer tissues showed the presence of *VEGF121*, *VEGF145*, *VEGF165*, and *VEGF189* splice variants and an array of novel mutations within the conserved expression site of the gene. Five samples exhibited single nucleotide changes and in one sample a two-nucleotide deletion was detected. In addition to this, a one- and two-base deletion with frame-shift and protein truncation in exon 3 at positions +172 and +171/172 were shown, respectively, as well as a transition mutation in exon 3 at position +248, and two transition mutations in exon 4 at positions +398 and +403. The biologic and clinical impact of these variants were not shown, however (50).

In conclusion, 936 T allele has shown to be associated with increased risk (43), advanced stage of disease (45), worse survival (47), and longer time to relapse (40), whereas other studies have shown no correlation with tumor size, grade, and stage (48), and no relation with survival (49) in patients with colorectal cancer. The -634 C allele was predictive of decreased risk (45) and better survival (47), and was not related with tumor size, grade, and stage (48). The *VEGF* -2578C > A SNP was not related with risk (44), survival (47), tumor size and stage (48), but -2578 CA and AA had a borderline protective role in Korean women (44). The results of the meta-analysis shown in Table 4 suggest that ORs for several genotypic comparisons were not homogeneous to be pooled across studies (*P*-value for test of homogeneity, < 0.05). The 95% CI for all the genotypic comparisons for which ORs were

homogeneous and could be pooled together included one, suggesting no significant correlation of selected *VEGF* SNPs with risk of colorectal cancer.

VEGF SNPs in Prostate Cancer

Like some solid tumors, progressive growth and metastasis of prostate cancer is also dependent on angiogenesis. *VEGF* expression and vascular density were significantly higher in prostate cancer tissue compared with benign prostate hypertrophy (BPH) or normal prostate tissue (5, 51). Metastatic prostate cells had more enhanced *VEGF* production and tumor vascularity than cells with low metastatic potential (52), suggesting the importance of *VEGF* in prostate cancer metastasis. Studies evaluating the role of *VEGF* SNPs in prostate cancer susceptibility, progression, and metastasis are summarized in Table 5.

A case-control study on 247 cancer patients and 263 controls from the UK examined SNPs in five cytokines, including *VEGF* -1154 G > A, and showed that -1154 AA genotype had a reduced risk of prostate cancer (OR = 0.45; ref. 53). Another small case-control study on the *VEGF* -1498C > T germline SNP in 96 prostate cancer patients and 119 controls of Taiwanese descent found that carriers of -1498T allele were at 2.3-times higher risk of prostate cancer than noncarriers (54). However, this SNP was not related with prognosis (i.e., clinical stage and pathological tumor grade) and response to hormonal therapy (54). A case-control study in Tunisian subjects evaluated the influence of germline -1154G > A, -634G > C, and 936C > T *VEGF* SNPs on risk and prognosis of prostate cancer in 101 patients and 100 age-matched healthy controls (55). Carriers of at least one -1154 A allele were at reduced risk (OR = 0.42) and lower susceptibility to high-grade (OR = 0.25) and advanced prostate cancer (OR = 0.37). The OR for -1154 GA and AA genotypes were 0.46 and 0.27, respectively (55), agreeing with McCarron and colleagues (53), despite differences in ethnicity. The *VEGF* -634 GC+CC genotypes were at increased risk of prostate cancer (OR = 1.95), and the -634GC and CC genotypes were related with high-grade tumors (OR = 3.83 and 4.89, respectively; ref. 55). Haplotype analysis showed that -1154A/-634G had reduced disease risk (OR = 0.48), high-grade tumor (OR = 0.22), and advanced disease (OR = 0.45) compared with -1154G/-634G (55). No correlation was observed between 936C > T SNP and risk or aggressiveness of tumor (55). These results suggest that both *VEGF* -1154G > A and -634G > C SNPs may act as an important genetic marker of risk and aggressiveness of prostate cancer in Tunisian subjects. A case-control study in 133 sporadic prostate cancer patients and 157 healthy controls of Turkish descent evaluated the germline -1498C > T SNP and found no relation with susceptibility to disease (56), which was contrary to the findings for Taiwanese subjects (54). Further, the -1498C > T SNP was not related with the tumor-node-metastasis (TNM) staging, Gleason scores, and serum prostate specific antigen (PSA) levels of patients (56). The impact of the same germline *VEGF* -1498C > T SNP

Table 5. Summary of case-control studies assessing the VEGF polymorphisms in relation to risk of prostate cancer

Polymorphism	Ethnicity	Case-Control	Genotype	Association of Cancer Risk with Respect to VEGF Genotype	OR	L95% CI	U95% CI	OR Adjusted	L95% CI	U95% CI	Reference No.
-1498C > T*	Taiwanese	96/119	CT vs. CC	No	0.4	0.1	1.9	0.3	0.1	1.5	(54)
			TT vs. CC	No	1.4	0.3	5.9	1.1	0.2	4.9	(56)
			T vs. C	Yes (increases risk)	2.3	1.4	3.8	2.2	1.3	3.8	
	Turkish	133/157	CT vs. CC	No	1.44	0.84	2.42				(56)
			TT vs. CC	No	1.28	0.51	3.2				
			(CT or TT) vs. CC	No	1.42	0.84	2.37				
Japanese	270/252	CT vs. TT	No				0.98	0.68	1.4	(51)	
		TT vs. CC [†]	No				1.02	0.97	1.39		
		(CT or TT) vs. CC [†]	No	1.03	0.57	1.88					
-1154G > A [§]	Tunisian	101/100	GA vs. GG	Yes (decreases risk)	0.46	0.24	0.87				(55)
			AA vs. GG	Yes (decreases risk)	0.27	0.08	0.83				
			(GA or AA) vs. GG	Yes (decreases risk)	0.42	0.23	0.76				
			(GA or GG) vs. AA [†]	No	0.39	0.12	1.14				
	UK Caucasian	238/263	GA vs. GG [‡]	No	1.05	0.72	1.55				(53)
			(GA or GG) vs. AA	Yes (decreases risk)	0.45	0.24	0.86				
-634G > C	Tunisian	101/100	GC vs. GG	No	1.88	0.98	3.62				(55)
			CC vs. GG	No	2.28	0.82	6.38				
			(GC or CC) vs. GG	Yes (increases risk)	1.95	1.04	3.65				
	Austrian	702/702	CC vs. GG [‡]	No	1.08	0.86	1.35				(58)
			CT vs. CC	No	0.68	0.33	1.37				
			TT vs. CC	No	1.82	0.13	51.93				
936C > T	Tunisian	101/100	(CT or TT) vs. CC	No	0.72	0.36	1.43				(55)
			TT vs. CC [†]	No	1.16	0.91	1.48				
			CC vs. AA [†]	No	0.96	0.75	1.24				
-2578C > A	Austrian	702/702	CC vs. AA [†]	No	0.96	0.75	1.24				(58)
-2489C > T	Austrian	702/702	CC vs. TT [‡]	No	0.97	0.75	1.25				(58)
-1498C > T	Austrian	702/702	CC vs. TT [‡]	No	1	0.77	1.28				(58)
-7C > T	Austrian	702/702	CC vs. TT [‡]	No	1.09	0.86	1.39				(58)
1612G > A	Austrian	702/702	GG vs. AA [‡]	No	0.95	0.74	1.23				(58)
-1154G/-634C	Tunisian	101/100	Ref:	No	1.12	0.67	1.88				(55)
-1154A/-634G	Tunisian		-1154G/-634G	Yes (decreases risk)	0.48	0.28	0.84				
-1154A/-634C	Tunisian			No	0.91	0.28	2.99				

Abbreviations: L, lower; U, upper; Ref, reference haplotype.

*Test for homogeneity of OR: pooled OR (95% CI); CT versus CC, $P = 0.069$, 0.80 (0.18–3.54); TT versus CC, $P = 0.90$, 1.05 (0.78–1.40); (CT or TT) versus CC, $P = 0.42$, 1.24 (0.84–1.83).

[†]OR inverted from original publication.

[‡]Calculated OR.

[§]Test for homogeneity of OR: pooled OR (95% CI); GA versus GG, $P = 0.029$, 0.72 (0.32–1.63); AA versus GG, $P = 0.41$, 0.40 (0.22–0.72); (GA or AA) versus GG, $P = 0.030$, 0.64 (0.30–1.38).

in Japanese population was examined in a case-control study enrolling 270 patients and 252 controls. Similar to the Turkish population (56), the -1498C > T SNP was not related with disease risk, tumor grade and stage, Gleason scores, VEGF expression, MVD, and age at diagnosis (51). The rate of PSA relapse after radical prostatectomy was

higher in patients with the 1498CC or CT genotype (51), but the TT genotype had significantly worse cancer-specific and overall survival (51).

Another study in Tunisian subjects determined the combined effect of four germline SNPs that are involved in the process of angiogenesis, i.e., VEGF -1154G > A,

VEGF -634G > C, *MMP9* -1562C > T, and *TSP1* -8831A > G, in predicting the risk of prostate cancer by enrolling 101 patients and 74 controls (57). The *VEGF* -1154AA, *VEGF* -634GG, *MMP9* -1562CC, and *TSP1* -8831AA genotypes were considered as low-risk genotypes and were used as reference, and the remaining genotypes for each SNP were considered as high-risk genotypes (57). Patients with one, two, or three high-risk genotypes in the *VEGF* -1154/*VEGF* -634/*MMP9* group had an increasingly high risk of prostate cancer (OR = 2.79, 4.57, and 7.11, respectively), compared with the zero high-risk genotype group (57). In the *VEGF* -1154/634/*TSP1* group, patients carrying three high-risk genotypes had six times higher risk of prostate cancer (OR = 6.0) and were 20 times more prone to have high-grade tumors when compared with reference groups (OR = 20.75; 57). Overall, their findings show that the gene-gene interactions between high-risk angiogenic polymorphisms may increase the risk of prostate cancer and tumor aggressiveness (57). Finally, a case-control study in 702 patients and 702 age-matched healthy controls from Austria analyzed seven *VEGF* SNPs (-2578C > A, -2489C > T, -1498C > T, -634G > C, -7C > T, 936C > T, and 1612G > A) and their haplotypes, and failed to find any relation between these SNPs and the risk of prostate cancer or *VEGF* plasma levels (58).

To conclude, studies from Austria, Japan, Taiwan, Tunisia, Turkey, and the UK evaluated the importance of *VEGF* SNPs in prostate cancer. The *VEGF* -1154 A allele seems to predict the reduced risk (53, 55), which was consistent among studies as shown by meta-analysis (Table 5), in which the pooled OR (95% CI) for -1154 AA genotype was 0.40 (0.22–0.72). The *VEGF* -1154A allele also imparted protection against high-grade tumor and tumor aggressiveness (55). Except for Taiwanese subjects, in which the -1498T allele predicted high risk of prostate cancer (54), the -1498C > T SNP was not related with risk (51, 56, 58) or disease prognosis (i.e., with tumor grade, stage, Gleason scores, *VEGF* expression, and MVD; ref. 51, 56, 58). The -634C allele was predictive of increased risk and high tumor grades (55) in Tunisian subjects but had no relation with risk in Austrian subjects (58). The results of meta-analysis showed no significant association between any other *VEGF* polymorphism and prostate cancer risk or clinical outcome.

Conclusions

Many studies have investigated the role of *VEGF* polymorphisms as a genetic determinant for susceptibility and outcome of breast, prostate, NSCL, and colorectal cancer. Several polymorphisms have been reported within the promoter (-2578C > A, -2489C > T, -1498C > T, and -1154G > A), 5'-UTR (-634G > C and -7C > T), and 3'-UTR (936C > T and 1612G > A) for the *VEGF* gene. The variant allele for -1154G > A and 936C > T results in lower *VEGF* expression, whereas the variant allele for -1498C > T and -7C > T results in increased concentrations of *VEGF* mRNA (15, 59). The functional role of -2578C > A and -634G > C is not in agreement among studies, which report low or high *VEGF* production for variant alleles (15). Results

of association studies of these polymorphisms with risk and outcome in breast, prostate, NSCL, and colorectal cancer are also variable (Tables 2-5, Supplementary Table 6). For example the -936T allele has been shown to impart protection against breast cancer (22, 26, 32), but conversely, increases the risk of colorectal cancer (43). Similarly, the -634C allele was predictive of higher risk of NSCLC (39) and prostate cancer (55), but predicts reduced risk for colon cancer (45), and has no relation with risk of breast cancer (24, 26–29, 31). The lack of consensus about the role of these SNPs could be because of their linkage with other unknown functional SNPs in the *VEGF* gene or unknown SNPs in other factors of the angiogenesis pathway. Perhaps, association with haplotypes will be more predictive than individual SNPs. Increased understanding of the functionality of these polymorphisms and their role in tumor biology is essential to discern the reasons for inconsistencies in the literature (60), which may also help in designing association studies in the future. Other possible reasons are differences in study population (i.e., ethnicity and tumor type), small sample size (insufficient power), and failure to correct for multiple testing when studying the correlation with more than one SNP, which is important to avoid chance correlations. However, the likelihood of chance correlations should be assessed with consideration to the biological plausibility of any observed correlation (60).

Independently, several of these studies reported an association between *VEGF* SNPs and the risk of cancer (Tables 2-5); however, the findings from these studies seem to be inconsistent. We did meta-analysis to look at the combined results from studies addressing the same hypothesis. With the exception of *VEGF* -1154G > A, no other *VEGF* SNP was found to be related with a risk of studied tumors. The *VEGF* -1154A allele was at a significantly reduced risk of prostate cancer (OR = 0.40; 95% CI 0.22–0.72). Haplotype -2578A/-1154A/-634G and -1154A/-634G predicted a reduced risk of breast (26) and prostate cancer (55).

The *VEGF*-mediated angiogenesis pathway involves a complex signaling network regulated by multiple factors and it is likely that final outcome is governed by interaction and cross-talk between these signaling molecules and/or receptors rather than the *VEGF* gene alone. In that case, a pathway approach, which includes combined analysis of variations in different genes in the angiogenesis pathway, may provide better predictive markers. For example carriers of three high risk genotypes, i.e., *VEGF* -1154AA/*VEGF* -634GG/*MMP9*CC, had a 7.11-times higher risk of prostate cancer than the *VEGF* -1154GG/*VEGF* -634CC/*MMP9*TT group (57). Genome-wide association (GWA) studies, done to identify the candidate genes and chromosomal sites linked to cancer, showed that the locus including the *VEGF* gene was not associated with a risk of breast, NSCL, colon, and prostate cancer (61–71). However, our meta-analysis suggests that *VEGF* SNPs predict the risk of prostate cancer, indicating their role in vascular growth, at the minimum in prostate cancer; although they may not be as strongly predictive (i.e., penetrant) as other alleles in the genome. It should also be noted that there are other

limitations to the GWA approach, such as multiple testing in small populations and low-density genotyping (72, 73), which may limit the identification of true susceptibility genes.

The majority of the published studies assessing the role of VEGF polymorphisms compare the cases with controls to identify the disease markers. There is a critical need to conduct more studies evaluating the relation of genotype with drug concentrations and response. These studies are equally important and helpful for identification of genetic markers to guide the choice of treatment and supportive care specific to each individual.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Ferrara N. VEGF and the quest for tumour angiogenesis factors. *Nat Rev Cancer* 2002;2:795–803.
- Schneider BP, Sledge GW, Jr. Drug insight: VEGF as a therapeutic target for breast cancer. *Nat Clin Pract Oncol* 2007;4:181–9.
- Heist RS, Zhai R, Liu G, et al. VEGF polymorphisms and survival in early-stage non-small-cell lung cancer. *J Clin Oncol* 2008;26:856–62.
- Yamaguchi T, Bando H, Mori T, et al. Overexpression of soluble vascular endothelial growth factor receptor 1 in colorectal cancer: Association with progression and prognosis. *Cancer Sci* 2007;98:405–10.
- Ferrer FA, Miller LJ, Andrawis RI, et al. Angiogenesis and prostate cancer: *in vivo* and *in vitro* expression of angiogenesis factors by prostate cancer cells. *Urology* 1998;51:161–7.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- Pharm GKB. Available from: <http://www.pharmgkb.org/>.
- Brogan IJ, Khan N, Isaac K, Hutchinson JA, Pravica V, Hutchinson IV. Novel polymorphisms in the promoter and 5' UTR regions of the human vascular endothelial growth factor gene. *Hum Immunol* 1999;60:1245–9.
- Tischer E, Mitchell R, Hartman T, et al. The human gene for vascular endothelial growth factor. Multiple protein forms are encoded through alternative exon splicing. *J Biol Chem* 1991;266:11947–54.
- Minchenko A, Salceda S, Bauer T, Caro J. Hypoxia regulatory elements of the human vascular endothelial growth factor gene. *Cell Mol Biol Res* 1994;40:35–9.
- Schultz A, Lavie L, Hochberg I, et al. Interindividual heterogeneity in the hypoxic regulation of VEGF: significance for the development of the coronary artery collateral circulation. *Circulation* 1999;100:547–52.
- Watson CJ, Webb NJ, Bottomley MJ, Brenchley PE. Identification of polymorphisms within the vascular endothelial growth factor (VEGF) gene: correlation with variation in VEGF protein production. *Cytokine* 2000;12:1232–5.
- Renner W, Kotschan S, Hoffmann C, Obermayer-Pietsch B, Pilger E. A common 936 C/T mutation in the gene for vascular endothelial growth factor is associated with vascular endothelial growth factor plasma levels. *J Vasc Res* 2000;37:443–8.
- Koukourakis MI, Papazoglou D, Giatromanolaki A, Bougioukas G, Maltezos E, Sivridis E. VEGF gene sequence variation defines VEGF gene expression status and angiogenic activity in non-small cell lung cancer. *Lung Cancer* 2004;46:293–8.
- Pander J, Gelderblom H, Guchelaar HJ. Pharmacogenetics of EGFR and VEGF inhibition. *Drug Discov Today* 2007;12:1054–60.
- Schneider BP, Miller KD. Angiogenesis of breast cancer. *J Clin Oncol* 2005;23:1782–90.
- Linderholm B, Tavelin B, Grankvist K, Henriksson R. Vascular endothelial growth factor is of high prognostic value in node-negative breast carcinoma. *J Clin Oncol* 1998;16:3121–8.
- Foekens JA, Peters HA, Grebenchtchikov N, et al. High tumor levels of vascular endothelial growth factor predict poor response to systemic therapy in advanced breast cancer. *Cancer Res* 2001;61:5407–14.
- Gasparini G, Toi M, Gion M, et al. Prognostic significance of vascular endothelial growth factor protein in node-negative breast carcinoma. *J Natl Cancer Inst* 1997;89:139–47.
- Weidner N, Semple JP, Welch WR, Folkman J. Tumor angiogenesis and metastasis—correlation in invasive breast carcinoma. *N Engl J Med* 1991;324:1–8.
- Weidner N, Folkman J, Pozza F, et al. Tumor angiogenesis: a new significant and independent prognostic indicator in early-stage breast carcinoma. *J Natl Cancer Inst* 1992;84:1875–87.
- Krippel P, Langsenlehner U, Renner W, et al. A common 936 C/T gene polymorphism of vascular endothelial growth factor is associated with decreased breast cancer risk. *Int J Cancer* 2003;106:468–71.
- Smith KC, Bateman AC, Fussell HM, Howell WM. Cytokine gene polymorphisms and breast cancer susceptibility and prognosis. *Eur J Immunogenet* 2004;31:167–73.
- Jin Q, Hemminki K, Enquist K, et al. Vascular endothelial growth factor polymorphisms in relation to breast cancer development and prognosis. *Clin Cancer Res* 2005;11:3647–53.
- Lu H, Shu XO, Cui Y, et al. Association of genetic polymorphisms in the VEGF gene with breast cancer survival. *Cancer Res* 2005;65:5015–9.
- Kataoka N, Cai Q, Wen W, et al. Population-based case-control study of VEGF gene polymorphisms and breast cancer risk among Chinese women. *Cancer Epidemiol Biomarkers Prev* 2006;15:1148–52.
- Jacobs EJ, Feigelson HS, Bain EB, et al. Polymorphisms in the vascular endothelial growth factor gene and breast cancer in the Cancer Prevention Study II cohort. *Breast Cancer Res* 2006;8:R22.
- Balasubramanian SP, Cox A, Cross SS, Higham SE, Brown NJ, Reed MW. Influence of VEGF-A gene variation and protein levels in breast cancer susceptibility and severity. *Int J Cancer* 2007;121:1009–16.
- Langsenlehner U, Wolf G, Langsenlehner T, et al. Genetic polymorphisms in the vascular endothelial growth factor gene and breast cancer risk. The Austrian "tumor of breast tissue: incidence, genetics, and environmental risk factors" study. *Breast Cancer Res Treat* 2008;109:297–304.
- Eroglu A, Ozturk A, Cam R, Akar N. Vascular endothelial growth factor gene 936 C/T polymorphism in breast cancer patients. *Med Oncol* 2008;25:54–5.
- Schneider BP, Radovich M, Sledge GW, et al. Association of polymorphisms of angiogenesis genes with breast cancer. *Breast Cancer Res Treat* 2008;111:157–63.
- Jakubowska A, Gronwald J, Menkiszak J, et al. The VEGF_936_C>T 3'UTR polymorphism reduces BRCA1-associated breast cancer risk in Polish women. *Cancer Lett* 2008;262:71–6.
- Schneider BP, Wang M, Radovich M, et al. Association of vascular endothelial growth factor and vascular endothelial growth factor receptor-2 genetic polymorphisms with outcome in a trial of paclitaxel compared with paclitaxel plus bevacizumab in advanced breast cancer: ECOG 2100. *J Clin Oncol* 2008;26:4672–8.
- Burstein HJ, Chen YH, Parker LM, et al. VEGF as a marker for outcome among advanced breast cancer patients receiving anti-VEGF therapy with bevacizumab and vinorelbine chemotherapy. *Clin Cancer Res* 2008;14:7871–7.
- Yuan A, Yu CJ, Chen WJ, et al. Correlation of total VEGF mRNA and protein expression with histologic type, tumor angiogenesis, patient survival and timing of relapse in non-small-cell lung cancer. *Int J Cancer* 2000;89:475–83.
- Kaya A, Ciledag A, Gulbay BE, et al. The prognostic significance of vascular endothelial growth factor levels in sera of non-small cell lung cancer patients. *Respir Med* 2004;98:632–6.
- Dudek AZ, Mahaseth H. Circulating angiogenic cytokines in patients with advanced non-small cell lung cancer: correlation with treatment response and survival. *Cancer Invest* 2005;23:193–200.
- Kaya A, Oner F, Fitoz S, Erden I, Numanoglu N. Metastatic lung cancer: presenting with ocular symptoms. *Tuber Toraks* 2005;53:386–9.
- Zhai R, Liu G, Zhou W, et al. Vascular endothelial growth factor genotypes, haplotypes, gender, and the risk of non-small cell lung cancer. *Clin Cancer Res* 2008;14:612–7.
- Lurje G, Zhang W, Schultheis AM, et al. Polymorphisms in VEGF and IL-8 predict tumor recurrence in stage III colon cancer. *Ann Oncol* 2008;19:1734–41.
- Kaio E, Tanaka S, Kitadai Y, et al. Clinical significance of angiogenic

- factor expression at the deepest invasive site of advanced colorectal carcinoma. *Oncology* 2003;64:61–73.
42. Des Guetz G, Uzzan B, Nicolas P, et al. Microvessel density and VEGF expression are prognostic factors in colorectal cancer. Meta-analysis of the literature. *Br J Cancer* 2006;94:1823–32.
43. Bae SJ, Kim JW, Kang H, Hwang SG, Oh D, Kim NK. Gender-specific association between polymorphism of vascular endothelial growth factor (VEGF 936 C>T) gene and colon cancer in Korea. *Anticancer Res* 2008;28:1271–6.
44. Park HM, Hong SH, Kim JW, et al. Gender-specific association of the VEGF -2578C > A polymorphism in Korean patients with colon cancer. *Anticancer Res* 2007;27:2535–9.
45. Chae YS, Kim JG, Sohn SK, et al. Association of vascular endothelial growth factor gene polymorphisms with susceptibility and clinicopathologic characteristics of colorectal cancer. *J Korean Med Sci* 2008;23:421–7.
46. Yamamori M, Taniguchi M, Maeda S, et al. VEGF T-1498C polymorphism, a predictive marker of differentiation of colorectal adenocarcinomas in Japanese. *Int J Med Sci* 2008;5:80–6.
47. Kim JG, Chae YS, Sohn SK, et al. Vascular endothelial growth factor gene polymorphisms associated with prognosis for patients with colorectal cancer. *Clin Cancer Res* 2008;14:62–6.
48. Hofmann G, Langsenlehner U, Renner W, et al. Common single nucleotide polymorphisms in the vascular endothelial growth factor gene and colorectal cancer risk. *J Cancer Res Clin Oncol* 2008;134:591–5.
49. Zhang W, Gordon M, Press OA, et al. Cyclin D1 and epidermal growth factor polymorphisms associated with survival in patients with advanced colorectal cancer treated with Cetuximab. *Pharmacogenet Genomics* 2006;16:475–83.
50. Uthoff SM, Duchrow M, Schmidt MH, et al. VEGF isoforms and mutations in human colorectal cancer. *Int J Cancer* 2002;101:32–6.
51. Fukuda H, Tsuchiya N, Narita S, et al. Clinical implication of vascular endothelial growth factor T-460C polymorphism in the risk and progression of prostate cancer. *Oncol Rep* 2007;18:1155–63.
52. Balbay MD, Pettaway CA, Kuniyasu H, et al. Highly metastatic human prostate cancer growing within the prostate of athymic mice overexpresses vascular endothelial growth factor. *Clin Cancer Res* 1999;5:783–9.
53. McCarron SL, Edwards S, Evans PR, et al. Influence of cytokine gene polymorphisms on the development of prostate cancer. *Cancer Res* 2002;62:3369–72.
54. Lin CC, Wu HC, Tsai FJ, Chen HY, Chen WC. Vascular endothelial growth factor gene-460 C/T polymorphism is a biomarker for prostate cancer. *Urology* 2003;62:374–7.
55. Sfar S, Hassen E, Saad H, Mosbah F, Chouchane L. Association of VEGF genetic polymorphisms with prostate carcinoma risk and clinical outcome. *Cytokine* 2006;35:21–8.
56. Onen IH, Konac E, Eroglu M, Guneri C, Biri H, Ekmekci A. No association between polymorphism in the vascular endothelial growth factor gene at position -460 and sporadic prostate cancer in the Turkish population. *Mol Biol Rep* 2008;35:17–22.
57. Sfar S, Saad H, Mosbah F, Chouchane L. Combined effects of the angiogenic genes polymorphisms on prostate cancer susceptibility and aggressiveness. *Mol Biol Rep* 2009;36:37–45.
58. Langsenlehner T, Langsenlehner U, Renner W, et al. Single nucleotide polymorphisms and haplotypes in the gene for vascular endothelial growth factor and risk of prostate cancer. *Eur J Cancer* 2008;44:1572–6.
59. Yamamori M, Sakaeda T, Nakamura T, et al. Association of VEGF genotype with mRNA level in colorectal adenocarcinomas. *Biochem Biophys Res Commun* 2004;325:144–50.
60. Daly AK, Day CP. Candidate gene case-control association studies: advantages and potential pitfalls. *Br J Clin Pharmacol* 2001;52:489–99.
61. Amos CI, Wu X, Broderick P, et al. Genome-wide association scan of tag SNPs identifies a susceptibility locus for lung cancer at 15q25.1. *Nat Genet* 2008;40:616–22.
62. Easton DF, Eeles RA. Genome-wide association studies in cancer. *Hum Mol Genet* 2008;17:R109–15.
63. Huang YT, Heist RS, Chirieac LR, et al. Genome-wide analysis of survival in early-stage non-small-cell lung cancer. *J Clin Oncol* 2009;27:2660–7.
64. Kim TM, Yim SH, Lee JS, et al. Genome-wide screening of genomic alterations and their clinicopathologic implications in non-small cell lung cancers. *Clin Cancer Res* 2005;11:8235–42.
65. Kronenberg F. Genome-wide association studies in aging-related processes such as diabetes mellitus, atherosclerosis and cancer. *Exp Gerontol* 2008;43:39–43.
66. Ripperger T, Gadzicki D, Meindl A, Schlegelberger B. Breast cancer susceptibility: current knowledge and implications for genetic counselling. *Eur J Hum Genet* 2009;17:722–31.
67. Roukos DH. Genome-wide association studies: how predictable is a person's cancer risk? *Expert Rev Anticancer Ther* 2009;9:389–92.
68. Seng KC, Seng CK. The success of the genome-wide association approach: a brief story of a long struggle. *Eur J Hum Genet* 2008;16:554–64.
69. Spinola M, Leoni VP, Galvan A, et al. Genome-wide single nucleotide polymorphism analysis of lung cancer risk detects the KLF6 gene. *Cancer Lett* 2007;251:311–6.
70. Tenesa A, Dunlop MG. New insights into the aetiology of colorectal cancer from genome-wide association studies. *Nat Rev Genet* 2009;10:353–8.
71. Wang Y, Broderick P, Webb E, et al. Common 5p15.33 and 6p21.33 variants influence lung cancer risk. *Nat Genet* 2008;40:1407–9.
72. Ambrosone CB. The promise and limitations of genome-wide association studies to elucidate the causes of breast cancer. *Breast Cancer Res* 2007;9:114.
73. Hardy J, Singleton A. Genomewide association studies and human disease. *N Engl J Med* 2009;360:1759–68.

Molecular Cancer Therapeutics

The role of vascular endothelial growth factor SNPs as predictive and prognostic markers for major solid tumors

Lokesh Jain, Craig A. Vargo, Romano Danesi, et al.

Mol Cancer Ther 2009;8:2496-2508. Published OnlineFirst September 15, 2009.

Updated version Access the most recent version of this article at:
doi:[10.1158/1535-7163.MCT-09-0302](https://doi.org/10.1158/1535-7163.MCT-09-0302)

Supplementary Material Access the most recent supplemental material at:
<http://mct.aacrjournals.org/content/suppl/2009/09/16/1535-7163.MCT-09-0302.DC1>

Cited articles This article cites 72 articles, 19 of which you can access for free at:
<http://mct.aacrjournals.org/content/8/9/2496.full#ref-list-1>

Citing articles This article has been cited by 7 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/8/9/2496.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/8/9/2496>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.