

Research Article

Delivery of 5-Azacytidine to Human Cancer Cells by Elaidic Acid Esterification Increases Therapeutic Drug Efficacy

Bodo Brueckner¹, Maria Rius¹, Maria Rivera Markelova², Iduna Fichtner³, Petter-Arnt Hals⁴, Marit Liland Sandvold⁴, and Frank Lyko¹

Abstract

Azacytidine is an established nucleoside drug that is well known for its ability to modulate epigenetic gene regulation by inhibition of DNA methylation. Despite recent advances in the clinical development of azacytidine, the use of the drug is limited by its low bioavailability and dependency on variably expressed nucleoside transporters for cellular uptake. We show here that CP-4200, an elaidic acid derivative of azacytidine, has strong epigenetic modulatory potency in human cancer cell lines, as evidenced by efficient depletion of DNA methyltransferase protein, genome-wide DNA demethylation, and robust reactivation of epigenetically silenced tumor suppressor genes. Importantly, however, the cellular uptake of CP-4200 was substantially less dependent on the nucleoside transporters that are known to be involved in azacytidine uptake. In agreement with this notion, CP-4200 showed a significantly higher antitumoral activity than azacytidine in an orthotopic mouse tumor model for acute lymphocytic leukemia. Together, these data represent a detailed characterization of the CP-4200 mode of action and suggest that elaidic acid modification improves the therapeutic efficacy of azacytidine. *Mol Cancer Ther*; 9(5); 1256–64. ©2010 AACR.

Introduction

Azacytidine (5-azacytidine) and decitabine (2'-deoxy-5-azacytidine) are two closely related cytidine analogues that have found increasing clinical use for cancer therapy (1). Both compounds are prodrugs that need to be metabolized to the corresponding nucleoside triphosphates before they can become incorporated into DNA and/or RNA (2). After incorporation into DNA, the fraudulent base acts as a suicide substrate for DNA methyltransferase (DNMT) enzymes. The formation of a covalent reaction intermediate between DNA and enzymes triggers the depletion of DNMTs from drug-treated cells. Continuing replication in the absence of functional DNMT enzymes causes progressive loss of methylation and thus provides the mechanistic foundation for the prominent demethylating activity of azanucleoside drugs.

Drug-induced DNA demethylation represents a central aspect of epigenetic cancer therapy (3). Human cancers are characterized by a substantial number of epigenetic mutations, including hypermethylated and epigenetically silenced tumor suppressor genes (4). Because of the close

association between DNA demethylation and gene reactivation (5), it is thus generally assumed that azacytidine has the ability to revert epigenetic mutations and thereby restore normal gene expression patterns. As such, azacytidine (like decitabine) represents an archetypical epigenetic drug.

Azacytidine is approved for the treatment of myelodysplastic syndrome and has also been used successfully for the clinical management of acute myeloid leukemia (6, 7). Substantial clinical responses are usually observed at comparably low doses, roughly an order of magnitude below the maximum tolerated doses (1). These doses are generally believed to minimize the toxicity of the drug and, at the same time, maximize demethylation responses. Indeed, it has been shown recently that azacytidine can induce pronounced DNA demethylation in patients when administered under the standard prescription schedule (7–9).

The clinical response rates to azanucleosides are usually in the range of 30% to 50%. The underlying reasons for clinical azacytidine resistances are only beginning to be investigated (10) and the complex mode of action suggests that a variety of cellular pathways can modulate the efficacy of these drugs (2). One pathway that has received particular attention is the deactivation of the compound, both by chemical hydrolysis and concomitant ring opening (11) and by enzymatic deamination (12), resulting in low plasma stability. A second pathway that is being investigated is the dependency on specific and variably expressed nucleoside transporters for cellular uptake (13). Factors associated with these pathways might be important mediators of clinical drug responses.

Authors' Affiliations: ¹Division of Epigenetics, German Cancer Research Center, Heidelberg, Germany; ²Experimental Pharmacology & Oncology GmbH; ³Max Delbrück Center for Molecular Medicine, Berlin, Germany; and ⁴Clavis Pharma, Oslo, Norway

Corresponding Author: Frank Lyko, Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 580, 69120 Heidelberg, Germany. Phone: 49-6221-423800; Fax: 49-6221-423802. E-mail: f.lyko@dkfz.de

doi: 10.1158/1535-7163.MCT-09-1202

©2010 American Association for Cancer Research.

Recently, two independent strategies have been used to improve the efficacy of azanucleosides. S110, an azadeoxycytidine-deoxyguanine dinucleotide, showed cellular demethylation activity that was similar to that of azacytidine (14). However, the dinucleotide was largely resistant to deamination by cytidine deaminase (14), a process that has been tightly linked to azacytidine instability (12). In an independent approach, cytidine analogues have been modified by esterification with a fatty acid. In this context, it has been shown that elaidic acid esterification of cytarabine, a cytosine analogue closely related to azacytidine, modified the cellular uptake mechanisms of the drug and rendered it more active in human solid tumor models (15–17). Because the drug must enter into cells to interact with its target, the knowledge of the transport mechanisms is a key aspect for understanding its extended mode of action. Two major families of human nucleoside transporters have been identified: the concentrative and the equilibrative nucleoside transporters (hCNTs/SLC28 and hENTs/SLC29 proteins). The members of these families play a major role in the cellular uptake and therapeutic efficacy of nucleoside analogues (18).

In this study, we have characterized the effects of CP-4200, an elaidic acid ester derivative of azacytidine (Fig. 1A). Our results show that CP-4200 effectively depletes DNMT1 in all human cancer cell lines analyzed. Drug-induced DNA demethylation was observed both at the whole-genome level and at specific genes and was accompanied by robust reactivation of epigenetically silenced tumor suppressor genes. Whereas these results underscored profound similarities of CP-4200 with the parent drug azacytidine, the cellular uptake mechanisms of these two drugs seemed fundamentally different. In addition, CP-4200 showed a significantly improved *in vivo*


efficacy when compared with azacytidine. Taken together, these results show a potent epigenetic regulatory activity for CP-4200 and suggest that elaidic acid esterification of azacytidine increases the therapeutic efficacy of the drug.

Materials and Methods

Cell culture. HCT116 (from Bert Vogelstein, Johns Hopkins University, Baltimore, MD), MCF-7 (American Type Culture Collection), and HL60 (American Type Culture Collection) cells were cultured under standard conditions in McCoy's 5a medium, DMEM, and RPMI 1640, respectively, supplemented with 5% L-glutamine and 10% FCS (Invitrogen). To analyze the effect of CP-4200 and 5-azacytidine, cells were cultivated in media supplemented with the compounds, as indicated. To determine IC₅₀ concentrations for CP-4200 and 5-azacytidine, cell viability was analyzed after 72 hours of drug incubation by using the CellTiter-Glo luminescent viability assay (Promega) according to the manufacturer's protocol. Madin-Darby strain II canine kidney (MDCK; from Kai Simons, European Molecular Biology Laboratory, Heidelberg, Germany) cells and MDCK cells stably expressing recombinant hCNT1 (MDCK-CNT1) or hCNT3 (MDCK-CNT3) were grown in MEM (Sigma) supplemented with 10% (v/v) fetal bovine serum and 100 units/mL penicillin/streptomycin, as described previously (13).

DNMT trapping assay. Cells were treated with cell line-specific IC₅₀ concentrations for the indicated time periods. Proteins were extracted using lysis buffer [PBS (pH 7.5), 1% Igepal 630, 0.1% SDS, and 0.5% sodium deoxycholate] containing 1× complete protease inhibitor cocktail (Roche) and incubated on ice for 40 minutes. The lysate was centrifuged at 4°C for 25 minutes at

Figure 1. CP-4200 causes DNMT1 depletion. A, structure of CP-4200. B, DNMT1 trapping assay. Cells were treated with cell line-specific IC₅₀ concentrations of CP-4200 or azacytidine (AZA) for 1 to 72 h. Protein extracts were probed for the presence of DNMT1 and GAPDH by Western blotting. co, nontreated control cells.



14,000 rpm. Cleared supernatants were frozen in liquid nitrogen and stored at -80°C . Equal amounts of protein were separated on 10% SDS-polyacrylamide gels and analyzed by Western blotting using standard procedures. The primary antibodies used were anti-DNMT1 (C17, 1:500, Santa Cruz Biotechnology) and anti-glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*; 6C5, 1:1,000, Santa Cruz Biotechnology). Primary antibodies were visualized by enhanced chemiluminescence (Perkin-Elmer) according to the manufacturer's protocol.

DNA methylation analysis. Genomic DNA was purified using the DNeasy Blood and Tissue Kit (Qiagen). Global DNA methylation levels were determined by capillary electrophoresis as described previously (19). HumanMethylation27 DNA Analysis BeadChip arrays (Illumina) were used for genome-wide methylation pattern analysis (20) according to the manufacturer's instructions. Briefly, bisulfite-converted DNA was hybridized to methylation-specific oligonucleotides and washed. The hybridized oligonucleotides were then minisequenced with hapten-labeled dideoxynucleotides. After staining, the BeadChip was scanned. Results were analyzed with BeadStudio software (Illumina); histograms and heatmaps were drawn with R (<http://www.R-project.org>).

RNA isolation and reverse transcription-PCR. RNA was isolated using the RNeasy RNA Isolation kit (Qiagen). cDNA was prepared with the Quantitect Reverse Transcription kit (Qiagen). For quantitative reverse transcription-PCR (RT-PCR), 1 μg of RNA was reverse transcribed according to the manufacturer's protocol. Each cDNA sample was analyzed in triplicate using the QuantiFast SYBR Green PCR kit (Qiagen). A Quantitect Primer Assay (Qiagen) was used for tissue inhibitor of metalloproteinase-3 (*TIMP-3*; Hs_TIMP3_1_SG, QT00046382). The *DAPK1* gene was amplified using the primers *DAPK1_fwd* (TCCAAAGAAGAGAATGACAATTCA) and *DAPK1_rev* (TGCCTGTTGTGTATCTTTAGGC). The housekeeping gene *GAPDH* was amplified using the primers *GAPDH_fwd* (ATTCCACCCATGGCAAATTC) and *GAPDH_rev* (TCTCGCTCCTGGAAGATGGT). Using *GAPDH* as a reference, relative quantification was done using Q-Gen software (21). Primer-specific efficiencies were calculated from serial cDNA dilutions using the dilution method of quantification on the LightCycler (Roche) instrument.

tRNA methylation analysis. RNA bisulfite sequencing of tRNA^{Asp} was done as described previously (22). For 454 sequencing, bisulfite sequence-specific PCR amplicons were gel purified and reamplified using primer pairs containing 454 linker sequences. 454 deep sequencing of tRNA-derived cDNAs was done by Agowa (Berlin, Germany).

Transport experiments and cytotoxicity assays in MDCK cells. The sensitivity of MDCK control, MDCK-CNT1, and MDCK-CNT3 cells to CP-4200 and azacytidine, respectively, was assessed with the CellTiter-Blue assay (Promega). MDCK cells were seeded (1×10^4 per well) in 384-well plates and incubated for 24 hours before

exposure to graded concentrations of each drug for 24 or 72 hours. S-(4-Nitrobenzyl)-6-thioinosine (NBTI; Biozol) was added at a subtoxic concentration (10 $\mu\text{mol/L}$) 20 minutes before drug incubation. The IC_{50} concentration was defined as the drug concentration required to reduce cell survival, as determined by the relative fluorescence of reduced resazurin, to 50%. For transport assays, MDCK cells were seeded in six-well plates at a density of 2×10^6 per well and cultured for 24 hours to confluence. Cells were induced with 10 mmol/L sodium butyrate for additional 24 hours, as described previously (23). Uptake studies examining the Na^+ -dependent uptake of [^3H]uridine (0.6 TBq/mmol; Moravek Biochemicals) were done using Na^+ -containing buffer (100 mmol/L NaCl, 2 mmol/L KCl, 1 mmol/L CaCl_2 , 1 mmol/L MgCl_2 , and 10 mmol/L HEPES, pH 7.5) and Na^+ -free buffer in which NaCl was replaced by equimolar choline chloride. Cells were first washed three times with Na^+ -free buffer and then incubated with 1 mL of Na^+ -containing buffer containing the labeled nucleoside at the indicated concentration in the absence or presence of unlabeled nucleoside analogues for 5 minutes. After incubation at 37°C , cells were washed three times with ice-cold Na^+ -free buffer. Intracellular radioactivity was determined by liquid scintillation counting, after lysing the cells with 0.2% SDS.

In vivo experiments. Human acute lymphatic leukemia ALL-SCID 6 cells (10^7 ; ref. 24) were injected i.v. into 48 nonobese diabetic/severe combined immunodeficient female mice on day 0. Groups of eight mice were randomized to treatment and control groups, respectively. Treatment was initiated on day 3 and repeated every 3 days for five times (q3dx5). The control group received solvent only (10% Tween 80/saline, i.p.). The following treatment modalities were tested: CP-4200 in Phares solubilization—15 mg/kg/inj., i.v., or 20 mg/kg/inj., i.v.; CP-4200 in 10% Tween 80/saline—15 mg/kg/inj., i.p., or 20 mg/kg/inj., i.p.; 5-azacytidine in 10% Tween 80/saline—5 mg/kg/inj., i.v. Mice were sacrificed on day 69 and spleens were resected and weighed.

Results


Azanucleosides are generally characterized by complex modes of action that are based on cytotoxicity and epigenetic modulation of gene expression (2). To determine the effect of CP-4200 on the proliferation of human cancer cell lines, we incubated a panel of three cell lines (HCT116 colon carcinoma, MCF-7 breast cancer, and HL-60 myeloid leukemia) with increasing concentrations of CP-4200 and measured the metabolic activity of the cells after 24, 48, and 72 hours of drug incubation. The results showed well-defined dose-response curves after 48 and 72 hours and established IC_{50} concentrations of 6, 15, and 2 $\mu\text{mol/L}$ for HCT116, MCF-7, and HL-60 cells, respectively (data not shown). These concentrations were slightly higher but overall similar to the corresponding

IC₅₀ concentrations for azacytidine (5, 4, and 1 μmol/L, respectively; data not shown).

To characterize the epigenetic activity of CP-4200, we analyzed the ability of the drug to deplete the DNMT1 protein. This effect is a prominent characteristic of azanucleosides and a consequence of their incorporation into DNA. DNA containing the modified base covalently traps DNMTs and triggers the degradation of DNMT

proteins (25). HCT116, MCF-7, and HL-60 cells were incubated with the respective cell line-specific IC₅₀ concentrations of CP-4200 or azacytidine to compare the effects between individual cell lines under equitoxic conditions. Protein extracts were prepared after 24 hours and DNMT1 protein was detected by Western blotting. The results showed a rapid CP-4200-dependent depletion of DNMT1 in all cell lines, which was highly similar to the

Figure 2. CP-4200 causes global and gene-specific demethylation in cancer cells. **A**, global DNA methylation levels, as determined by capillary electrophoresis. Bars, SD from three or more measurements. **B**, array-based analysis of gene-specific DNA methylation. Histograms show the numbers of probes (y axis) over the methylation scores (x axis). Methylation scores range from 0 (completely unmethylated) to 1 (completely methylated). Blue histograms, results from untreated cells; orange histograms, methylation patterns from treated cells.



DNMT1 depletion observed with azacytidine (Fig. 1B). Differences in the kinetics of DNMT1 depletion between cell lines have been observed before and are probably related to differences in the metabolic activation of the drug (9). These results strongly suggest that intracellular CP-4200 can be metabolized to aza-dCTP and become incorporated into DNA. The observed trapping also predicted that CP-4200 has demethylating activity in human cancer cell lines.

To directly investigate the demethylating activity of the drug, HCT116, MCF-7 and HL-60 cells were incubated with 1 $\mu\text{mol/L}$ CP-4200. Cells were harvested after 24 hours, and after 72 hours, and genomic DNA was isolated. Total cytosine methylation levels were quantified by capillary electrophoresis, which established control methylation levels of 3.1% for HCT116 cells, 2.5% for MCF-7 cells, and 2.8% for HL-60 cells (Fig. 2A). These methylation levels were significantly decreased after 72 hours in all cell lines (Fig. 2A), which confirmed the demethylating activity of CP-4200. A parallel analysis of genomic DNA isolated from the same cell lines treated with 1 $\mu\text{mol/L}$ azacytidine revealed similar effects (Fig. 2A), thus suggesting that the epigenetic activity of CP-4200 is similar to that of the parent drug azacytidine.

To characterize the demethylating activity of CP-4200 in greater detail, we used Illumina HumanMethylation27 DNA Analysis BeadChip arrays to interrogate the methylation status of 27,578 CpG sites across the human genome. The methylation profiles from all three cell lines showed that the majority of sites were unmethylated, with methylation scores ranging from 0 to 0.2 (Fig. 2B). In addition, the methylation profiles also showed a second, smaller peak between 0.8 and 1, which corresponds to highly or completely methylated sites (Fig. 2B). After 72 hours of drug treatment, CP-4200 (1 $\mu\text{mol/L}$) caused a distinct shift in the hypermethylation peaks toward lower methylation scores (Fig. 2B), which illustrates the demethylation of highly methylated sites. This effect was observed in all three cell lines and was similar to the demethylation effects induced by 1 $\mu\text{mol/L}$ azacytidine (Fig. 2B). Our results thus provide an important confirmation for CP-4200-mediated DNA demethylation and suggest that the drug induces widespread demethylation at numerous hypermethylated loci. These findings also suggest that CP-4200 might be able to revert epigenetic mutations.

Hypermethylation of the *TIMP-3* tumor suppressor gene represents a well-described epigenetic mutation in HCT116 cells (26). The *TIMP-3* promoter CpG island is represented on the array by four independent probes that were strongly methylated in control HCT116 cells (Fig. 3A). Methylation was distinctly reduced after 72-hour treatment with CP-4200 (Fig. 3A). Similar results could also be observed in DNA obtained from HCT116 cells treated with azacytidine under the same conditions (Fig. 3A). We then used quantitative RT-PCR to determine the amount of *TIMP-3* mRNA in HCT116 control and drug-treated cells. The results showed that *TIMP-3* is




Figure 3. CP-4200 causes reactivation of epigenetically silenced tumor suppressor genes. **A**, array-based methylation analysis of *TIMP-3* in HCT116 cells treated with 1 $\mu\text{mol/L}$ CP-4200 or azacytidine. Numbers indicate individual methylation probes associated with the *TIMP-3* promoter (target IDs: 1-cg24080529, 2-cg05260966, 3-cg05288803, 4-cg22578204). **B**, quantitative RT-PCR analysis of *TIMP-3* expression in HCT116 cells treated with 1 $\mu\text{mol/L}$ CP-4200 or azacytidine for 24 or 72 h as indicated. **C**, array-based methylation analysis of *DAPK1* in HL-60 cells treated with 1 $\mu\text{mol/L}$ CP-4200 or azacytidine. Numbers indicate individual methylation probes associated with the *DAPK1* promoter (target IDs: 5-cg19734228, 6-cg13814950, 7-cg22571217, 8-cg15746719). **D**, quantitative RT-PCR analysis of *DAPK1* expression in HL-60 cells treated with 1 $\mu\text{mol/L}$ CP-4200 or azacytidine for 24 or 72 h as indicated. *GAPDH* expression was used as a reference and expression levels were normalized to *TIMP-3* or *DAPK1* (= 1) in control cells. Columns, mean from three experiments; bars, SE.

silenced in control HCT116 cells (Fig. 3B). The expression level was not detectably altered after 24 hours of treatment with 1 $\mu\text{mol/L}$ CP-4200, but the gene became strongly upregulated after 72 hours of drug treatment (Fig. 3B). Again, similar effects were observed with the parent drug azacytidine (Fig. 3B), and a tight correlation between *TIMP-3* demethylation and activation could be observed in all experiments.

To confirm CP-4200-mediated reversion of epigenetic mutations in an independent model, we analyzed the *DAPK-1* epimutation in HL-60 cells (27). The array contains four independent probes that are closely associated with the *DAPK1* promoter CpG, and these probes indicated high levels of methylation in control HL-60 cells (Fig. 3C). Methylation was distinctly reduced after 72-hour treatment with CP-4200 (Fig. 3C). Quantitative RT-PCR confirmed that *DAPK-1* expression was not detectable in control HL-60 cells (Fig. 3D), but robust

induction was detectable after 72 hours of drug treatment (Fig. 3D). Again, similar effects were observed with the parent drug azacytidine both by DNA methylation analysis and by expression analysis (Fig. 3C and D). Together, these results convincingly illustrate the ability of CP-4200 to revert established epigenetic mutations in human cancer cell lines.

It has been shown recently that azacytidine inhibits DNMT2-mediated tRNA methylation, and it has been suggested that this effect might be an important aspect of the antiproliferative drug effects (28). To investigate whether CP-4200 also inhibits tRNA methylation, we used RNA bisulfite sequencing of tRNA^{Asp}, an established substrate of DNMT2. Sequencing of 300 to 1,000 molecules per RNA sample showed variable CP-4200-dependent demethylation of C38, the primary target residue of DNMT2 in tRNA^{Asp} (Fig. 4). More specifically, CP-4200 treatment induced weak demethylation in HCT116 cells, but readily detectable demethylation in MCF-7 cells after 72 hours of drug incubation and in HL-60 cells after 24 hours of drug incubation (Fig. 4). These observations establish tRNA^{Asp} demethylation as a nonepigenetic molecular biomarker for CP-4200 drug activity. Similar to the effects observed with azacytidine, the kinetics of CP-4200-induced tRNA demethylation seemed to differ between individual cell lines, which is consistent with the comparably high dynamics of drug-induced tRNA demethylation (28).


It has been shown before that elaidic acid esterification of nucleoside drugs can modulate their cellular uptake mechanisms (15, 17, 29). Transport proteins that have been associated with azacytidine uptake include hCNT1 (13), hCNT3, (30) and hENT1 (31). To elucidate whether CP-4200 enters into the cells by similar pathways, we used MDCK cells, a standard model system for transport studies that is characterized by a basal Ent-dependent nucleoside drug uptake (32). When MDCK control cells were incubated with increasing concentrations of CP-4200 for 24 hours, we obtained a comparably high IC₅₀ value of 202 μmol/L, indicating low-affinity CP-

4200 uptake by Ent proteins (Fig. 5A). After 72 hours of drug incubation, the IC₅₀ value decreased to 40 μmol/L. The cytotoxicity did not depend on the basal Ent activity of MDCK control cells because the cellular drug sensitivity was increased by the addition of the ENT inhibitor NBTI (Fig. 5A). In contrast, azacytidine caused a noticeable cytotoxicity already after 24 hours of drug incubation, with an IC₅₀ value of 34 μmol/L (Fig. 5A). This effect was unchanged after 72 hours of drug incubation and the cytotoxicity decreased in cells pretreated with NBTI, with IC₅₀ values strongly increasing to 120 μmol/L (Fig. 5A). These results confirm Ent-dependent uptake of azacytidine in MDCK cells and suggest that CP-4200 is transported by different pathways.

To further characterize cellular CP-4200 uptake, we also compared the cytotoxic effects of CP-4200 and azacytidine between MDCK control cells and MDCK cells expressing recombinant hCNT1 (MDCK-CNT1). After 24 hours of drug incubation, MDCK-CNT1 cells showed a 1.9-fold reduction in the IC₅₀ value for CP-4200 (relative to MDCK control cells) but a 170-fold reduced IC₅₀ value for azacytidine (Fig. 5B). Similarly, increased cytotoxicity was also observed in MDCK-CNT3 cells with 2.7- and 170-fold reductions in the IC₅₀ values for CP-4200 and azacytidine, respectively (data not shown). These results indicate that CP-4200 is not a high-affinity substrate for hCNT1 and hCNT3 and further illustrate the differences between CP-4200 and its parent drug azacytidine in their cellular uptake mechanisms.

To further confirm these findings, we measured the uptake of 1 μmol/L [³H]uridine in MDCK-CNT1 and MDCK-CNT3 cells for 5 minutes in the absence or presence of CP-4200, azacytidine, and decitabine at 100 μmol/L concentrations (Fig. 5C). Under these experimental conditions, the nucleoside drugs compete with the radioactively labeled uridine for cellular uptake, if they share the same transport system. The results showed that azacytidine and decitabine were able to inhibit the Na⁺-dependent uptake of uridine in MDCK-CNT1 and MDCK-CNT3 after 5 minutes (Fig. 5C). In contrast, the

Figure 4. CP-4200 inhibits tRNA methylation at cytosine 38 (C38) of tRNA^{Asp}. HCT116, MCF-7, and HL-60 cells were treated with 1 μmol/L CP4200 (CP) or 5-azacytidine for 24 or 72 h. Columns, C38 methylation levels of tRNA^{Asp} as determined by 454 bisulfite sequencing. Black columns, methylation levels from CP-4200-treated cells; dark gray columns, methylation levels from azacytidine-treated cells; light gray columns, nontreated cells.






Figure 5. Cellular uptake of CP-4200. A, cytotoxicity assay in MDCK control cells. MDCK-Co cells were incubated with increasing concentrations of CP-4200 (black columns) or azacytidine (gray columns) for 24 or 72 h. NBTI was added at 10 $\mu\text{mol/L}$ concentration 20 min before and during drug incubation. The IC_{50} values were calculated as the nucleoside analogue concentration required to reduce cell viability by 50%. B, cytotoxicity assay in MDCK control and MDCK-CNT1 cells after 24-h incubation with CP-4200 and azacytidine. C, transport assay. Cells were incubated with 1 $\mu\text{mol/L}$ [^3H]uridine in Na^+ -containing or Na^+ -free buffer and in the absence (control) or presence of 100 $\mu\text{mol/L}$ CP-4200, azacytidine, and decitabine for 5 min. Na^+ -dependent intracellular substrate accumulation was calculated as mean \pm SD as determined from triplicate measurements that were reproduced independently at least once.

presence of CP-4200 did not cause inhibition of the hCNT1- and hCNT3-mediated uptake of [^3H]uridine. These results again suggest that CP-4200 is not a high-affinity substrate for hCNT1 and hCNT3 and thus further confirm that the cellular uptake mechanisms for CP-4200 and azacytidine are different.

Having shown that the elaidic acid modification of azacytidine modifies the cellular uptake mechanism of the drug without affecting its epigenetic potency, we also analyzed the therapeutic efficacy of CP-4200. To this end, an orthotopic *in vivo* tumor model was used (24). Forty-eight mice were randomized to six different treatment and control groups, respectively, and treatment was started at day 3 after *i.v.* injection of acute lymphocytic leukemia cells (Fig. 6A). In the course of the experiment, it was observed that the combination of Phares solubilization and *i.v.* application led to body weight loss, whereas the combination of Tween 80/saline solubiliza-

tion and *i.p.* administration seemed to be better tolerated (data not shown). After 69 days, mice were sacrificed and spleen weights were measured to determine the effect of the drug on splenomegaly, a prominent characteristic of the tumor model. The results showed that all four treatment modalities induced a significant decrease of the spleen weight with respect to the control group, which was independent of the route of application and dose (Fig. 6B). Notably, the decrease in splenomegaly consistently seemed to be more pronounced for CP-4200 than for an equitoxic dose of azacytidine, and the differences were statistically significant for three of four groups (Fig. 6B). These results suggest an increased therapeutic efficacy of CP-4200 when compared with azacytidine.

Discussion

Over the past few years, azacytidine has been approved for the treatment of myelodysplastic syndrome and has found increasing clinical use for the treatment of myeloid leukemias (33). Attempts to develop azacytidine for the clinical management of other tumor indications have remained largely unsuccessful. The underlying reasons are complex and include historical factors as well as specific biological features of myeloid leukemias (1). However, it is also becoming clear that efficient intracellular accumulation of azacytidine depends on the expression of specific transport proteins. As such, it is reasonable to assume that tissues with




Figure 6. Therapeutic efficacy of CP-4200 in an orthotopic mouse model of acute lymphocytic leukemia. A, treatment schedule. Eight mice per group were inoculated with 10^7 ALL-SCID 6 cells. Treatment started at day 3 and was repeated five times every 3 d. After 69 d, mice were sacrificed and spleen weights were determined. B, effect of CP-4200 on splenomegaly. * and **, $P < 0.05$, significant decreases compared with the solvent control and with both the solvent control and the azacytidine control, respectively (Mann-Whitney U test).

a low level of intracellular drug uptake and/or a high level of drug efflux are less sensitive to the effects of azacytidine.

The esterification of nucleoside drugs with fatty acids represents an established strategy for modifying the cellular uptake of these drugs. CP-4200 is an elaidic acid ester of azacytidine that has been synthesized with the intention of rendering drug uptake less dependent on conventional nucleoside transport systems. Indeed, our results indicate that the cellular uptake mechanisms of azacytidine and CP-4200 seem to be fundamentally different. More specifically, the endogenously expressed Ent proteins of MDCK cells were found to be important for the cellular uptake of azacytidine, but seemed to be largely dispensable for the uptake of CP-4200. Furthermore, our data also show that CP-4200 is not a high-affinity substrate for hCNT1 and hCNT3, whereas these proteins have been shown to efficiently mediate cellular uptake of azacytidine (13, 30). It is possible that CP-4200 enters cells through the membrane, either by a passive, concentration-dependent uptake mechanism or by an active transport pathway (34). However, alternative possibilities, including the extracellular cleavage of CP-4200 by plasma membrane lipases, followed by a secondary uptake of the free azacytidine group through hCNT1 or hCNT3 should also be considered in future studies.

Protection from inactivation by cytidine deaminase and efficient metabolic activation by cytidine kinase are important biochemical processes that modulate the potency of azacytidine (2). It was shown recently that elaidic acid esterification of the cytidine analogue gemcitabine strongly increases resistance to cytidine deaminase-mediated drug inactivation (35). As such, it is very likely that CP-4200 is a similarly poor substrate for cytidine deaminase. In addition, our results show that CP-4200 effectively depletes DNMT1 protein, which suggests that the added metabolic steps required for the activation of CP-4200 do not reduce the concentration of the active metabolite in human cancer cells. Our results also show that elaidic acid esterification does not affect the epigenetic potency of the drug. Genome-wide DNA methylation profiling of three different human cancer cell lines treated with CP-4200 showed widespread DNA demethylation of hypermethylated markers. In agreement with

this finding, CP-4200 also caused efficient reactivation of epigenetically silenced tumor suppressor genes. Overall, we noticed that the epigenetic modulatory effects of CP-4200 seemed to be similar to that of azacytidine. The low dependency on nucleoside transporters, combined with the high epigenetic modulatory potency, establishes CP-4200 as an intriguing drug candidate for epigenetic cancer therapy.

Although azacytidine has shown significant clinical benefits in the treatment of myeloid leukemias, only little clinical activity has been shown in solid tumors and in lymphocytic leukemias. A good example is provided by acute lymphocytic leukemia, in which epigenetic lesions have been found to be prevalent (36–39) but in which clinical activity of demethylating agents has only been shown for isolated cases (40). We have tested azacytidine and CP-4200 in an orthotopic ALL mouse tumor model. Our results showed a stronger therapeutic effect for CP-4200 in all treatment groups, with the difference being statistically significant in three of four groups. These results further support the notion that CP-4200 could be effective for the treatment of tumor entities that show little or no effect with azacytidine.

Disclosure of Potential Conflicts of Interest

F. Lyko and the Experimental Pharmacology & Oncology GmbH received a major commercial research grant from Clavis Pharma. No other potential conflicts of interest were disclosed.

Acknowledgments

We thank the DKFZ Genomics and Proteomics Core Facility for technical support in the Infinium methylation analysis and Bettina Helfert, Katharina Hanna, Johanna Hummel-Eisenbeiß, and Monika Becker for technical assistance.

Grant Support

Clavis Pharma. Maria Rius was supported by the Peter und Traudl Engelhorn-Stiftung zur Förderung der Biotechnologie und Gentechnik.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 12/23/2009; revised 02/13/2010; accepted 03/08/2010; published OnlineFirst 05/04/2010.

References

- Issa JP, Kantarjian HM. Targeting DNA methylation. *Clin Cancer Res* 2009;15:3938–46.
- Stresemann C, Lyko F. Modes of action of the DNA methyltransferase inhibitors azacytidine and decitabine. *Int J Cancer* 2008;123:8–13.
- Yoo CB, Jones PA. Epigenetic therapy of cancer: past, present and future. *Nat Rev Drug Discov* 2006;5:37–50.
- Esteller M. Cancer epigenomics: DNA methylomes and histone-modification maps. *Nat Rev Genet* 2007;8:286–98.
- Egger G, Liang G, Aparicio A, Jones PA. Epigenetics in human disease and prospects for epigenetic therapy. *Nature* 2004;429:457–63.
- Silverman LR, Demakos EP, Peterson BL, et al. Randomized controlled trial of azacitidine in patients with the myelodysplastic syndrome: a study of the cancer and leukemia group B. *J Clin Oncol* 2002;20:2429–40.
- Soriano AO, Yang H, Faderl S, et al. Safety and clinical activity of the combination of 5-azacytidine, valproic acid, and all-*trans* retinoic acid in acute myeloid leukemia and myelodysplastic syndrome. *Blood* 2007;110:2302–8.
- Gore SD, Baylin S, Sugar E, et al. Combined DNA methyltransferase and histone deacetylase inhibition in the treatment of myeloid neoplasms. *Cancer Res* 2006;66:6361–9.
- Stresemann C, Bokelmann I, Mahlknecht U, Lyko F. Azacytidine causes complex DNA methylation responses in myeloid leukemia. *Mol Cancer Ther* 2008;7:2998–3005.
- Qin T, Jelinek J, Si J, Shu J, Issa JP. Mechanisms of resistance

- to 5-aza-2'-deoxycytidine in human cancer cell lines. *Blood* 2009; 113:659–67.
11. Rogstad DK, Herring JL, Theruvathu JA, et al. Chemical decomposition of 5-aza-2'-deoxycytidine (decitabine): kinetic analyses and identification of products by NMR, HPLC, and mass spectrometry. *Chem Res Toxicol* 2009;22:1194–204.
 12. Chabot GG, Bouchard J, Mompalmer RL. Kinetics of deamination of 5-aza-2'-deoxycytidine and cytosine arabinoside by human liver cytidine deaminase and its inhibition by 3-deazauridine, thymidine or uracil arabinoside. *Biochem Pharmacol* 1983;32:1327–8.
 13. Rius M, Stresemann C, Keller D, et al. Human concentrative nucleoside transporter 1-mediated uptake of 5-azacytidine enhances DNA demethylation. *Mol Cancer Ther* 2009;8:225–31.
 14. Yoo CB, Jeong S, Egger G, et al. Delivery of 5-aza-2'-deoxycytidine to cells using oligodeoxynucleotides. *Cancer Res* 2007;67:6400–8.
 15. Breistol K, Balzarini J, Sandvold ML, et al. Antitumor activity of P-4055 (elaidic acid-cytarabine) compared to cytarabine in metastatic and s.c. human tumor xenograft models. *Cancer Res* 1999; 59:2944–9.
 16. Bergman AM, Kuiper CM, Voorn DA, et al. Antiproliferative activity and mechanism of action of fatty acid derivatives of arabinofuranosylcytosine in leukemia and solid tumor cell lines. *Biochem Pharmacol* 2004;67:503–11.
 17. Galmorini CM, Myhren F, Sandvold ML. CP-4055 and CP-4126 are active in ara-C and gemcitabine-resistant lymphoma cell lines. *Br J Haematol* 2009;144:273–5.
 18. Molina-Arcas M, Trigueros-Motos L, Casado FJ, Pastor-Anglada M. Physiological and pharmacological roles of nucleoside transporter proteins. *Nucleosides Nucleotides Nucleic Acids* 2008;27:769–78.
 19. Stach D, Schmitz OJ, Stilgenbauer S, et al. Capillary electrophoretic analysis of genomic DNA methylation levels. *Nucleic Acids Res* 2003;31:e2.
 20. Bibikova M, Le J, Barnes B, et al. Genome-wide DNA methylation profiling using Infinium assay. *Epigenomics* 2009;1:177–200.
 21. Muller PY, Janovjak H, Miserez AR, Dobbie Z. Processing of gene expression data generated by quantitative real-time RT-PCR. *Biotechniques* 2002;32:1372–4, 6, 8–9.
 22. Schaefer M, Pollex T, Hanna K, Lyko F. RNA cytosine methylation analysis by bisulfite sequencing. *Nucleic Acids Res* 2009;37:e12.
 23. Cui Y, Konig J, Buchholz JK, Spring H, Leier I, Keppler D. Drug resistance and ATP-dependent conjugate transport mediated by the apical multidrug resistance protein, MRP2, permanently expressed in human and canine cells. *Mol Pharmacol* 1999;55: 929–37.
 24. Borgmann A, Baldy C, von Stackelberg A, et al. Childhood ALL blasts retain phenotypic and genotypic characteristics upon long-term serial passage in NOD/SCID mice. *Pediatr Hematol Oncol* 2000;17:635–50.
 25. Weisenberger DJ, Velicescu M, Cheng JC, Gonzales FA, Liang G, Jones PA. Role of the DNA methyltransferase variant DNMT3b3 in DNA methylation. *Mol Cancer Res* 2004;2:62–72.
 26. Bachman KE, Herman JG, Corn PG, et al. Methylation-associated silencing of the tissue inhibitor of metalloproteinase-3 gene suggest a suppressor role in kidney, brain, and other human cancers. *Cancer Res* 1999;59:798–802.
 27. Raval A, Tanner SM, Byrd JC, et al. Downregulation of death-associated protein kinase 1 (DAPK1) in chronic lymphocytic leukemia. *Cell* 2007;129:879–90.
 28. Schaefer M, Hagemann S, Hanna K, Lyko F. Azacytidine inhibits RNA methylation at DNMT2 target sites in human cancer cell lines. *Cancer Res* 2009;69:8127–32.
 29. Balzarini J, Degreve B, Andrei G, et al. Superior cytostatic activity of the ganciclovir elaidic acid ester due to the prolonged intracellular retention of ganciclovir anabolites in herpes simplex virus type 1 thymidine kinase gene-transfected tumor cells. *Gene Ther* 1998;5: 419–26.
 30. Rius M, Keller D, Brom M, Hummel-Eisenbeiss J, Lyko F, Keppler D. Vectorial transport of nucleoside analogs from the apical to the basolateral membrane in double-transfected cells expressing the human concentrative nucleoside transporter hCNT3 and the export pump ABCC4. *Drug Metab Dispos* 2010. DOI:10.1124/dmd.110. 032664.
 31. Huang Y, Anderle P, Bussey KJ, et al. Membrane transporters and channels: role of the transportome in cancer chemosensitivity and chemoresistance. *Cancer Res* 2004;64:4294–301.
 32. Hammond JR, Stolk M, Archer RG, McConnell K. Pharmacological analysis and molecular cloning of the canine equilibrative nucleoside transporter 1. *Eur J Pharmacol* 2004;491:9–19.
 33. Issa JP. DNA methylation as a therapeutic target in cancer. *Clin Cancer Res* 2007;13:1634–7.
 34. Su X, Abumrad NA. Cellular fatty acid uptake: a pathway under construction. *Trends Endocrinol Metab* 2009;20:72–7.
 35. Bergman AM, Adema AD, Balzarini J, et al. Antiproliferative activity, mechanism of action and oral antitumor activity of CP-4126, a fatty acid derivative of gemcitabine, in *in vitro* and *in vivo* tumor models. *Invest New Drugs* 2010. DOI:10.1007/s10637-009-9377-7.
 36. Garcia-Manero G, Daniel J, Smith TL, et al. DNA methylation of multiple promoter-associated CpG islands in adult acute lymphocytic leukemia. *Clin Cancer Res* 2002;8:2217–24.
 37. Hoshino K, Quintas-Cardama A, Yang H, Sanchez-Gonzalez B, Garcia-Manero G. Aberrant DNA methylation of the Src kinase Hck, but not of Lyn, in Philadelphia chromosome negative acute lymphocytic leukemia. *Leukemia* 2007;21:906–11.
 38. Roman-Gomez J, Cordeu L, Agirre X, et al. Epigenetic regulation of Wnt-signaling pathway in acute lymphoblastic leukemia. *Blood* 2007;109:3462–9.
 39. Kuang SQ, Tong WG, Yang H, et al. Genome-wide identification of aberrantly methylated promoter associated CpG islands in acute lymphocytic leukemia. *Leukemia* 2008;22:1529–38.
 40. Yanez L, Bermudez A, Richard C, Bureo E, Iriando A. Successful induction therapy with decitabine in refractory childhood acute lymphoblastic leukemia. *Leukemia* 2009;23:1342–3.

Molecular Cancer Therapeutics

Delivery of 5-Azacytidine to Human Cancer Cells by Elaidic Acid Esterification Increases Therapeutic Drug Efficacy

Bodo Brueckner, Maria Rius, Maria Rivera Markelova, et al.

Mol Cancer Ther 2010;9:1256-1264. Published OnlineFirst May 4, 2010.

Updated version Access the most recent version of this article at:
doi:[10.1158/1535-7163.MCT-09-1202](https://doi.org/10.1158/1535-7163.MCT-09-1202)

Cited articles This article cites 40 articles, 18 of which you can access for free at:
<http://mct.aacrjournals.org/content/9/5/1256.full#ref-list-1>

Citing articles This article has been cited by 4 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/9/5/1256.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://mct.aacrjournals.org/content/9/5/1256>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.